

TECHNICAL REPORT

May 2017

**Dive Distribution and Group Size Parameters for Marine
Species Occurring in the U.S. Navy's Atlantic and Hawaii-
Southern California Training and Testing Study Areas**



**Naval Undersea Warfare Center Division
Newport, Rhode Island**

Suggested citation: U.S. Department of the Navy (2017). Dive Distribution and Group Size Parameters for Marine Species Occurring in the U.S. Navy's Atlantic and Hawaii-Southern California Training and Testing Areas. Technical report prepared by Naval Undersea Warfare Center, Newport

Distribution A. Unclassified. Approved for public release: distribution unlimited.

EXECUTIVE SUMMARY

An important element of the Navy's comprehensive environmental planning is the acoustic effects analysis executed with the Navy Acoustic Effects Model (NAEMO) software. NAEMO was developed to estimate the possible impacts of anthropogenic sound on marine animals, combining established acoustic propagation modeling with data regarding the distribution and abundance of marine species. This report recommends species-typical static depth distributions and group size information for all marine mammal and sea turtle species that occur in the Atlantic Fleet Training and Testing (AFTT) and Hawaii – Southern California Training and Testing (HSTT) Study Areas that will be modeled using NAEMO.

Contents

1	Introduction	1
1.1	The Navy Acoustic Effects Model	1
1.2	Data Inputs.....	1
2	Marine Mammal and Sea Turtle Depth Distributions	2
2.1	Surrogate Species and Study Areas	2
2.2	Marine Mammal and Sea Turtle Dive Behavior Summaries.....	12
2.2.1	Cetaceans.....	12
2.2.2	Carnivores	41
2.2.3	Sirenians	51
2.2.4	Sea Turtles	51
2.3	Conclusions	61
3	Marine Mammal Group Size Information.....	63
3.1	Introduction	63
3.2	Atlantic Fleet Training and Testing Study Area Group Sizes.....	63
3.3	Hawaii-Southern California Training and Testing Study Area Group Sizes	67
4	References.....	71

ACRONYMS AND ABBREVIATIONS

%	Percent
AFTT	Atlantic Fleet Training and Testing
DTAG	Digital Acoustic Recording Tag
HRC	Hawaii Range Complex
HSTT	Hawaii- Southern California Training and Testing
m	meter(s)
min	minute(s)
NAEMO	Navy Acoustic Effects Model
SD	standard deviation
U.S.	United States

1 Introduction

The United States (U.S.) Navy is required to assess potential impacts of Navy-generated sound in the water on protected marine species in compliance with applicable laws and regulations, including the National Environmental Policy Act, Executive Order 12114, the Marine Mammal Protection Act, and the Endangered Species Act. This report describes the methods and analytical approach to quantifying the depth distributions in the water column and group sizes of marine mammals and sea turtles to be used within the Navy Acoustic Effects Model (NAEMO).

1.1 The Navy Acoustic Effects Model

NAEMO is the standard model used by the Navy to estimate potential impacts to marine species from impulsive and non-impulsive sound sources used during Navy training and testing activities. NAEMO combines marine species distribution information with environmental parameters, propagation characteristics, sound source parameters, and typical training or testing scenarios in order to assess the level of behavioral disturbance, hearing impacts (including both temporary and permanent threshold shifts), and other injuries predicted for individual marine mammals and sea turtles likely to be in the vicinity of Navy training and testing activities.

1.2 Data Inputs

NAEMO first uses location-specific density (more detailed information regarding species density is available in density technical reports by U.S. Department of the Navy (In Prep.-b, In Prep.-c) and group size information to patchily distribute a given marine species into a simulation area. The depth distribution data is then used to place animals in the water column at the depths at which they are typically found. An animal is reassigned a new depth every four minutes throughout the simulation based on the depth distribution for that species. Where available, seasonal or geographically-specific depth and group size information is used.

Density data are not available for all taxa of concern for Navy activities (Section 2). In addition to available marine mammal and sea turtle data, specific information about environmental conditions and projected Navy activities within a study area is needed to run NAEMO and quantify potential impacts to marine mammals and sea turtles. This environmental data includes information about bathymetry, seafloor composition (e.g., rock, sand), and factors that vary throughout the year such as wind speed and sound velocity profiles. The details of Navy training and testing activities are also collected, including location, frequency, and source characteristics. For more detailed information about the NAEMO model, consult the Quantitative Analysis Technical Report (U.S. Department of the Navy In Prep.-a).

2 Marine Mammal and Sea Turtle Depth Distributions

The best available science from literature reviews was used to obtain species-specific depth distribution information for the Atlantic Fleet Training and Testing (AFTT) and Hawaii-Southern California Training and Testing (HSTT) Study Areas (Figure 1). Journal articles, books, technical reports, cruise reports, funding agency reports, theses, dissertations, and raw data from individual researchers were assessed for this report.

As described in Section 1.1, depth distribution data are combined with species density data during the NAEMO modeling process. Densities were obtained from the Navy Marine Species Density Database (U.S. Department of the Navy In Prep.-b, In Prep.-c). In some instances, density data were provided for guilds of species, rather than for individual species (e.g., pilot whales, *Kogia* spp., AFTT and HSTT beaked whales, AFTT seals, HSTT sea turtles, and AFTT hardshell sea turtles are all groups of two or more species). These multi-species guilds were created because observers could not differentiate closely related species at sea, or because the sample sizes of the species observed individually were too small to incorporate into density modeling. For these cases, a single representative species was chosen or a composite depth distribution was created to correspond to the multi-species groupings contained in the density data.

The information required for representing a species in NAEMO specifically focuses on the percent of time each animal spends in the water column, defined here as a range of depths extending from the surface to the maximum dive depth of each species. Percentage values may be slightly above or below 100 percent, due to decimal rounding, especially when animals spend smaller percentages of time close to their maximum dive depth. Rather than round down to zero, the deep bins are often rounded up in order to show a fraction of a percentage when it has been recorded that a species is capable of reaching that depth bin. For pinniped species, time spent hauled out of the water is not represented (this is accounted for in the density data). Depth distributions contain percent time spent in the water only, either at the surface or in given depth bins.

2.1 Surrogate Species and Study Areas

Depth distribution data within this report are based upon species-specific tagging data obtained during literature review. If tagging data were not available for a particular species, data for the most similar species were used in the form of a surrogate. A species will generally only be considered a surrogate for modeling if the species is closely related (within the same genus or family), feeds on similar prey, or has a distribution in similar water types (e.g., continental shelf waters). The exception to the general surrogate selection is the two species of *Kogia* spp. (dwarf and pygmy sperm whales), for which there are no other species in their family to choose as surrogates. Therefore, a species from another family within their suborder (Odontoceti) was chosen as a surrogate. Surrogate species (if required) for all species are provided in Table 1 for both AFTT and HSTT Study Areas.




Figure 1. Atlantic Fleet and Hawaii-Southern California Training and Testing Study Areas

Table 1. Marine Mammal and Sea Turtle Species Occurring in the Atlantic Fleet Training and Testing (AFTT) and Hawaii-Southern California Training and Testing (HSTT) Study Areas. Potential presence is indicated with an X.

Species Name	Common Name	Surrogate species	AFTT	HSTT		Section
				Southern California portion of the HSTT Study Area	Hawaii Range Complex	
Cetaceans						
Family Balaenidae						
<i>Eubalaena glacialis</i>	North Atlantic right whale	N/A	X	-	-	2.2.1.1.1
Family Balaenopteridae						
<i>Balaenoptera acutorostrata</i>	Common minke whale	N/A	X	X	X	2.2.1.2.1
<i>Balaenoptera borealis</i>	Sei whale	Bryde's whale (<i>Balaenoptera edeni</i>)	X	X	X	2.2.1.2.6
<i>Balaenoptera edeni</i>	Bryde's whale	N/A	X	X	X	2.2.1.2.5
<i>Balaenoptera musculus</i>	Blue whale	N/A	X	X	X	2.2.1.2.2
<i>Balaenoptera physalus</i>	Fin whale	N/A	X	X	X	2.2.1.2.3
<i>Megaptera novaeangliae</i>	Humpback whale	N/A	X	X	X	2.2.1.2.4
Family Delphinidae						
<i>Delphinus capensis</i>	Long-beaked common dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	-	X	-	2.2.1.3.14, 2.2.1.3.13
<i>Delphinus delphis</i>	Short-beaked common dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X	X	-	2.2.1.3.15
<i>Feresa attenuata</i>	Pygmy killer whale	Risso's dolphin (<i>Grampus griseus</i>)	X	X	X	2.2.1.3.10
<i>Globicephala macrorhynchus</i>	Short-finned pilot whale	N/A	Modeled as <i>Globicephala</i> spp.	X	X	2.2.1.3.5
<i>Globicephala melas</i>	Long-finned pilot whale	N/A	Modeled as <i>Globicephala</i> spp.	-	-	2.2.1.3.6

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Species Name	Common Name	Surrogate species	AFTT	HSTT		Section
				Southern California portion of the HSTT Study Area	Hawaii Range Complex	
<i>Globicephala</i> spp.	Pilot whales (long-finned and short-finned pilot whales)	Short-finned pilot whale (<i>Globicephala macrorhynchus</i>)	X	-	-	2.2.1.3.7
<i>Grampus griseus</i>	Risso's dolphin	N/A	X	X	X	2.2.1.3.9
<i>Lagenodelphis hosei</i>	Fraser's dolphin	Short-finned pilot whale (<i>Globicephala macrorhynchus</i>)	X	-	X	2.2.1.3.8
<i>Lagenorhynchus acutus</i>	Atlantic white-sided dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X	-	-	2.2.1.3.16
<i>Lagenorhynchus albirostris</i>	White-beaked dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X	-	-	2.2.1.3.17
<i>Lagenorhynchus obliquidens</i>	Pacific white-sided dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	-	X	-	2.2.1.3.18
<i>Lissodelphis borealis</i>	Northern right whale dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	-	X	-	2.2.1.3.19
<i>Orcinus orca</i>	Killer whale	N/A	X	X	X	2.2.1.3.1
<i>Peponocephala electra</i>	Melon-headed whale	Risso's dolphin (<i>Grampus griseus</i>)	X	-	X	2.2.1.3.11
<i>Pseudorca crassidens</i>	False killer whale	Risso's dolphin (<i>Grampus griseus</i>)	X	-	X	2.2.1.3.12
<i>Stenella attenuata</i>	Pantropical spotted dolphin	N/A	X	-	X	2.2.1.3.13
<i>Stenella clymene</i>	Clymene dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X	-	-	2.2.1.3.20
<i>Stenella coeruleoalba</i>	Striped dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X	X	X	2.2.1.3.20

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Species Name	Common Name	Surrogate species	AFTT	HSTT		Section
				Southern California portion of the HSTT Study Area	Hawaii Range Complex	
<i>Stenella frontalis</i>	Atlantic spotted dolphin	N/A	X	-	-	2.2.1.3.2
<i>Stenella longirostris</i>	Spinner dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X	-	X	2.2.1.3.21
<i>Steno bredanensis</i>	Rough-toothed dolphin	N/A	X	-	X	2.2.1.3.3
<i>Tursiops truncatus</i>	Common bottlenose dolphin	N/A	X	X	X	2.2.1.3.4
Family Eschrichtiidae						
<i>Eschrichtius robustus</i>	Gray whale	N/A	-	X	-	2.2.1.4.1
Family Kogiidae						
<i>Kogia breviceps</i>	Pygmy sperm whale	Short-finned pilot whale (<i>Globicephala macrorhynchus</i>)	Modeled as <i>Kogia</i> spp. guild	Modeled as <i>Kogia</i> spp. guild	X	2.2.1.5.1
<i>Kogia sima</i>	Dwarf sperm whale	Short-finned pilot whale (<i>Globicephala macrorhynchus</i>)	Modeled as <i>Kogia</i> spp. guild	Modeled as <i>Kogia</i> spp. guild	X	2.2.1.5.2
<i>Kogia</i> spp. (<i>Kogia breviceps</i> and <i>Kogia sima</i>)	<i>Kogia</i> spp.	Short-finned pilot whale (<i>Globicephala macrorhynchus</i>)	X	X	-	2.2.1.5.3
Family Phocoenidae						
<i>Phocoena phocoena</i>	Harbor porpoise	N/A	X	-	-	2.2.1.6.1
<i>Phocoenoides dalli</i>	Dall's porpoise	N/A	-	X	-	2.2.1.6.2
Family Physeteridae						
<i>Physeter macrocephalus</i>	Sperm whale	N/A	X	X	X	2.2.1.7.1
Family Ziphiidae						
<i>Berardius bairdii</i>	Baird's beaked whale	Cuvier's beaked whale (<i>Ziphius cavirostris</i>)	-	X	-	2.2.1.8.6

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Species Name	Common Name	Surrogate species	AFTT	HSTT		Section
				Southern California portion of the HSTT Study Area	Hawaii Range Complex	
<i>Hyperoodon ampullatus</i>	Northern bottlenose whale	Cuvier's beaked whale (<i>Ziphius cavirostris</i>)	X	-	-	2.2.1.8.7
<i>Indopacetus pacificus</i>	Longman's beaked whale	Blainville's beaked whale (<i>Mesoplodon densirostris</i>)	-	-	X	2.2.1.8.2
<i>Mesoplodon densirostris</i>	Blainville's beaked whale	N/A	-	Modeled as HSTT beaked whale guild	X	2.2.1.8.1
<i>Ziphius cavirostris</i>	Cuvier's beaked whale	N/A	-	X	X	2.2.1.8.5
<i>Mesoplodon</i> spp. (<i>M. densirostris</i> , <i>M. ginkgodens</i> , <i>M. stejnegeri</i> , <i>M. carlhubbsi</i> , <i>M. perrini</i> , <i>M. peruvianis</i>)	HSTT beaked whale guild (Blainville's beaked whale, ginkgo-toothed beaked whale, Stejneger's beaked whale, Hubb's beaked whale, Perrin's beaked whale, and pygmy beaked whale)	Blainville's beaked whale (<i>Mesoplodon densirostris</i>)	-	-	X	2.2.1.8.3

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Species Name	Common Name	Surrogate species	AFTT	HSTT		Section
				Southern California portion of the HSTT Study Area	Hawaii Range Complex	
<i>Ziphius cavirostris</i> , <i>Mesoplodon densirostris</i> , <i>M. bidens</i> , <i>M. europaeus</i> , <i>M. mirus</i>	AFTT beaked whale guild (Cuvier's beaked whale, Blainville's beaked whale, Sowerby's beaked whale, Gervais' beaked whale, and True's beaked whale)	Blainville's beaked whale (<i>Mesoplodon densirostris</i>)	X	-	-	2.2.1.8.4
Carnivores						
Family Otariidae						
<i>Arctocephalus townsendi</i>	Guadalupe fur seal	N/A	-	X	-	2.2.2.1.1
<i>Callorhinus ursinus</i>	Northern fur seal	N/A	-	X	-	2.2.2.1.2
<i>Zalophus californianus</i>	California sea lion	N/A	-	X	-	2.2.2.1.3
Family Phocidae						
<i>Cystophora cristata</i>	Hooded seal	N/A	X	-	-	2.2.2.2.1
<i>Halichoerus grypus</i>	Grey seal	N/A	Modeled as AFTT seal guild	-	-	2.2.2.2.4
<i>Monachus schauinslandi</i>	Hawaiian monk seal	N/A	-	-	X	2.2.2.2.2
<i>Mirounga angustirostris</i>	Northern elephant seal	N/A	-	X	-	2.2.2.2.3
<i>Pagophilus groenlandicus</i>	Harp seal	N/A	X	-	-	2.2.2.2.7
<i>Phoca vitulina</i>	Harbor seal	N/A	Modeled as AFTT seal guild	X	-	2.2.2.2.5

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Species Name	Common Name	Surrogate species	AFTT	HSTT		Section
				Southern California portion of the HSTT Study Area	Hawaii Range Complex	
<i>Pusa hispida</i>	Ringed seal	N/A	X	-	-	2.2.2.2.8

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Species Name	Common Name	Surrogate species	AFTT	HSTT		Section
				Southern California portion of the HSTT Study Area	Hawaii Range Complex	
<i>Halichoerus grypus</i> , <i>Phoca vitulina</i>	AFTT Seal guild (harbor seal and grey seal)	Composite of grey seal and harbor seal	X	-	-	2.2.2.2.6
Sirenians						
Family Trichechidae						
<i>Trichechus manatus</i>	West Indian manatee	N/A	X	-	-	2.2.3.1.1
Sea Turtles						
Family Dermochelyidae						
<i>Dermochelys coriacea</i>	Leatherback sea turtle	N/A	X	-	Modeled as HSTT sea turtle guild	2.2.4.1.1
Family Cheloniidae						
<i>Chelonia mydas</i>	Green sea turtle	N/A	Modeled as AFTT hardshell sea turtle guild	X	Modeled as HSTT sea turtle guild	2.2.4.2.1
<i>Eretmochelys imbricata</i>	Hawksbill sea turtle	N/A	Modeled as AFTT hardshell sea turtle guild	-	Modeled as HSTT sea turtle guild	2.2.4.2.5
<i>Caretta caretta</i>	Loggerhead sea turtle	N/A	X	-	Modeled as HSTT sea turtle guild	2.2.4.2.6
<i>Lepidochelys kempii</i>	Kemp's ridley sea turtle	Green sea turtle (<i>Chelonia mydas</i>)	X	-	-	2.2.4.2.2
<i>Lepidochelys olivacea</i>	Olive ridley sea turtle	N/A	Modeled as AFTT hardshell sea turtle guild	-	Modeled as HSTT sea turtle guild	2.2.4.2.3

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Species Name	Common Name	Surrogate species	AFTT	HSTT		Section
				Southern California portion of the HSTT Study Area	Hawaii Range Complex	
<i>Chelonia mydas</i> , <i>Lepidochelys olivacea</i> , <i>Eretmochelys imbricata</i>	AFTT hardshell sea turtle guild (green, olive ridley, hawksbill)	Composite of all unidentified sightings of all hardshell turtle species	X	-	-	2.2.4.2.4
<i>Chelonia mydas</i> , <i>Eretmochelys imbricata</i> , <i>Caretta caretta</i> , <i>Lepidochelys olivacea</i> , <i>Dermochelys coriacea</i>	HSTT sea turtle guild (green, hawksbill, loggerhead, olive ridley, leatherback)	Composite of green, hawksbill, olive ridley, loggerhead, and leatherback sea turtles	-	-	X	2.2.4.2.7

2.2 Marine Mammal and Sea Turtle Dive Behavior Summaries

This section discusses the depth distributions that were constructed for each species or surrogate species based on the best available science. Ideally, depth distributions would be specific to different locations; however, sometimes diving data were not available for the precise locations within AFTT and HSTT. Marine mammal and sea turtle dive behaviors are not easily stereotyped, but a species' behavior can generally be quantified by using an average percentage of time that an animal will typically spend within a range of depths, or depth bin. For each species, a distribution throughout the water column is presented, along with a list of the references that are the source of the data and an explanation about how these references were used to create the distribution. Depth bins are given in meters (m). Depending on the species, the distribution may cover a larger or smaller range of depths, such as for a shallow diving fur seal or a deep diving sperm whale. Likewise, depth bins may be smaller near the surface or larger at greater depths (e.g., 20 m bins near the surface where the animal spends more time or 100 m bins at the deepest depths the animal can reach). For certain species (e.g., North Atlantic right whale, humpback whale, and sperm whale), more than one depth distribution is given due to documented seasonal or geographic differences in diving behavior. Individual species are listed within each order.

2.2.1 Cetaceans

2.2.1.1 Family Balaenidae

2.2.1.1.1 *Eubalaena glacialis*, North Atlantic Right Whale

North Atlantic right whales migrate between their feeding grounds in temperate and sub-polar shelf waters and near-shore waters, and their breeding grounds in sub-tropical near-shore waters (Kenney 2002). Their prey consists entirely of zooplankton, particularly large copepods. Right whales are skim feeders that swim open-mouthed, both at the surface and at depth, through patches of congregated prey (Kenney 2002; Kenney et al. 2001; Mate and Nieukirk 1992).

Due to a separation of behaviors based on location within a foraging ground or within a breeding ground, two separate representative depth distributions were compiled for North Atlantic right whales. Foraging grounds include all waters off the North American seaboard that are north of Cape Hatteras, North Carolina. In order to build a representative depth distribution for the North Atlantic right whale on foraging grounds, depth distributions for 46 whales from the Bay of Fundy were provided by D. Nowacek and A. McGregor (2010). The Bay of Fundy data reflect skim feeding both at the surface and at depth (Nowacek and McGregor 2010). Parks et al. (2011) examined dive behavior during the spring feeding season in Cape Cod Bay (Parks et al. 2011). Parks et al. (2011) reported whales spent on average 84 percent of time within 3 m of the surface, while Nowacek and McGregor (2010) reported whales spent on average 32.89 percent of time in the top 10 m. The average of these percentages amounts to 58.45 percent of the whales' time spent in the surface bin (0-10 m). The remaining depth bins provided by Nowacek and McGregor (2010) were redistributed proportionally to account for the remaining 41.56 percent of time. The resulting depth distribution for right whales on foraging grounds is given in Table 2. This distribution is consistent with maximum dive depths reported in the literature of 174 m (Baumgartner and Mate 2003) and 272 m (Mate and Nieukirk 1992).

Table 2. Percentage of Time at Depth for the North Atlantic Right Whale on Foraging Grounds¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–10	58.45	110–120	5.19
10–20	2.39	120–130	4.79
20–30	1.66	130–140	3.88
30–40	1.22	140–150	3.31
40–50	2.08	150–160	1.94
50–60	1.07	160–170	0.84
60–70	0.97	170–180	0.28
70–80	2.29	180–190	0.02
80–90	2.33	190–200	0.02
90–100	2.62	200–210	0.01
100–110	4.65	210–220	0.01

¹Based on data from Nowacek and McGregor (2010) and Parks et al. (2011)

In order to build a representative depth distribution for North Atlantic right whales on breeding grounds, depth distributions for six whales in the South Atlantic Bight were provided by Nowacek and McGregor (2010) (Table 3). The South Atlantic Bight encompasses the continental shelf from Cape Hatteras, North Carolina south to Cape Canaveral, Florida. The depth distribution in Table 3 will also be used for both migration periods and any other movements to the south of Cape Hatteras.

Table 3. Percentage of Time at Depth for the North Atlantic Right Whale on Breeding Grounds¹

Depth Bin (m)	% of Time at Depth
0–4	46.05
4–8	29.16
8–12	9.73
12–16	14.76
16–24	0.31

¹Based on data from Nowacek and McGregor (2010)

2.2.1.2 Family Balaenopteridae

2.2.1.2.1 *Balaenoptera acutorostrata*, Common Minke Whale

Minke whales are widely distributed throughout the world oceans, occurring in coastal and continental shelf waters, the deeper waters along continental slopes, and further seaward (Dorsey et al. 1990; Øien 1990). Fish (e.g., capelin, sandlance, and herring) and planktonic crustaceans (e.g., krill) are the main components of the minke whale diet (Haug et al. 1995). Minke whales feed by side-lunging into schools of prey as well as gulping large amounts of water (Jefferson et al. 2008).

Little data have been collected on the dive behavior of minke whales. In order to build a representative depth distribution for minke whales, data from Figure 2 in Blix and Folkow (1995) were used. Blix and Folkow (1995) presented a time-depth record for a single minke whale tagged off the west coast of Svalbard, a Norwegian archipelago. This animal was predominantly foraging between 25 and 50 m. Two depth bins and the time spent within each depth bin were estimated, with the resulting depth distribution shown. The depth distribution data are derived from a short (75 minutes [min]) dive profile of a single animal, in which two behaviors are represented, cruising (52 percent of time) and foraging (48 percent of time), however, the amount of time spent in these two behaviors can vary significantly among individuals (Blix and Folkow 1995). The depth distribution for common minke whales is given in Table 4.

Table 4. Percentage of Time at Depth for the Minke Whale¹

Depth Bin (m)	% of Time at Depth
0–25	79.7
25–65	20.3

¹Based on data from Blix and Folkow (1995)

More recent data suggest that the common minke whale is capable of diving to greater depths than depicted by this distribution. For example, minke whales in the Antarctic have been associated with krill patches found at a median depth of 118 m (Friedlaender et al. 2009b). Off Scotland, minke whales are found where patches of pre-spawning herring occur at depths between 100 and 150 m (MacLeod et al. 2003b), while off the coast of California, tagged minke whales dove to 130 m (Southall et al. 2014), although in both cases whales spend the majority of time in the top 25 m of the water column. There is also limited evidence that minke whales may exhibit diurnal variation in diving behavior (Joyce et al. 1990; Stockin et al. 2001). The depth distribution shown in Table 4 will be considered representative for common minke whales until more information becomes available.

2.2.1.2.2 *Balaenoptera musculus*, Blue Whale

Blue whales have a cosmopolitan distribution, living in both coastal and offshore waters (Jefferson et al. 2008). Blue whales track the diel vertical migration of their prey and feed almost exclusively on euphausiids, or krill (Sears 2002). Although surface feeding has been observed during the daylight, it is more usual for blue whales to dive to at least 100 m into layers of euphausiid concentrations during daylight hours and feed nearer the surface at night (Sears and Perrin 2008).

In order to build a representative depth distribution for blue whales, data from Figures 4 and 8 in Oleson et al. (2007), as well as Figure 2 from Acevedo-Gutiérrez et al. (2002) were used. Oleson et al. (2007) provided graphs of the percent time at depth of 38 blue whales off the coast of California in Figure 8. The data for the non-vocal, AB callers, and D callers were averaged together to get a general depth distribution. However, percentage of time at the surface was ignored by this study. By incorporating the average number of surfacing events over time in the sample dive profile from Figure 4 in Oleson et al. (2007) and the average time spent at surfacing events from Figure 2 in the Acevedo-Gutiérrez et al. (2002) study, a percent time spent in the surface bin could be estimated (20.9 percent). The remaining bins from the Oleson et al. (2007) study were redistributed proportionally to account for the remaining 79.1 percent of time. The depth distribution for blue whales is given in Table 5.

Table 5. Percentage of Time at Depth for the Blue Whale¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–5	20.9	155–165	2.21
5–15	12.74	165–175	2.37
15–25	9.89	175–185	2.21
25–35	6.41	185–195	2.29
35–45	5.30	195–205	1.42
45–55	3.96	205–215	1.11
55–65	3.48	215–225	1.11
65–75	2.77	225–235	1.03
75–85	2.21	235–245	0.87
85–95	2.37	245–255	0.71
95–105	2.45	255–265	0.71
105–115	2.06	265–275	0.32
115–125	1.98	275–285	0.24
125–135	2.06	285–295	0.16
135–145	2.14	295–305	0.16
145–155	2.29	305–315	0.16

¹Based on data from Oleson et al. (2007) and Acevedo-Gutiérrez et al. (2002)

While other studies did not include depth distributions for blue whales, they did provide additional information to categorize dive behavior. Blue whales in the Gulf of St. Lawrence conducted foraging dives to 150 m, where feeding lunges were observed (Doniol-Valcroze et al. 2011). Similarly, a study conducted in Monterrey Bay found that blue whales fed on the most concentrated patches of krill at depths of 130 to 150 m (Schoenherr 1991). Blue whales off central California foraged at depths between 130 and 300 m (Calambokidis et al. 2008; Croll et al. 2001), while in southern California, dive depths ranged from 50 to 350 m (Acevedo-Gutiérrez et al. 2002; Croll et al. 2001; De Vos et al. 2012; Goldbogen et al. 2012; Goldbogen et al. 2013; Mate et al. 2016; Oleson et al. 2007; Southall et al. 2014). Seven whales tagged off the coast of Southern California dove to a mean depth of 140 m and a maximum depth of 204 m during foraging, while to only a mean depth of 67.6 m during non-foraging dives (Croll et al. 2001). These data are consistent with the depth distribution in Table 5.

2.2.1.2.3 *Balaenoptera physalus*, Fin Whale

The fin whale occurs in greatest concentrations in cold and temperate waters around the globe, and are commonly found seaward of the continental slope (Aguilar 2002). Prey species include euphausiids (Laidre et al. 2010; Ruchonnet et al. 2006; Vikingsson 1997), schooling fish such as herring and capelin (Nottestad et al. 2002), and cephalopods (Flinn et al. 2002). A 2001 study has shown that dense prey concentrations are typically found at depths greater than 100 m off the coast of California (Croll et al. 2001).

In order to build a representative depth distribution for fin whales, data from Figure 4a in Croll et al. (2001) and the text of Goldbogen et al. (2006) were used. Due to the lack of data on time spent at depth, the data from Croll et al. (2001) will be used as a proxy for percentage of time spent at depth. Croll et al. (2001) found that, amongst the 15 tagged fin whales, there was a maximum dive depth of 316 m. Foraging dives were deeper and longer in duration than non-feeding dives. Goldbogen et al. (2006) reported that tagged whales spent 40 percent of time in the top 50 m. Time spent at depths below 50 m were extracted from dive profiles presented in Croll et al. (2001) to represent the remaining 60 percent of time. The depth distribution for fin whales is given in Table 6.

Table 6. Percentage of Time Spent at Depth for the Fin Whale¹

Depth Bin (m)	% of Time at Depth
0–50	40
50–70	13.9
70–90	9.6
90–110	8.14
110–130	11.43
130–150	8.41
150–170	3.48
170–190	2.01
190–210	1.37
210–230	0.55
230–250	0.64
250–270	0.18
270–290	0.09
290–310	0.18

¹Based on data from Croll et al. (2001) and Goldbogen et al. (2006)

While other studies did not include depth distributions for fin whales, they did provide additional information to categorize dive behavior. Off southern California foraging dives of 100 to 300 m (Acevedo-Gutiérrez et al. 2002; Goldbogen et al. 2006; Mate et al. 2016) have been recorded. In the Ligurian Sea, a maximum dive to over 470 m was noted (Panigada et al. 1999), while Southall et al. (2014) reported that dives by fin whales rarely exceeded 250 m. These data are consistent with the depth distribution in Table 6.

2.2.1.2.4 *Megaptera novaeangliae*, Humpback Whale

Humpback whales have a cosmopolitan distribution in the coastal and continental shelf waters of the globe. They migrate between mid- to high-latitude foraging grounds and low-latitude breeding grounds (Clapham 2002). Humpback whales feed on a variety of organisms, including euphausiids and small schooling fish (Hain et al. 1982; Hazen et al. 2009; Laerm et al. 1997).

Due to a separation of behaviors based on location within a foraging ground or within a breeding ground, two separate representative depth distributions were compiled for humpback whales. In order to build a representative depth distribution for humpback whales on foraging grounds, data from Figure 3.8 as well as the text of Dietz et al. (2002) were used. Dietz et al. (2002) shows the number of dives per hour to specific depth bins. The data from Dietz et al. (2002) will be used as a proxy for percentage of time spent at depth for the six whales tagged off the foraging grounds of West Greenland. While Figure 3.8 begins at a depth of 8 m, the text states that the average time spent at the surface is 83.3 percent during mid-day and 75 percent at midnight, resulting in an average surface time of 79.2 percent. The data from Dietz et al. (2002) were then redistributed proportionally to account for the remaining 20.8 percent of time. The depth distribution for humpback whales on foraging grounds is given in Table 7.

Table 7. Percentage of Time at Depth for Humpback Whales on Foraging Grounds¹

Depth Bin (m)	% of Time at Depth
0–8	79.2
8–20	11.65
20–35	2.70
35–50	1.04
50–100	1.56
100–150	1.56
150–200	1.25
200–300	1.25
300–400	0.21

¹Based on data from Dietz et al. (2002)

Dive depths on the Greenland foraging grounds are consistent with the depth of feeding reported by Goldbogen et al. (2008) off central California and Dolphin (1987b) off Alaska. Dolphin (1987a), however, reported that 75 percent of feeding dives were to less than 60 m, and Friedlaender et al. (2009a) found evidence of bottom feeding in the shallower water (less than 50 m) of the Gulf of Maine. While the depth distribution in Table 8 has a maximum depth of 400 m, over 95 percent of the time is in the top 50 m. Therefore, the depth distribution in Table 7 is consistent with these studies as well.

Humpback whales have major breeding grounds in several locations, including the West Indies, Hawaii, Mexico, and Japan (Clapham 2002). In order to build a representative depth distribution for humpback whales on breeding grounds, data from Table 3 in Baird et al. (2000) were used. Baird et al. (2000) reported the time at depth data for 10 whales in Hawaiian waters. While all 10 whales were thought to be males, the whales were engaged in a variety of behaviors, including escorting females and calves. Therefore, the depth distribution in Table 9 represents the best estimate of time spent at depth by whales on a breeding ground. Baird et al (2000) found that, on average, about 40 percent of a whale’s time was spent in the top 10 m and about 90 percent of the time was spent in the top 100 m. The depth distribution for humpback whales on breeding grounds is given in Table 8.

Table 8. Percentage of Time at Depth for the Humpback Whale on Breeding Grounds¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–10	39.55	90–100	1.55
10–20	26.51	100–110	1.39
20–30	11.65	110–120	1.31
30–40	4.25	120–130	0.92
40–50	3.04	130–140	0.72
50–60	2.47	140–150	0.30
60–70	2.14	150–160	0.23
70–80	1.66	160–170	0.15
80–90	1.97	170–180	0.09

¹Based on data from Baird et al. (2000)

2.2.1.2.5 *Balaenoptera edeni*, Bryde’s Whale

Bryde’s whales are found in tropical and temperate waters, with separate coastal and offshore forms (Best 2001; Weir 2007). There is ongoing debate about the taxonomic relationship between two morphotypes — the larger *-brydei* form, found within the AFTT and HSTT Study Areas, and the smaller *-edeni* form, found outside of the AFTT and HSTT study areas (Sasaki et al. 2006). These two forms are genetically distinct, and

are differentiated by geographic distribution, inshore/offshore habitat preferences, and size. However, for both morphotypes, which are not easily distinguished at sea, the scientific name *B. edeni* is commonly used. The main prey of Bryde's whales include pelagic schooling fish species, such as sardines, mackerel, and herring (Siciliano et al. 2004), as well as cephalopods and small crustaceans (Kato 2002; Omura 1962).

In order to build a representative depth distribution for Bryde's whales, data from Table 1 in Alves et al. (2010) were used. Alves et al. (2010) reported a distribution of time spent in shallow versus deep dives for two whales tagged with a time-depth recorder near Madeira Island, Spain. Though these data are not strictly an indication of time spent in the two different depth bins (time spent diving to 40-292 m includes time passing through the 0-40 m depth bin), the data are the best available approximation of time spent at depth. The depth distribution for Bryde's whales is given in Table 9.

Table 9. Percentage Time at Depth for the Bryde's Whale^{1,2}

Depth Bin (m)	% of Time at Depth
0-40	84.5
40-292	15.5

¹Based on data from Alves et al. (2010)

²This depth distribution is also representative of the sei whale

2.2.1.2.6 *Balaenoptera borealis*, Sei Whale

Sei whales have a cosmopolitan distribution, migrating between high latitude feeding grounds and low latitude breeding grounds (Horwood 2002). Sei whales are capable of diving for between 5 and 20 min (Reeves et al. 2002) to feed on plankton, predominantly copepods and euphausiids, which occur between the surface and depths around 150 m (Budylenko 1978; Flinn et al. 2002). They may also feed on small schooling fish and cephalopods by both gulping and skimming.

Little data have been collected on the dive behavior of sei whales. Sei whales are not thought to be deep divers. Baumgartner et al. (2011) found that sei whales were absent during times when copepods were at depth, suggesting that sei whales may only be able to feed effectively on copepod aggregations when they are at or near the surface. In addition, Baumgartner and Fratantoni (2008) observed low calling rates during the night when copepods were at the surface, and higher calling rates during the day when copepods were at depth. This study speculated that sei whales reduced calling rates to accommodate nighttime feeding on the copepod aggregations at the surface, and increased calling rates during the day when copepods migrated to deeper depths where they were unavailable as prey to the sei whales.

Due to a lack of available data on the dive behavior of sei whales, they will be represented by a surrogate species: the Bryde's whale (Section 2.2.1.2.5). The Bryde's whale is the closest relative to the sei whale (Sasaki et al. 2005); these species are of similar body size (Horwood 2002) and feed on similar prey in the Northern Hemisphere (Flinn et al. 2002; Mizroch et al. 1984). While sei whales differ from other Balaenopterids in their prey preference for copepods, this preference means that, like Bryde's whales, sei whales are not thought to be deep divers and spend most of their time near the surface (Alves et al. 2010). Foraging sei whales and Bryde's whales utilize similar water depths (Alves et al. 2010; Baumgartner et al. 2011). The depth distribution for the sei whale can be found in Table 9.

2.2.1.3 Family Delphinidae

2.2.1.3.1 *Orcinus orca*, Killer Whale

Killer whales have a cosmopolitan distribution, but are most commonly observed in temperate, coastal waters (Ford 2002). Killer whales feed on a variety of prey, although most populations exhibit some degree of

dietary specialization. In the northeastern Pacific and Antarctic, sympatric populations in each location are socially (and, in some cases, reproductively) isolated by foraging specializations for fish or marine mammal species (Ford et al. 1998; Pitman and Ensor 2003; Saulitis et al. 2000).

Due to a separation of diving behaviors based on preferred prey of the killer whale, two separate representative depth distributions were compiled for killer whales: fish-eating killer whales and mammal-eating killer whales. Fish-eating killer whales have been studied more extensively than mammal-eating ecotypes, although there is still limited published information on diving behavior of either. Fish-eating killer whales will either chase individual prey at the surface, or collectively herd schooling fish towards the surface (Domenici et al. 2000; Nøttestad et al. 2002). Mammal-eating killer whales have different foraging strategies than fish-eating killer whales (Barrett-Lennard et al. 1996; Pitman and Ensor 2003). Mammal-eating killer whales often attempt to capture prey from below, where a prey's silhouette against brighter surface waters may improve detection. Miller et al. (2010) found deeper dives for mammal-eating killer whales occurred during the day.

In order to build a representative depth distribution for fish-eating killer whales, data from Figure 2 in Sivle et al. (2012), Figure 1e in Kvadsheim et al. (2012), as well as plots from post sonar exposure and/or silent pass of the ship from Miller et al. (2011) were used. Since all three studies analyzed the potential effects of sonar on the dive behavior of killer whales, only dive profiles from periods of time when no sonar was active were used. Depth distributions were extracted from the presented dive profiles. Individual depth distributions for each animal were averaged to create the representative depth distribution in Table 10. This representative depth distribution is consistent with Baird (1994) and Shapiro (2008), who reported that fish-eating resident killer whales spent the vast majority of their time in the top 20 m. The depth distribution for killer whales is given in Table 10. The deepest killer whale dive recorded thus far was to 264 m by Baird et al. (2005a), who tagged a total of 34 Southern Resident killer whales. However, the average of all the tagged killer whale deepest dives was 141 m, which is consistent with the depth distribution in Table 10 below.

Table 10. Percentage of Time at Depth for Fish-Eating Killer Whales¹

Depth Bin (m)	% of Time at Depth
0–20	81.51
20–40	8.60
40–60	4.45
60–80	2.37
80–100	2.12
100–120	0.53
120–140	0.32
140–150	0.18

¹Based on data from Kvadsheim et al. (2012), Miller et al. (2011), and Sivle et al. (2012)

In order to build a representative depth distribution for mammal-eating killer whales (Table 11), data from Figure 5 in Miller et al. (2010) and Figure 2.5B from Baird (1994) were used. Based on visual inspection and interpretation of the Figures provided by Miller et al. (2010), an average dive distribution was created for the presented whales; visual inspection was also used to build a depth distribution from the Figure presented by Baird (1994). The calculated data from Miller et al. (2010) was weighted by a factor of 11 when creating the final dive profile for mammal-eating killer whales, due to the presence of 11 animals in that study compared to one presented by Baird (1994).

Table 11. Percentage of Time at Depth for Mammal-Eating Killer Whales¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–10	42.91	50–60	2.46
10–20	23.32	60–70	1.66
20–30	13.94	70–80	0.82
30–40	8.27	80–100	0.36
40–50	5.97		

¹Based on data from Baird (1994) and Miller et al. (2010)

2.2.1.3.2 *Stenella frontalis*, Atlantic Spotted Dolphin

The Atlantic spotted dolphin is usually found in shallow, continental shelf waters. Atlantic spotted dolphins feed on both mesopelagic fish and squid, as well as benthic invertebrates (Perrin et al. 1994a). In the Bahamas, the Atlantic spotted dolphin diet consists of, in order of highest consumption: flying fish (Family Exocoetidae), squid (*Doryteuthis spp.*), needlefish (Family Belonidae), and ballyhoo/half beaks (Family Hemiramphidae) (Herzing and Elliser 2013).

In order to build a representative depth distribution for Atlantic spotted dolphins, data from Table 1 in Davis et al. (1996) were used. Davis et al. (1996) tracked a tagged dolphin in water with a mean depth of 32.6 m (range 12–63 m), and indicated that it was consistently diving deep enough to reach the seafloor. Davis et al. (1996) reported no diel pattern in the depth of dives. The depth distribution for Atlantic spotted dolphins is given in Table 12.

Table 12. Percentage of Time at Depth for the Atlantic Spotted Dolphin¹

Depth Bin (m)	% of Time at Depth
0–10	76.2
10–20	11.4
20–30	8.3
30–40	2.9
40–60	1.4

¹Based on data from Davis et al. (1996)

This representative depth distribution is consistent with tagging results from Griffin (2005), who reported dolphins diving to the seafloor at 30 m, with animals spending most of their time in the top 10 m. While Atlantic spotted dolphins are often observed in deeper water than 60 m (Davis et al. 1998; Herzing and Elliser 2013), Herzing and Elliser (2013) observed Atlantic spotted dolphins near Little Bahama Bank spending most of the day in shallow water and moving into deeper water when the deep scattering layer rises after dark to forage.

2.2.1.3.3 *Steno bredanensis*, Rough-toothed Dolphin

Rough-toothed dolphins are commonly found in waters along the continental shelf in tropical and warmer temperate waters (Davis et al. 1998). Rough-toothed dolphins have been reported feeding on squids and fishes near the surface, which may indicate that they primarily make shallow dives (Lodi and Hetzel 1999; Pitman and Stinchcomb 2002).

In order to build a representative depth distribution for rough-toothed dolphins, data from Table 9 in Wells et al. (2008) were used. Wells et al. (2008) reported time-at-depth data from four rehabilitated and released adult rough-toothed dolphins in the Atlantic Ocean and presented the percentage of dives to greater than 2 m for each animal. While these data underestimate surface time (since an animal had to dive below 2 m

depth for the tag to save the data), they indicate that rough-toothed dolphins spend the majority of their time in the upper 25 m of the water column. Only two of the four dolphins (three dives total) reached the 200–300 m depth bin, and dives were generally shallowest during the daytime. The data from Wells et al. (2008) were averaged across all four animals. The depth distribution for rough-toothed dolphins is given in Table 13.

Table 13. Percentage of Time at Depth for the Rough-Toothed Dolphin¹

Depth bin (m)	% of Time at Depth
0–10	78.0
10–25	16.2
25–50	3.8
50–75	0.9
75–100	0.3
100–150	0.1
150–200	0.01
200–300	0.01

¹Based on data from Wells et al. (2008)

Other than the Wells et al. (2008) study, little data have been collected on the dive behavior of the rough-toothed dolphin. An early study by Watkins et al. (1987) reported rough-toothed dolphins rubbing against a deployed hydrophone at a depth of 70 m.

2.2.1.3.4 *Tursiops truncatus*, Bottlenose Dolphin

Bottlenose dolphins have a cosmopolitan distribution in the tropical and temperate waters of the world (Wells and Scott 2002). They reside in estuarine, coastal, and offshore continental shelf and slope waters. Populations vary in their migratory and foraging behavior (Wells and Scott 2002). Bottlenose dolphins feed primarily on fish species, with squid and other invertebrates contributing to their diet as well (National Marine Fisheries Service 2015). Due to the range of habitats in which bottlenose dolphins are found, prey species may be epipelagic, pelagic, mesopelagic, or benthic in origin, depending on the region and habitat (Mead and Potter 1990; Rossbach and Herzing 1997; Shane 1990; Wells and Scott 1999). The presence of deep-sea fish in the stomachs of some offshore animals suggests that they can dive to depths greater than 500 m (Reeves et al. 2002).

Little data have been collected on the dive behavior of bottlenose dolphins. In order to build a representative depth distribution for bottlenose dolphins, data from Table 4 in Klatsky (2004) were used. Dolphins 39999 and 40000 were tagged for 30 and 48 hours, respectively, whereas Dolphin 40001 was tracked for 45 days, providing 792 hours of dive data. So, while Klatsky (2004) presents the percentage of time at depth for three individuals, only the data from Tag 40001 were considered for the depth distribution in order to provide the most comprehensive view of bottlenose dolphin diving behavior. Klatsky (2004) reported that the maximum recorded depth of a dolphin was 492 m, therefore, that used as the maximum depth associated with this depth distribution. The depth distribution for bottlenose dolphins is given in Table 14.

Table 14. Percentage of Time at Depth for the Bottlenose Dolphin¹

Depth bin (m)	% of Time at Depth
0–6	64.6
6–10	3.9
10–26	5.7
26–50	2.5
50–76	2.0
76–100	1.7
100–150	3.6
150–200	2.5
200–250	2.1
250–300	2.2
300–350	1.9
350–400	1.7
400–450	1.6
400–492	4.0

¹Based on data from Klatsky et al. (2004)

These data are consistent with animals foraging up to 500 m off Hawaii, and spending the majority of their time between the surface and 50 m (Baird et al. 2014).

2.2.1.3.5 *Globicephala macrorhynchus*, Short-finned Pilot Whale

Short-finned pilot whales occur in tropical and warm-temperate waters along the continental shelf and slope (Davis et al. 1998). Short-finned pilot whales feed pre-dominantly on squid, but they also feed on octopus and fish occasionally (Mintzer et al. 2008; Reeves et al. 2002). On the U.S. Pacific coast the neritic cephalopod, *Loligo* spp., is the dominant prey (Mintzer et al. 2008). Short-finned pilot whales feed on vertically migrating prey, diving deep during dusk and dawn and staying near surface at night (Baird et al. 2003).

In order to build a representative depth distribution for short-finned pilot whales, data from Figure 9 in Wells et al. (2013) were used. Wells et al. (2013) tagged two male pilot whales after a mass stranding event in the Florida Keys. However, one of the individual tags stopped transmitting after 16 days, but the other tag transmitted for a total of 67 days; thus, the representative depth distribution contains only the Wells et al. (2013) data from the individual with the longer transmission time. Due to the lack of data on time spent at depth, the proportion of dives made to specific depth ranges from Wells et al. (2013) will be used as a proxy for percentage of time spent at depth. The depth distribution for short-finned pilot whales is given in Table 15.

Table 15. Percentage of Time at Depth for the Short-Finned Pilot Whale^{1,2}

Depth Bin (m)	% of Time at Depth
0–2	32.25
2–50	46.75
50–100	3.00
100–200	5.00
200–300	4.25
300–400	2.75
400–500	1.75
500–600	1.75
600–700	1.75
700–800	0.5
800–900	0.125
900–1,000	0.125

¹Based on data from Wells et al. (2013)

²This depth distribution is also representative of the following species: the long-finned pilot whale, pilot whale guild, Fraser’s dolphin, pygmy sperm whale, dwarf sperm whale, and *Kogia* spp. guild

Aguilar Soto et al. (2008) reported a maximum dive depth of 1019 m for 23 whales near the Canary Islands, which is similar to the deepest depth bin in Table 15. While the maximum depth in Jensen et al. (2011) (roughly 700 m) is hundreds of meters shallower than in the distribution above, the total time spent deeper than this depth constitutes a very small percentage of the whale’s total time.

2.2.1.3.6 *Globicephala melas*, Long-finned Pilot Whale

Long-finned pilot whales are commonly found in cold and temperate continental shelf and slope waters (Buckland et al. 1993; Payne and Heinemann 1993). Long-finned pilot whales prey on epipelagic, mesopelagic, and demersal squid and fish (Desportes and Mouritsen 1993; Gannon et al. 1997).

In order to build a representative depth distribution for long-finned pilot whales, data from Table 2 from Heide-Jorgensen et al. (2002) were used. The maximum reported dive depth was 828 m. Data from the individual whales of different sexes and sizes, as well as in a variety of locations, were rounded to sum to 100 percent of each individual whale’s time, then averaged together to create the depth distribution given in Table 16.

Table 16. Percentage of Time at Depth for the Long-Finned Pilot Whale¹

Depth Bin (m)	% of Time at Depth
0–17	74.4
18–35	5.2
36–53	2.2
54–101	3.8
102–149	2.8
150–197	1.8
198–299	3.4
300–401	2.6
402–599	2.9
600–797	0.9

¹Based on data from Heide-Jorgensen et al. (2002)

Maximum reported dive depths for long-finned pilot whales range differ by location, including a depth of 513 m off the coast of Norway (Aoki et al. 2013) and a depth of 648 m in the Mediterranean Sea (Baird et al. 2002). Long-finned pilot whales show some diurnal variation in diving behavior, with the deepest dives occurring shortly after sunset when the whales may target vertically-migrating prey (Baird et al. 2002). Mate et al. (2005) reported whales spending between 5.6 and 47.4 percent of their time at the surface, while Hooker et al. (2011) found long-finned pilot whales spending an average of 67 percent of time in the top 10 m of the water column. The representative depth distribution in Table 16 is consistent with distribution data reported in Table 1 in Nawojchik et al. (2003), Figures 3a and 8a in Aoki et al. (2013), Figure 3 in Sivle et al. (2012), and plots from post sonar exposure and silent passes of a ship in Miller et al. (2011).

Despite the presence of depth distribution data for long-finned pilot whales, there is a lack of available separate density data each species of pilot whale. As a result, the two species must be represented by one depth distribution. After comparing the depth distribution given in Table 16 with the depth distribution for the short-finned pilot whale given in Table 15, the short-finned pilot whale (Section 2.2.1.3.5) data was selected as the representative depth distribution for use. The short-finned pilot whale data represents a majority of time spent in the top 50 m of the water column rather than at depth. Both pilot whales are members of the same subfamily, Globicephalinae, feed on mesopelagic prey, and are similar in size. The depth distribution for long-finned pilot whales can be found in Table 15.

2.2.1.3.7 Pilot Whale Guild

Within the AFTT Study Area, the distributions of short-finned and long-finned pilot whales overlap. As these two species are difficult to distinguish when viewing at sea, much of the density data from surveys are reported as generic “pilot whales.” Because the short-finned pilot whale depth distribution found in Table 15 was selected as the representative depth distribution for both species of pilot whale, these data also represent the depth distribution for the pilot whale guild. Within this guild, the best available estimate of the population of short-finned pilot whales in the western North Atlantic is 37,112 short-finned pilot whales. As compared to the best available estimate of long-finned pilot whales (at 5,636 long-finned pilot whales), the short-finned pilot whale has a significantly larger population size in the AFTT Study Area (Barlow 2010; Barlow and Forney 2007; Forney 2007; Waring et al. 2016). Short-finned pilot whales also inhabit a larger geographic range, spanning from the western North Atlantic to the Gulf of Mexico, while long-finned pilot whales are only found in the western North Atlantic. The representative depth distribution for pilot whales can be given in Table 15.

2.2.1.3.8 *Lagenodelphis hosei*, Fraser’s Dolphin

Fraser’s dolphins are commonly found in the tropics worldwide in waters deeper than 1,000 m. While they do have occasional strandings in temperate waters, those are thought to be extralimital occurrences (Louella and Dolar 2002; Reeves et al. 2002).

Little data have been collected on the dive behavior of Fraser’s dolphins. Robison and Craddock (1983) reported that the mesopelagic fish, shrimp, and squid species that were found in the stomachs of three dolphins typically inhabit depths between 250 and 500 m. Fraser’s dolphins in the Sulu Sea, off the coast of the Philippines, were found to feed on vertically migrating species in the upper 200 m of the water column, as well as on non-migrating species found at depths below 600 m (Dolar et al. 2003). Fraser’s dolphins have also been seen herding fish near the water’s surface (Watkins et al. 1994).

Due to the lack of available data on the diving behavior of the Fraser’s dolphin, it will be represented by a surrogate species: the short-finned pilot whale (Section 2.2.1.3.5). Dolar et al. (1999) found that Fraser’s dolphins have myoglobin concentrations consistent with those of other deep-diving marine mammals, and relative muscle masses much greater than those of other dolphins their size, both of which are indicative of

enhanced diving ability. Therefore, despite their smaller size, the Fraser's dolphin will be modeled as a short-finned pilot whale, another species in the family Delphinidae which feeds on mesopelagic and bathypelagic prey at similar deep depths (Desportes and Mouritsen 1993; Gannon et al. 1997). The depth distribution for the Fraser's dolphin is given in Table 15.

2.2.1.3.9 *Grampus griseus*, Risso's Dolphin

Risso's dolphins are commonly found in temperate and tropical waters along continental slopes (Azzellino et al. 2008; Baumgartner 1997; Green et al. 1992). Although little is known about their foraging or diving behavior, vertically migrating cephalopods are presumed to be the primary food source for Risso's dolphins (Clarke and Pascoe 1985).

In order to build a representative depth distribution, data from Figure 5 in Wells et al. (2009) were used. Wells et al. (2009) reported on the movement and diving behavior of a rehabilitated adult male Risso's dolphin that stranded on the Gulf coast of Florida. Based on Figure 5, the depth distribution for Risso's dolphins was estimated for four 6-hour blocks of time. The tagged animal in this study travelled through waters with a mean depth of 548 m (range 3–2300 m), and was therefore likely not diving close to the seafloor. The deepest dive recorded on the tag was in the 400–500 m depth range, and less than 0.1 percent of dives were deeper than 200 m (Wells et al. 2009). The average time spent in these depth bins was calculated for the representative depth distribution. The depth distribution for the Risso's dolphin is given in Table 17.

Table 17. Percentage of Time at Depth for the Risso's Dolphin^{1,2}

Depth Bin (m)	% of Time at Depth
0–1	24.8
1–2	13.5
2–10	16.5
10–50	43.5
50–100	1.2
100–150	0.1
150–600	0.4

¹Based on data from Wells et al. (2009)

²This depth distribution is also representative of the following species: the pygmy killer whale, melon-headed whale, and false killer whale

2.2.1.3.10 *Feresa attenuata*, Pygmy Killer Whale

Pygmy killer whales inhabit tropical and subtropical waters of the continental slope and waters farther offshore (Donahue and Perryman 2002). Analyses of their stomach contents indicate that their primary prey include cephalopods and fish, although other marine mammals also constitute some portion of their diet (Mignucci-Giannoni et al. 1999; Perryman and Foster 1980). Some shallow water prey (from less than 200 m deep) have been reported in the stomachs of stranded pygmy killer whales, though these prey species may have been consumed as animals moved closer to shore prior to stranding (Sekiguchi et al. 1992; Zerbini and de Oliveira Santos 1997). Characteristics of their echolocation clicks indicate that pygmy killer whales could detect fish and cephalopod prey at distances of 50–200 m (Madsen et al. 2004b).

Due to a lack of available data on the dive behavior of pygmy killer whales, they will be represented by a surrogate species: the Risso's dolphin (Section 2.2.1.3.9). The closest relatives to the pygmy killer whale for which diving behavior have been studied are three members of the subfamily Globicephalinae: the short-finned pilot whale, the long-finned pilot whale, and the Risso's dolphin (LeDuc et al. 1999). The pygmy killer whale (at 2.3 m long) is closer in size to the Risso's dolphin (4 m) than to the pilot whales (6 m). Both pygmy

killer whales and Risso's dolphins are found in deep water and feed on squid and other cephalopods. The depth distribution for the pygmy killer whale can be found in Table 17.

2.2.1.3.11 *Peponocephala electra*, Melon-headed Whale

Melon-headed whales occur in oceanic tropical and subtropical waters (Perryman et al. 1994). Melon-headed whales feed on a variety of mesopelagic fish (e.g., myctophids) and cephalopod species (Brownell Jr. et al. 2009; Gross et al. 2009; Jefferson and Barros 1997).

No data have been collected on the dive behavior of the melon-headed whale, although some inferences can be made from the primary prey species. Myctophids are vertical migrators that descend to depths of 700–3,000 m during the day and rise to 200 m or less at night (Clarke 1973), when melon-headed whales are thought to feed (Brownell Jr. et al. 2009). Cephalopods are also vertical migrators, descending to depths of 400–700 m during the day and rising to 100–150 m at night (Young 1978).

Due to the lack of available data on the diving behavior of the melon-headed whale, it will be represented by a surrogate species: the Risso's dolphin (Section 2.2.1.3.9). Both species are members of the same subfamily, Globicephalinae, feed on mesopelagic prey, and are similar in size. The closest relative to the melon-headed whale is actually the pilot whale; however, pilot whales are more than twice the size of melon-headed whales (LeDuc et al. 1999). The small size of melon-headed whales may indicate that they do not dive as deeply as their larger relatives; thus, the Risso's dolphin is a more suitable surrogate species for dive behavior. The depth distribution for the melon-headed whale is given in Table 17.

2.2.1.3.12 *Pseudorca crassidens*, False Killer Whale

False killer whales inhabit tropical and temperate waters along and offshore of the continental slope (Odell and McClune 1999). Stomach content analyses have revealed that false killer whales feed on oceanic cephalopods (Alonso et al. 1999; Andrade et al. 2001), while observations indicate that they consume a variety of prey (including fish and other marine mammals) both at depth and at the surface (Acevedo-Gutiérrez et al. 1997; Perryman and Foster 1980; Stacey et al. 1994).

Little data have been collected on the dive behavior of the false killer whale. Cummings and Fish (1971) estimated that false killer whales would be capable of diving to depths up to 500 m. Based on measurements of their echolocation clicks, whales may detect large fish at up to 200 m distance and cephalopods at about half that distance (Madsen et al. 2004a), which may suggest false killer whales are capable of diving to at least 200 m. Unpublished time-depth recorder data of a single whale showed that all dives to deeper than 100 m occurred during the day, with a maximum depth exceeding 234 m (Baird 2009). Dives during the nighttime remained within the top 100 m of the water column. Ligon and Baird (2001) reported that three instrumented whales showed a maximum diving depth of 53 m, with an average dive depth range of 8–12 m; however, the time of day that the dives occurred was not reported.

Due to the lack of available data on the diving behavior of the false killer whale, it will be represented by a surrogate species: the Risso's dolphin (Section 2.2.1.3.9). The closest relatives to the false killer whale for which diving behavior have been studied are three members of the subfamily Globicephalinae: the short-finned pilot whale, the long-finned pilot whale, and the Risso's dolphin (LeDuc et al. 1999). False killer whales are in between these species in size, but the limited data suggest that false killer whales do not dive as deep as pilot whales. Risso's dolphins and false killer whales also both feed on pelagic cephalopods (Clarke and Pascoe 1985). The depth distribution for the false killer whale is given in [Table 17](#).

2.2.1.3.13 *Stenella attenuata*, Pantropical Spotted Dolphin

Pantropical spotted dolphins are found in warm temperate and tropical waters over the continental slope and offshore in deeper waters (Perrin and Hohn 1994). Pantropical spotted dolphins feed on both epipelagic and mesopelagic fish and squid (Wang et al. 2003). In general, pantropical spotted dolphins dive deeper at night, foraging on prey associated with vertical migrations of the deep scattering layer (Robertson and Chivers 1997; Scott and Chivers 2009).

In order to build a representative depth distribution for pantropical spotted dolphins, data from Figure 4 and Table 2 in Baird et al. (2001) and Figure 9 and Table 2 in Scott and Chivers (2009) were used. While Baird et al. (2001) looked at pantropical spotted dolphin diving behavior around the Hawaiian Islands, Scott and Chivers (2009) recorded data on these dolphins in pelagic waters. Baird et al. (2001) reported pantropical spotted dolphins spend on average 88.5 percent of their time within 10 m of the surface during the day. Baird et al. (2001) reported the daytime average percentage of time in two meter intervals for the top 10 meters. For the Baird et al. (2001) nighttime and Scott and Chivers (2009) data, the percentage of time determined for the top 10 meters was uniformly distributed across these two meter intervals. Daytime and nighttime averages were calculated for the Baird et al. (2001) data, and these were then averaged with the Scott and Chivers (2009) data. The resulting mean daytime and nighttime depth distribution data are presented in Table 18. Baird reported maximum daytime and nighttime dive depths at 122 m and 213 m, respectively (Baird et al. 2001); however, Scott and Chivers (2009) calculated that dives to more than 120 m accounted for less than 0.1 percent of all dives. They also noted that daytime dives were primarily shallow and above the thermocline (Scott and Chivers 2009). The depth distribution for pantropical spotted dolphins is given in Table 18.

Table 18. Percentage of Time at Depth for the Pantropical Spotted Dolphin^{1,2}

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–2	20.4	70–80	0.6
2–4	10.7	80–90	0.6
4–6	8.6	90–100	0.4
6–8	9.0	100–110	0.4
8–10	9.5	110–120	0.3
10–20	21.3	120–130	0.1
20–30	8.8	130–140	0.1
30–40	3.8	140–150	0.1
40–50	2.5	150–160	0.1
50–60	1.9	160–170	0.1
60–70	1.1		

¹Based on data from Baird et al. (2001) and Scott and Chivers (2009)

²This depth distribution is also representative of the following species: the long-beaked common dolphin, short-beaked common dolphin, Atlantic white-sided dolphin, white-beaked dolphin, Pacific white-sided dolphin, Northern Right whale dolphin, clymene dolphin, striped dolphin, and spinner dolphin

2.2.1.3.14 *Delphinus capensis*, Long-beaked Common Dolphin

It was not until the mid-1990s that the long-beaked common dolphin was separated from the short-beaked common dolphin as a distinct species (Heyning and Perrin 1994). Long-beaked common dolphins are thought to be coastal foragers, feeding mostly on pelagic fish, particularly those in the families Scombridae, Scianidae, and Serranidae (Niño-Torres et al. 2006) as well as krill and squid (Reeves et al. 2002).

No data have been collected on the dive behavior of the long-beaked common dolphin. Due to the lack of available data on the diving behavior of the long-beaked common dolphin, it will be represented by a

surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). The closest relative to the long-beaked common dolphin is the short-beaked common dolphin, which is also represented by the pantropical spotted dolphin. While the long-beaked common dolphin is considered a more coastal species than the short-beaked common dolphin, in many areas their distributions overlap. The depth distribution of the long-beaked common dolphin can be found in Table 18.

2.2.1.3.15 *Delphinus delphis*, Short-beaked Common Dolphin

While species of common dolphins are sympatric in some nearshore continental shelf waters, short-beaked common dolphins are typically found in deeper waters along the continental slope (Cañadas and Hammond 2008; Heyning and Perrin 1994; Jefferson et al. 2009; Rosel et al. 1994; Selzer and Payne 1988). They feed on epipelagic and mesopelagic fish and squid (Selzer and Payne 1988), and also forage at night on vertically migrating prey associated with the deep scattering layer (Evans 1994; Neumann and Orams 2003; Ohizumi et al. 1998; Pusineri et al. 2007).

Little data have been collected on the dive behavior of the short-beaked common dolphin. Evans (1975; 1994) described the late afternoon and evening diving behavior of an adult female short-beaked common dolphin in the Pacific Ocean. Before 1730 the dolphin mostly remained in the top 10 m, at which time it switched to a pattern of regular dives to 50 m, with a maximum dive depth of just over 200 m (Evans 1974, 1994).

Due to the lack of available data on the diving behavior of the short-beaked common dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). Pantropical spotted dolphins also make shallower dives during the day than at night, when they forage on vertically-migrating prey associated with the deep scattering layer (Scott and Chivers 2009). During the day, pantropical spotted dolphins spend 94 percent of their time in the top 20 m of the water column, while at night 95 percent of their time is spent in the top 50 m (Baird et al. 2001). Evans (1994) reported the maximum dive depth for three short-beaked common dolphins was 257 m. Similarly, maximum dive depths for pantropical spotted dolphins are 122 m for daytime and 213 m for nighttime (Baird et al. 2001). Pantropical spotted dolphins and common dolphins are members of the same subfamily, Delphinidae (LeDuc et al. 1999) and their behavior shows clear similarities in diving pattern, foraging behavior, and water column usage. The depth distribution of the short-beaked common dolphin is given in Table 18.

2.2.1.3.16 *Lagenorhynchus acutus*, Atlantic White-sided Dolphin

Atlantic white-sided dolphins are found in cold temperate and sub-polar waters, ranging from the continental shelf to further offshore of the continental slope (Palka et al. 1997). They feed on both epipelagic and mesopelagic fish and squid (Couperus 1997; Craddock et al. 2009; Doksaeter et al. 2008; Weinrich et al. 2001).

Little data have been collected on the dive behavior of Atlantic white-sided dolphins. One stranded and rehabilitated dolphin with a satellite tag provided basic dive behavior information (Mate et al. 1994). Over a six day period, no dives were longer than four minutes, and no surface durations lasted longer than one minute; the Atlantic white-sided dolphin spent 89 percent of its time beneath the surface at unknown depths. White-sided dolphins caught in bottom trawl and sink gillnet fisheries as bycatch were recovered in nets fished at a mean depth of 189.8 m (range 55–503 m), although the exact depths at which the animals were entangled are unknown (Craddock et al. 2009).

Due to the lack of available data on the diving behavior of the Atlantic white-sided dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). Both species are members of the same family, Delphinidae, and forage on similar vertically migrating prey species that come

to the surface layer (above 200 m) at night (Scott and Chivers 2009; Wang et al. 2003). The depth distribution for the Atlantic white-sided dolphin is given in Table 18.

2.2.1.3.17 *Lagenorhynchus albirostris*, White-beaked Dolphin

White-beaked dolphins occur only in the temperate and subarctic North Atlantic Ocean, generally in coastal and continental shelf waters (Kinze 2002). Fish species constitute the primary prey of white-beaked dolphins (e.g., Atlantic cod, haddock, herring, and hake); however, cephalopods and benthic crustaceans are also part of their diet (Ostrom et al. 1993; Reeves et al. 1999). They have also been observed cooperatively herding schooling fish species at the surface (Reeves et al. 1999).

Little data have been collected on the dive behavior of the white-beaked dolphin. White-beaked dolphins generally prefer shallow waters less than 200 m deep (Jefferson et al. 2008). In the 1970s and 1980s, white-beaked dolphins off the northeastern U.S. Atlantic coast may have shifted habitats with Atlantic white-sided dolphins. During this time, white-beaked dolphins, which were normally found in inshore waters, moved offshore due to an increase in sand lance on the continental shelf and a decline in herring.

Due to the lack of available data on the diving behavior of the white-beaked dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). The closest relative to the white-beaked dolphin is the Atlantic white-sided dolphin, which is also represented by the pantropical spotted dolphin. The depth distribution for the white-beaked dolphin can be found in Table 18.

2.2.1.3.18 *Lagenorhynchus obliquidens*, Pacific White-sided Dolphin

Pacific white-sided dolphins inhabit cold temperate waters of the North Pacific, in both offshore and coastal waters (Brownell et al. 1999; Waerebeek and Würsig 2002). Their primary prey species include mesopelagic fish and cephalopods, as well as epipelagic fish in shallower waters (Brownell et al. 1999; Kajimura and Loughlin 1988; Miyazaki et al. 1991; Morton 2000; Walker and Jones 1994).

Little data have been collected on the dive behavior of the Pacific white-sided dolphin. Hall (1970) trained a captive Pacific white-sided dolphin to dive to a depth of 214 m. However, Black (1994) reported that in coastal waters, 70 percent of dives were shorter than 20 seconds in duration, and dives longer than 90 seconds were rare, indicating that most dives are shallow. Heise (1997) similarly reported that 70 percent of foraging dives were less than 15 seconds in duration. Therefore, Pacific white-sided dolphins are not considered deep divers. This species is thought to feed mostly at night or in the morning (Stroud et al. 1981) when their mesopelagic prey rise to surface waters.

Due to the lack of available data on the diving behavior of the Pacific white-sided dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). Pantropical spotted dolphins spend the majority of their time in the top 50 m, and their maximum diving depths are within the range of the dive depth of the trained Pacific white-sided dolphin (Baird et al. 2001; Hall 1970; Scott and Chivers 2009). Pantropical spotted dolphins and Pacific white-sided dolphins also feed on similarly migrating prey species. The depth distribution for the Pacific white-sided dolphin can be found in Table 18.

2.2.1.3.19 *Lissodelphis borealis*, Northern Right Whale Dolphin

The northern right whale dolphin is abundant in deep, temperate waters across the North Pacific Ocean (Forney and Barlow 1998; Jefferson and Newcomer 1993b; Leatherwood and Walker 1979; Rankin et al. 2007). They are known to commonly associate with Pacific white-sided dolphins and Risso's dolphins (Forney and Barlow 1998; Jefferson and Newcomer 1993b), with which they show dietary overlap (Walker and Jones 1994). Northern right whale dolphins near the southern California coast feed principally on cephalopods and

a diverse variety of myctophid fish (Jefferson and Newcomer 1993a; Jefferson et al. 1994; Leatherwood and Walker 1979).

Little data have been collected on the dive behavior of the Northern right whale dolphin. Some evidence based on stomach contents suggests that northern right whale dolphins may dive as deep as 200 m (Fitch and Brownell 1968; Jefferson et al. 1994). Individual northern right whale dolphins have been observed to dive for brief periods (10 to 75 seconds), but can also remain submerged for up to 6.5 min (Cruickshank and Brown 1981; Leatherwood and Walker 1979). Northern right whale dolphins have comparatively low muscle myoglobin content among odontocetes, suggesting they are not deep divers (Noren and Williams 2000).

Due to the lack of available data on the diving behavior of the northern right whale dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). Of the two dolphins with which the northern right whale dolphin associates the most, the Risso's dolphin is considered a much deeper diver than the Pacific white-sided dolphin. Because northern right whale dolphins have low muscle myoglobin content and are thought to feed on prey only as deep as 200 m, they are thought to be shallower divers. Therefore, due to dietary similarity and frequent association with the Pacific white-sided dolphin, the northern right whale dolphin will be represented by the same surrogate species, the pantropical spotted dolphin. The depth distribution for the Northern right whale dolphin can be found in Table 18.

2.2.1.3.20 *Stenella coeruleoalba*, Striped Dolphin

Striped dolphins prefer tropical and warm temperate waters and have an oceanic distribution, with most observations occurring beyond the continental shelf (Archer II 2002; Cañadas et al. 2002; Davis and Fargion 1996; Davis et al. 1998; Perrin et al. 1994b). Striped dolphins primarily feed on small, pelagic, vertically migrating prey (Blanco et al. 1995). Stomach contents analyses suggest that foraging occurs mostly in the dusk and early evening hours (Ringelstein et al. 2006). Their distribution in the North Atlantic Ocean is associated with a mesopelagic prey community comprised of fish and cephalopod species (Doksaeter et al. 2008).

Little data has been collected on the dive behavior of the striped dolphin. A single striped dolphin carrying a time-depth recorder dove to a mean depth of 22.6 m (standard deviation [SD]=17.5) during the day and 126.7 m (SD=120.9) at night, with a maximum dive depth of 705 m (Minamikawa et al. 2003).

Due to the lack of available data on the diving behavior of the striped dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). The observed pattern of shallow daytime shallow diving and deeper nighttime diving reported in Minamikawa et al. (2003) is consistent with similar diving behavior seen in short-beaked common dolphins (Section 2.2.1.3.15) and their surrogate species, pantropical spotted dolphins (Section 2.2.1.3.13) which is also in the genus *Stenella*. Additionally, all three species occur in similar water depths (Davis et al. 1998). However, it is acknowledged that the striped dolphin may dive to deeper depths on average, due to the deep maximum dive depth recorded by Minamikawa et al. (2003). The depth distribution for striped dolphins is given in Table 18.

2.2.1.3.21 *Stenella longirostris*, Spinner Dolphin

Spinner dolphins typically reside in tropical pelagic waters, although they have a coastal distribution around the Hawaiian and French Polynesia island chains (Benoit-Bird and Au 2003). The prey of spinner dolphins consists of vertically migrating mesopelagic fish, cephalopods, and crustaceans, as well as pelagic organisms concentrated in near-surface waters with a shallow thermocline (Lammers 2004; Reilly 1990).

No data have been collected on the dive behavior of the spinner dolphin. Many of the vertically migrating prey of the spinner dolphin spend daytime hours at depths from 700–3,000 m, but ascend to depths between the surface and 200 m at night. Spinner dolphins in Hawaiian waters mostly forage in deep water at

dusk and early evening, but dive to shallow depths due to the vertical migration of their prey at night (Benoit-Bird and Au 2003; Lammers 2004).

Due to the lack of available data on the diving behavior of the spinner dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). Pantropical spotted dolphins also forage mostly at night on vertically migrating fish and cephalopod prey and their foraging dives are primarily limited to the upper 200 m of the water column (Baird et al. 2001). Gross et al. (2009) found no niche differentiation between the two species. The depth distribution for striped dolphins can be found in Table 18.

2.2.1.3.22 *Stenella clymene*, Clymene Dolphin

Clymene dolphins are found in deep tropical and warm temperate waters beyond the continental shelf (Jefferson 2002a). The limited information available regarding their prey suggests they primarily feed on squid and vertically migrating myctophid fish, which are associated with the deep scattering layer (Perrin et al. 1981).

No data has been collected on the dive behavior of the Clymene dolphin. Some authors have speculated that their size limits them to the upper 250 m of the water column (Davis et al. 1998), although preliminary evidence suggests similarly-sized striped dolphins are capable of diving much deeper (Minamikawa et al. 2003).

Due to the lack of available data on the diving behavior of the Clymene dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.3.13). Also in the genus *Stenella* and a shallower diver amongst the dolphin species, striped and spinner dolphins are the Clymene dolphin's closest relatives. Thus, the Clymene dolphin will be represented by the same surrogate species, the pantropical spotted dolphin. The depth distribution for the Clymene dolphin is given in Table 18.

2.2.1.4 Family Eschrichtiidae

2.2.1.4.1 *Eschrichtius robustus*, Gray Whale

Gray whales are distributed coastally throughout the Pacific Ocean, migrating annually between Arctic and subtropical waters (Jones and Swartz 2002; Swartz 1986). Gray whales forage within the water column with modified skimming techniques to capture neritic fish, and will scrape along the benthos to acquire benthic fish, squid, annelids, crustaceans, and mollusks (Darling et al. 1998; Dunham and Duffus 2002; Jones and Swartz 2002; Nerini 1984). Gray whales have been reported foraging in water up to 120 m deep (Cacchione et al. 1987; Dunham and Duffus 2002; Würsig et al. 1986), although in many areas whales forage in waters less than 20 m deep (Guerrero 1989; Ljungblad et al. 1987; Malcolm and Duffus 2000; Malcolm et al. 1995; Stewart et al. 2001; Woodward and Winn 2006).

In order to build a representative depth distribution for gray whales, data from Figure 3 in Malcolm et al. (1995) were used. Malcolm et al. (1995) reported the percentage of time at depth for a single foraging whale carrying a tag for over 8 hours in waters off British Columbia in an area with a mean bottom depth of 18 m. The majority of dives (76 percent) were ventilation dives (to a mean depth of 2.3 m), while 13 percent were feeding dives (to a mean depth of 16.7 m). The whale appeared to spend little time at intermediate depths within the water column, spending most of its time either at the surface breathing or at the bottom feeding. Due to the large size of gray whales, the data from Malcolm et al. (1995) was summed into four-meter bins for the representative depth distribution. The depth distribution for gray whales is given in Table 19.

Table 19. Percentage of Time at Depth for the Gray Whale¹

Depth Bin (m)	% of Time at Depth
0–4	39.0
4–8	8.5
8–12	7.0
12–16	16.0
16–20	28.0
20–22	1.5

¹Based on data from Malcom et al. (1995)

While other studies did not include depth distributions for gray whales, they did provide additional information to categorize dive behavior. The representative depth distribution data compare to a later study with a larger sample size of whales, where 79 percent of dives by whales off Vancouver Island were to a mean depth of 2.2 m, and 15 percent of dives were to a mean depth of 12–19 m (Malcolm and Duffus 2000). Woodward and Winn (2006) and Woodward (2006) similarly reported that six whales feeding along the central British Columbia coast had a mean dive depth of 11 m (range 2.4–28.9 m). The percentage of time near the surface (19.5 percent from 0–2 m) is also consistent with other studies in the same region (14.2 percent and 17.5 percent) (Stelle et al. 2008) and in the Bering Sea (22 percent) (Würsig et al. 1986). Furthermore, the dive depth is similar to the reported foraging depths in British Columbia and other regions (Guerrero 1989; Ljungblad et al. 1987; Malcolm and Duffus 2000; Malcolm et al. 1995; Stewart et al. 2001; Woodward and Winn 2006). Stewart et al. (2001) described the diving behavior post-release for a rehabilitated calf in southern California. All dives were less than 20 m deep, and 85 percent of dives were less than 10 m deep. An earlier release of a post-rehabilitated calf in the same area documented a much deeper maximum diving depth (170 m) and an average diving depth of approximately 50 m (Evans 1974), which is deeper than the that in the representative depth distribution.

2.2.1.5 Family Kogiidae

2.2.1.5.1 *Kogia breviceps*, Pygmy Sperm Whale

Pygmy sperm whales have a cosmopolitan distribution in all temperate and tropical waters (Bloodworth and Odell 2008). Mid- and deep-water cephalopods predominantly contribute to the diet of the pygmy sperm whale (Beatson 2007; Bloodworth and Odell 2008; Fernandez et al. 2009; McAlpine et al. 1997; Ross 1979; Santos et al. 2006).

Little data have been collected on the dive behavior of pygmy sperm whales. Sightings of pygmy sperm whales in the North Atlantic are most common in waters ranging from 400 to 1,000 m in depth (Clarke 2003; Scott et al. 2001; Waring et al. 2006). Based on the analysis of the stomach contents of whales stranded in New Zealand, Beatson (2007) concluded that pygmy sperm whales feed at shallower depths within the water column than sperm whales, although some prey species are found at depths greater than 600 m. Ploen (2004) found that prey species from the stomachs of stranded pygmy sperm whales in South Africa are found at depths below 300 m.

Due to the lack of available data on the diving behavior of the pygmy sperm whale, it will be represented by the surrogate species: the short-finned pilot whale (Section 2.2.1.3.5). The short-finned pilot whale is another primarily squid-eating species (Mintzer et al. 2008; Reeves et al. 2002) which forages deep in the water column (Jensen et al. 2011). The broad similarity in prey types and oceanic habitat suggests similarity in diving behavior to the short-finned pilot whale. The depth distribution for the pygmy sperm whale can be found in Table 15.

2.2.1.5.2 *Kogia sima*, Dwarf Sperm Whale

Dwarf sperm whales have a cosmopolitan distribution in all temperate and tropical waters (Willis and Baird 1998). Little data have been collected on the dive behavior of dwarf sperm whales. There is some indication that dwarf sperm whales have a more coastal distribution than pygmy sperm whales, and prey often include more continental shelf and slope species than those of the pygmy sperm whale (Ross 1979; Wang et al. 2002). The preferred prey species of dwarf sperm whales are found deep in the water column, with some species found below 400 m (Wang et al. 2002).

Due to lack of available data on the diving behavior of the dwarf sperm whale, it will be represented by the surrogate species: the short-finned pilot whale (Section 2.2.1.3.5). The broad similarity in depth at which preferred prey can be found and oceanic habitat suggests similarity in diving behavior to the short-finned pilot whale. In addition, the dwarf sperm whale's closest relative, the pygmy sperm whale, is also represented by the short-finned pilot whale. The depth distribution for the dwarf sperm whale can be found in Table 15.

2.2.1.5.3 *Kogia* spp. Guild

Within the AFTT Study Area, the two *Kogia* species are difficult to distinguish when viewed at sea and are found in overlapping areas within the AFTT study area, as well as the Southern California portion of the HSTT Study Area. Therefore, much of the density data from surveys is combined for both species into a generic *Kogia* spp. guild. Because the depth distributions for both *Kogia* species are represented by the short-finned pilot whale, these data also represent the depth distribution for the *Kogia* spp. guild. Therefore, the depth distribution for *Kogia* spp. whales can be found in Table 15.

2.2.1.6 Family Phocoenidae

2.2.1.6.1 *Phocoena phocoena*, Harbor Porpoise

Harbor porpoises inhabit temperate and sub-arctic continental shelf waters in the northern hemisphere. Their diet consists primarily of fish, including both pelagic schooling and benthic species (Bjorge and Tolley 2002; Recchia and Read 1989). Cephalopods, crustaceans, euphausiids, and polychaetes also contribute to their overall diet (Recchia and Read 1989; Smith and Read 1992; Walker et al. 1998).

In order to build a representative depth distribution for harbor porpoises, data from Figure 3 in Otani et al. (1998), Figures 2 and 3 in Westgate et al. (1995), Figure 1 in Otani et al. (2000), Figures 2, 3, and 4 in Cooper et al. (1993), and Figure 7 in Westgate and Read (1998) were used. Cooper et al. (1993) reported that porpoises in the Bay of Fundy were capable of diving to 150 m, but spent most of their time in the top 50 m of the water column. Westgate et al. (1995) reported porpoises diving to a maximum of 226 m, but that average dive depth for individual porpoises ranged from 14–41 m, and the depth range with the greatest proportion of dives was 2–10 m. Otani et al. (1998) found that harbor porpoises off the coast of Japan spent 74–86 percent of their time in the top 20 m of the water column, with an average dive depth of 12–19 m. The time at depth was visually inspected and averaged from all of the above Figures to create the depth distribution for the harbor porpoise. This was done to include a total of 14 different harbor porpoises into the dive distribution. While data from Cooper (1993), Otani (1998), and Westgate et al. (1995) show the number or frequency of dives to specific depth bins, this data will be used as a proxy for percentage of time spent at depth. The depth distribution for harbor porpoises is given in Table 20.

Table 20. Percentage of Time at Depth for the Harbor Porpoise¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–10	39.02	110–120	0.26
10–20	17.79	120–130	0.35
20–30	12.74	130–140	0.34
30–40	10.14	140–150	0.23
40–50	6.90	150–160	0.16
50–60	4.61	160–170	0.06
60–70	2.52	170–180	0.06
70–80	1.50	180–190	0.06
80–90	1.37	190–200	0.06
90–100	0.98	200–210	0.06
100–110	0.31	210–226	0.06

¹Based on data from Cooper (1993), Otani et al. (2000), Otani et al. (1998), Westgate and Read (1998), and Westgate et al. (1995)

While other studies did not include depth distributions for harbor porpoises, they did provide additional information to categorize dive behavior. In a study by Linnenschmidt et al. (2013) three harbor porpoises were tagged; two of the harbor porpoises showed consistent diving activity throughout the day while one harbor porpoise showed a diel diving pattern with few dives during the day. Teilmann et al. (2013) found harbor porpoises spent more time between 0 to 2 m at night than during the day; this may be due to the movement of their prey throughout the water column (e.g., herring [*Clupea harengus*] and sprat [*Sprattus sprattus*]) (Cardinale et al. 2003).

2.2.1.6.2 *Phocoenoides dalli*, Dall's Porpoise

Dall's porpoises can be found in the subarctic and cool temperate North Pacific Ocean, including the Bering Sea, Okhotsk Sea, and Sea of Japan (Jefferson 2002b). Primary prey species include epipelagic and mesopelagic schooling fish and cephalopod species (Jefferson 1988; Ohizumi et al. 2000; Stroud et al. 1981; Walker 1996).

In order to build a representative depth distribution for the Dall's porpoise, data from Figures 3 and 4 in Baird and Hanson (1998) were used. Baird and Hanson (1998) tagged three Dall's porpoises with time-depth recorders in the waters between Washington State and British Columbia. Each animal had a median dive depth of less than 40 m, and maximum dive depths ranged from 197–278 m. Data from the tagged Dall's porpoises were averaged together to create the representative depth distribution below. The depth distribution for Dall porpoises is given in Table 21.

Table 21. Percentage of Time at Depth for the Dall's Porpoise¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0-1	5.33	50-60	4.75
1-2	15.67	60-70	4.67
2-3	8.83	70-80	3.83
3-4	3.17	80-90	2.25
4-5	3.75	90-100	2.00
5-6	1.67	100-110	1.08
6-7	1.00	110-120	0.67
7-8	1.33	120-130	0.42
8-9	1.25	130-140	0.50
9-10	1.83	140-150	0.42
10-11	1.33	150-160	0.33
11-20	11.67	160-170	0.25
20-30	8.33	170-180	0.33
30-40	6.83	180-190	0.25
40-50	6.17	190-200	0.08

¹Based on data from Baird and Hanson (1998)

The representative depth distribution is consistent with stomach contents analyses which suggest that Dall's porpoises feed high in the water column on vertically migrating mesopelagic species, but occasionally forage on deeper benthic prey (Jefferson 1988; Ohizumi et al. 1998).

2.2.1.7 Family Physeteridae

2.2.1.7.1 *Physeter macrocephalus*, Sperm Whale

The sperm whale has a cosmopolitan distribution, preferring deeper waters seaward of the continental shelf edge (Whitehead 2002). Females and immature males tend to inhabit tropical and temperate waters below 40° latitude, while maturing and adult males move to higher latitudes, occurring in polar waters as adults (Whitehead 2002). Sperm whales feed on cephalopod species, primarily squid, as well as mesopelagic and demersal fish and occasionally crustaceans (Fiscus et al. 1989; Flinn et al. 2002; Kawakami 1980; Martin and Clarke 1986).

To account for published differences in the foraging dive behavior of whales in different regions, separate depth distributions were generated for the Atlantic Ocean, Gulf of Mexico, and the Pacific Ocean. In general, time spent at depth for the regions is consistent with foraging dives to 800–1,200 m in the Western North Atlantic Ocean (Sivle et al. 2012; Teloni et al. 2008; Watwood et al. 2006), and 400–1,300 m in the Western North Pacific Ocean (Amano and Yoshioka 2003; Aoki et al. 2012; Aoki et al. 2007). Overall, sperm whales typically spend 70–80 percent of their time between 20 and 400 m (Sivle et al. 2012; Teloni et al. 2008). At mid- and low latitudes, females and immature animals undertake stereotypic dives lasting about 45 min and to depths between 400 and 1,200 m (Teloni et al. 2008; Watwood et al. 2006). Off Japan, females and immature sperm whales performed similarly stereotyped dive patterns to 1,400 m, lasting 30–50 min (Aoki et al. 2012). Radically different dive behavior has been observed at high latitudes, where mature males undertake dives lasting up to 60 min and to depths of nearly 1,900 m (Sivle et al. 2012; Teloni et al. 2008).

In order to build a representative depth distribution for sperm whales in the Atlantic Ocean, data from Figure 4 in Sivle et al. (2012), Figure 2 in Teloni et al. (2008), and Figure 2 and raw data from Watwood et al. (2006) were used. Sivle et al. (2012) found that the four tagged sperm whales in their study made regular deep dives to depths of 200–1,500 m, averaging a duration of 25 min, followed by a period of 5–15 min of shallow diving

close to the surface. Sivle et al. (2012) found that sperm whales spend 80 percent of their time deeper than 20 m. Teloni et al. (2008) recorded the dives of four male sperm whales. Dive duration was between 6 and 60 min and dives ranged in depth between 14 and 1,860 m, with a median depth of 175 m. Most surfacing events (92 percent) lasted less than 15 min (Teloni et al. 2008). Watwood et al. (2006) reported that for whales in the Atlantic Ocean, Gulf of Mexico, and Ligurian Sea, typical foraging dives lasted 45 min, regardless of location. Data was extracted from the depth profiles given by Sivle et al. (2012) and Teloni et al. (2008). The resulting data were then averaged across the four whales recorded in the Teloni et al. (2008) study. The data from Sivle et al. (2012), Teloni et al. (2008), and Watwood et al. (2006) were then averaged together for the resulting representative depth distribution for sperm whales in the Atlantic Ocean, which is given in Table 22.

Table 22. Percentage of Time at Depth for the Sperm Whale in the Atlantic Ocean¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–50	29.25	701–800	3.30
51–100	19.14	801–900	2.93
101–150	13.48	901–1000	1.83
151–200	3.33	1001–1100	1.44
201–250	2.72	1101–1200	1.18
251–300	2.29	1201–1300	1.38
301–350	2.35	1301–1400	0.57
351–400	1.39	1401–1500	0.50
401–450	1.17	1501–1600	0.49
451–500	1.03	1601–1700	0.59
501–600	2.94	1701–1800	1.03
601–700	5.52	1801–1900	0.15

¹Based on data from Sivle et al. (2012), Teloni et al. (2008), and Watwood et al. (2006)

For the representative depth distribution of sperm whales in the Gulf of Mexico, data from Watwood et al. (2006) were used. In the Gulf of Mexico, 29 tagged whales spent 28 percent of their time within 10 m of the surface (Watwood et al. 2006). The data represented in Figure 2 was obtained as raw data from the author, then binned, resulting in the depth distribution for sperm whales in the Gulf of Mexico given in Table 23.

Table 23. Percentage of Time at Depth for the Sperm Whale in the Gulf of Mexico¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–50	35.65	501–550	7.72
51–100	1.89	551–600	7.33
101–150	1.85	601–650	4.74
151–200	1.77	651–700	6.24
201–250	1.78	701–750	3.24
251–300	1.83	751–800	1.08
301–350	2.37	801–850	0.27
351–400	4.41	851–900	0.55
401–450	5.98	901–950	1.38
451–500	8.93	951–1000	0.43

¹Based on data from Watwood et al. (2006)

To build a representative depth distribution for sperm whales in the Pacific Ocean, data from Aoki et al. (2007) were used. Aoki et al. (2007) tagged four whales off the coast of Japan. The mean dive depth for nighttime was 515 m averaged over the two tag locations; the mean dive depth for the daytime was 749.5 m.

While this may suggest a diel diving pattern that follows the availability of prey, the pattern seems to depend largely on location (Aoki et al. 2012). The depth distribution for sperm whales in the Pacific Ocean is given in Table 24.

Table 24. Percentage of Time at Depth for the Sperm Whale in the Pacific Ocean¹

Depth Bin (m)	% of Time at Depth
1–50	30.24
51–100	4.77
101–150	3.29
151–200	3.30
201–250	3.13
251–300	2.55
301–350	2.60
351–400	3.60
401–450	5.31
451–500	6.48
501–600	9.76
601–700	6.46
701–800	8.58
801–900	8.38
901–1000	1.26
1001–1100	0.34

¹Based on data from Aoki et al. (2012)

2.2.1.8 Family Ziphiidae

2.2.1.8.1 *Mesoplodon densirostris*, Blainville’s Beaked Whale

Blainville’s beaked whales inhabit deep temperate and tropical waters of the world’s oceans (Pitman 2002b). Little is known about prey species, but the diet of Blainville’s beaked whales includes mesopelagic cephalopods, fish, and crustaceans (Herman et al. 1994; Hickmott 2005; MacLeod et al. 2003a; Mead 1989).

In order to build a representative depth distribution for Blainville’s beaked whales, data were acquired from Figures 3a and 3b from Arranz et al. (2011), Figure 6 from Baird et al. (2005b), Figures 3a and 3b from Baird et al. (2006), Figure 1 from Barlow et al. (2013), Digital Acoustic Recording Tag (DTAG) data from Johnson and Aguilar de Soto (2008b) and Johnson and Aguilar de Soto (2008a), DTAG data from Tyack (2010, and Figure 1b from Tyack et al. (2006. Arranz et al (2011) tagged 9 whales to collect acoustic and movement data, looking to study buzz and click behaviors during dives; Figures 3a and 3b show the dive profile of a male Blainville’s beaked whale over a period of 17 hours. Baird et al. (2005b) tagged four Blainville’s beaked whales, and presented cumulative percentage of time spent at depth for two individuals: an adult female with young calf, the daytime data for a large sub-adult or adult female, and the nighttime data for the same sub-adult or adult female. Different data from that same female whale were used to create another set of dive profiles after a 22.6 hour deployment, as published in Baird et al. (2006). Barlow et al. (2013) used DTAGs to collect acoustic information for beaked whales in an attempt to estimate density and abundance; Figure 1 is a typical dive profile for a tagged whale captured over a 15 hour period. Raw DTAG data for two animals in the Canary Islands, provided by Johnson and Aguilar de Soto (Johnson and Aguilar de Soto 2008a, 2008b), were binned, as well as raw DTAG data from one animal in the Bahamas by Tyack (2010). Tyack et al. (2006) used DTAGs to create a representative dive profile for Blainville’s beaked whale in an attempt to study how depth impacts foraging tactics. Data from each source were arranged into 100 m bins, and those bins

were averaged together to create a representative depth distribution. The depth distribution for Blainville's beaked whales is given in Table 25.

Table 25. Percentage of Time at Depth for the Blainville's Beaked Whale^{1,2}

Depth Bin (m)	% of Time at Depth
1–100	54.3
100–200	10.2
200–300	3.8
300–400	3.2
400–500	3.7
500–600	3.4
600–700	3.8
700–800	4.2
800–900	4.7
900–1,000	3
1,000–1,100	2.2
1,100–1,200	1.8
1,200–1,300	1.1
1,300–1,400	0.8
1,400–1,500	0.1

¹Based on data from Arranz et al. (2011), Baird et al.(2005b), Baird et al. (2006), Barlow et al. (2013), Johnson and Aguilar de Soto (Johnson and Aguilar de Soto 2008a, 2008b), Tyack et al. (2006), and Tyack (2010)

² This depth distribution is also representative of the Longman's beaked whale, the HSTT beaked whale guild, and the AFTT beaked whale guild

2.2.1.8.2 *Indopacetus pacificus*, Longman's Beaked Whale

Longman's beaked whales inhabit generally warm, deep, pelagic waters of tropical and subtropical regions. Little is known about the Longman's beaked whale feeding behaviors. Similar to other beaked whales who dive deep to forage for food, their diet most likely consists primarily of cephalopods (Yamada 2002).

Little data have been collected on the dive behavior of the Longman's beaked whale. The existence of Longman's beaked whales had previously only been known from the skeletal remains of stranded animals (Moore 1972; Pitman 2002a); however, some live sightings have been recognized as this species (Dalebout et al. 2003; Pitman et al. 1999). Dive times have been found to last from 11 to 33 min, although one dive may have been over 45 min (Anderson et al. 2006; Gallo-Reynoso and Figueroa-Carranza 1995).

Due to the lack of available data on the diving behavior of the Longman's beaked whale, it will be represented by a surrogate species: the Blainville's beaked whale (Section 2.2.1.8.1). Though originally placed in the *Mesoplodon* genus, differences in features of the skull have led to the reclassification of this species into its own genus, *Indopacetus*; the previous classification into the same genus, and the current classification into the same family, is the primary reason for using Blainville's beaked whale as the surrogate species. The depth distribution for the Longman's beaked whale can be found in Table 25.

2.2.1.8.3 HSTT Beaked Whale Guild, *Mesoplodon* spp.

Within the Southern California portion of the HSTT Study Area, a beaked whale guild has been established. This guild was created to account for the difficulties associated with identifying beaked whales to the species level at sea. The beaked whale guild includes: Blainville's beaked whale, Ginkgo-toothed beaked whale (*Mesoplodon ginkgodens*), Stejneger's beaked whale (*Mesoplodon stejnegeri*), Hubb's beaked whale (*Mesoplodon carlshubbi*), Perrin's beaked whale (*Mesoplodon perrini*), and pygmy beaked whale

(*Mesoplodon peruvianus*). The depth distribution used for this group is that of the Blainville's beaked whale. The depth distribution for the beaked whale guild in the HSTT Study Area can be found in Table 25.

2.2.1.8.4 AFTT Beaked Whale Guild

Within the AFTT Study Area, all beaked whales are represented as one group due to the difficulty of identifying beaked whales at sea to the individual species level. The AFTT beaked whale guild is a composite of all species that would be present in the Study Area; including Cuvier's beaked whale (*Ziphius cavirostris*), Blainville's beaked whale, Sowerby's beaked whale (*Mesoplodon bidens*), Gervais' beaked whale (*Mesoplodon europaeus*), and True's beaked whale (*Mesoplodon mirus*). While both Blainville's and Cuvier's beaked whales are both located within the AFTT Study Area, the depth distribution for Blainville's beaked whales (Section 2.2.1.8.1) was used as the representative species for this group. Blainville's beaked whales spend a greater percentage of time in the upper 100 m of water than Cuvier's, making the model a more conservative estimate of affected beaked whales. The depth distribution of beaked whales in the AFTT study area can be found in Table 25.

2.2.1.8.5 *Ziphius cavirostris*, Cuvier's Beaked Whale

Cuvier's beaked whales inhabit slope waters with steep gradients around the world's oceans, with the exception of the high polar seas (Heyning 1989). Stomach contents analyses indicate that prey species include mesopelagic and benthic cephalopods, fish, and crustaceans (Heyning 1989; Hickmott 2005; Santos et al. 2001a). It appears however, that Cuvier's beaked whales eat mostly squid, and the majority of prey are open-ocean species that occur well below the surface, including on or near the seafloor in deep waters (Reeves et al. 2002).

In order to build a representative depth distribution for the Cuvier's beaked whale (Table 26), data from Figure 1 from Aguilar de Soto et al. (2006), Figures 5 from Baird et al. (2005b), Figure 3c from Baird et al. (2006), Figure 2a from Baird et al. (2008), Figure 1 from Barlow et al. (2013), DTAG data from Johnson and Sturlese (Johnson and Sturlese 2008a, 2008b), Figure 1a from Kvadsheim et al. (2012), and Figures 2a and 3a from Schorr et al. (2014). Aguilar de Soto et al. (2006) presented a time-depth profile of a Cuvier's beaked whale off Italy over a 15.6 hour period. Baird et al. (2005b) presented the cumulative percentage of time spent at depth for an adult female during the day and at night. Similarly, Baird et al. (2008) looked at diel variation in Cuvier's beaked whale diving behavior, presenting the cumulative percentage of time spent at depth for two tagged whales during both the day and night. Barlow et al. (Barlow et al. 2013) used DTAGs to collect acoustic information for beaked whales in an attempt to estimate density and abundance; Figure 1 is a typical dive profile for a tagged whale captured over a 15 hour period. Raw DTAG data collected on multiple occasions for an animal in Liguria, Italy, provided by Johnson and Sturlese (Johnson and Sturlese 2008a, 2008b), were binned. Kvadsheim et al. (2012) presented changes in dive behavior in response to sonar; as this report is portraying normal Cuvier's beaked whale behavior, the portion of the dive profile provided by Kvadsheim et al. (2012) in which sonar was used has been omitted from the typical depth distribution calculated. Schorr et al. (2014) provided multi-day tag data for Cuvier's beaked whales, allowing for observation of diel patterns in dive behavior. The depth distributions from these studies were averaged together to create the representative depth distribution below. The depth distribution for Cuvier's beaked whales is given in Table 26.

Table 26. Percentage of Time at Depth for Cuvier’s Beaked Whale^{1,2}

Depth Bin (m)	% of Time at Depth
0–100	31.9
100–200	10.3
200–300	11.7
300–400	5.5
400–500	4.1
500–600	3.9
600–700	5.1
700–800	4.1
800–900	3.4
900–1,000	4.8
1,000–1,100	4.3
1,100–1,200	3.3
1,200–1,300	2.2
1,300–1,400	1.4
1,400–1,500	1.1
1,500–1,600	0.3
1,600–1,700	0.9
1,700–1,800	0.5
1,800–1,900	0.8

¹Based on data from Aguilar de Soto et al. (2006), Baird et al. (2005b), Baird et al. (2006), Baird et al. (2008), Barlow et al. (2013), Johnson and Sturlese (Johnson and Sturlese 2008a, 2008b), Kvadsheim et al. (2012), and Schorr et al. (2014)

²This depth distribution is also representative of the Baird’s beaked whale and the Northern bottlenose whale

This representative depth distribution is consistent with foraging dives from 689 to 1,888 m in the Mediterranean Sea (Tyack et al. 2006) and to 1,450 m off Hawaii (Baird et al. 2006). Based on the representative depth distribution in Table 26, Cuvier’s beaked whales spent 31.9 percent of their time between 0–100 m and 36.1 percent of time deeper than 500 m; these values remain consistent with the data reported by Baird et al. (2008), in which Cuvier’s beaked whales spend 12.4–51.1 percent of their time spent at depths less than 50 m, and between 33.9–52.1 percent of time at depths greater than 500 m.

2.2.1.8.6 *Berardius bairdii*, Baird’s Beaked Whale

Baird’s beaked whales inhabit temperate waters of the North Pacific Ocean and adjoining seas, primarily in the deep waters offshore of the continental shelf (Balcomb 1989; Kasuya 1986). This species consumes benthic and epibenthic fish and cephalopods, and occasionally feeds on mesopelagic species as well (Balcomb 1989; Kasuya 2002; Walker et al. 2002).

Little data have been collected on the dive behavior of Baird’s beaked whales. Stomach contents analysis suggests that whales are feeding at depths of 800–1,200 m off Japan, feeding on prey at or near the seafloor (Reeves et al. 2002; Walker et al. 2002). Minamaka (2007) reported that one animal carrying a time-depth recorder dove down to a maximum depth 1,777 m, with dives lasting up to 64.4 min, which is similar to the maximum dive duration of 67 min observed by Kasuya (1986). The maximum dive depth reported by the first deployment of a multi-sensor tag on this species was given as roughly 1400 m (Stimpert et al. 2014).

Due to the lack of data on the diving behavior of the Baird’s beaked whale, it will be represented by a surrogate species: Cuvier’s beaked whale (Section 2.2.1.8.5). The Cuvier’s beaked whale is also a member of the subfamily Ziphiidae, and the feeding habits and types of prey for the two species are similar. The diving pattern of Baird’s beaked whales appears very similar to other beaked whales, in which a long duration, deep

dive is followed by shorter duration, shallow dives (Minamikawa et al. 2007; Tyack et al. 2006). The depth distribution for the Baird's beaked whale can be found in Table 26.

2.2.1.8.7 *Hyperoodon ampullatus*, Northern Bottlenose Whale

Northern bottlenose whales are found in deep, cold temperate and subpolar waters of the North Atlantic Ocean (Gowans 2002). The primary prey of bottlenose whales is squid, principally those species of the genus *Gonatus* (Hooker et al. 2001; Santos et al. 2001b). They also eat herring, various deep sea fish, shrimp, and sometimes sea cucumbers and starfish (Reeves et al. 2002).

Little data have been collected on the dive behavior of Northern bottlenose whales. Hooker and Baird (1999) reported on the diving behavior of two individuals from Canadian waters. Both individuals routinely dove deeper than 800 m, reaching a maximum dive depth of 1,453 m.

Due to the lack of available data on the diving behavior of the Northern bottlenose whale, it will be represented by a surrogate species: the Cuvier's beaked whale (Section 2.2.1.8.5). Cuvier's beaked whales are also deep-diving members of the family Ziphiidae, and are of similar size to female Northern bottlenose whales (Leatherwood and Reeves 1983). Additionally, the primary prey for the Northern bottlenose whale is squid, as is the primary prey for Cuvier's beaked whales. The depth distribution for the Northern bottlenose whale can be found in Table 26.

2.2.2 Carnivores

2.2.2.1 Family Otariidae

2.2.2.1.1 *Arctocephalus townsendi*, Guadalupe Fur Seal

The Guadalupe fur seal breeds at only two rookery locations: the east side of Isla Guadalupe and the southeast side of Isla Bonito del Este (Maravilla-Chavez and Lowry 1999). Single animals have occasionally been seen at islands along the coast of Baja California and California (Stewart et al. 1987). Squid, teleost fish, and crustaceans comprise the diet of the Guadalupe fur seal (Auriolles-Gamboa and Camacho-Ríos 2007; Belcher and Lee 2002; Riedman 1990).

Little data have been collected on the dive behavior of Guadalupe fur seals. In order to build a representative depth distribution for Guadalupe fur seals, data from Figure 2 in Lander et al. (2000) were used. Lander et al. (2000) obtained information on the diving behavior of Guadalupe fur seals by satellite tagging a single female following rehabilitation off southern California. Most dives were performed at night and lasted 2–4 min (mean dive time of 1.7 min, SD=1.0 min), with a maximum dive duration of 18 min (Lander et al. 2000). The mean dive depth was 15.7 m (SD=11.8 m). The maximum recorded dive depth was 130 m, although most dives were less than 20 m deep. While Lander et al. (2000) reported only the number of dives to various depth bins, this is used as a proxy for the time spent at depth. The bins over 40 m were presented together as they represented a small percentage of time at depth. The depth distribution for Guadalupe fur seals is given in Table 27.

Table 27. Percentage of Time at Depth for the Guadalupe Fur Seal¹

Depth Bin (m)	% of Time at Depth
0–20	92.0
20–40	7.0
40–160	1.0

¹Based on data from Lander et al. (2000)

Data from Gallo-Reynoso et al. (2008) also studied the diving behavior of one adult female, finding a mean dive depth of 17 m (SD=10 m) with an average dive duration of 2.6 min (SD=1.4 min).

2.2.2.1.2 *Callorhinus ursinus*, Northern Fur Seal

Northern fur seals occupy the pelagic waters of the North Pacific Ocean, Bering Sea, and Sea of Japan, ranging coastally as far south as Baja California, Mexico and Japan, and with an at-sea southern limit around 35°N (Gelatt and Lowry 2008). Northern fur seals are known to feed in the deep waters along the continental shelf break, as well as shallower waters of the shelf itself (Gentry 2002; Ponganis et al. 1992). The diet of northern fur seals varies regionally and seasonally, but is comprised principally of finfish (e.g., Pacific herring, sand lance, capelin, myctophids) and squid; they will occasionally feed upon other prey such as birds and crustaceans (Ream et al. 2005; Riedman and Estes 1990).

In order to build a representative depth distribution for the Northern fur seals, data from Table 1 in Kooyman et al. (1976) were used. The dive behavior and physiology of the northern fur seal was among the earliest pinniped species to be tagged and tracked (Kooyman et al. 1976). Table 1 indicates the number of dives to specific depth bins; thus this data from Kooyman et al. (1976) will be used as a proxy for percentage of time spent at depth. Kooyman et al. (1976) found that shallow dives (0–20 m) lasted less than 1 min in duration. Deeper dives, from 20–140 m, lasted from 2–5 min. The maximum dive depth in this study was 190 m. The average interval between dives was 17 min. The depth distribution for Northern fur seals is given in Table 28.

Table 28. Percentage of Time at Depth for the Northern Fur Seal¹

Depth Bin (m)	% of Time at Depth
0–20	48.36
20–50	42.32
50–80	5.94
80–110	0.96
110–140	1.99
140–170	0.39
170–200	0.04

¹Based on data from Kooyman et al. (1976)

While other studies did not include depth distributions for northern fur seals, they did provide additional information to categorize dive behavior. Because northern fur seals spend the vast majority of their lives at sea (87–90 percent of the year), only coming ashore to breed for 35–45 days from June to August, most of the tagging studies examine the dive behavior of females on excursions from breeding colonies in the Bering Sea (Baker 2007; Baker and Donohue 2000; Gentry 2002; Goebel et al. 1991; Kooyman et al. 1976; Kuhn et al. 2009; Ream et al. 2005). In general, adult females on foraging excursions generally follow one of three dive profiles: shallow, deep, or mixed depth. Shallow-diving seals show a crepuscular pattern with dive depths varying according to movement of the deep scattering layer. Deep-diving seals show no temporal pattern, apparently ignoring the diel movements of vertically migrating prey, and have no consistent change in depth within bouts. Mixed-depth divers alternate between dive profiles, perhaps shifting prey types (Gentry et al. 1986). Female northern fur seals dive mostly at night (68 percent) (Gentry et al. 1986). Individuals may be consistent in their diving behavior, presumably choosing prey sources at different depths, as evidenced by unique fatty acid profiles specific to a differentiated prey type (deep versus vertically migrating species, for example); however, other evidence points towards a seasonal shift in dive behavior (Gentry 2002; Gentry et al. 1986; Hobson and Sease 1998; Hobson et al. 1997). Radio-tracking studies suggest that deep-diving patterns are used while foraging on the continental shelf, while shallow-diving patterns occur over deeper waters off the shelf break or in the Aleutian Basin (Gentry et al. 1986; Goebel et al. 1991).

Female northern fur seals have been recorded diving to a maximum depth of 256 m (Ponganis et al. 1992), although they most frequently dive between 50 and 60 m (Gentry et al. 1986). The activity budgets for adult females on foraging trips was 17 percent of time spent resting, 26 percent of time spent diving, and 57 percent of time spent in surface active time (Gentry et al. 1986).

Male northern fur seal diving behavior has also been examined (Sterling and Ream 2004) and has revealed that like parturient females, juvenile males exhibit shallow versus deep dive patterns based upon foraging location (deep dives in water less than 200 m deep, shallow nighttime dives in waters up to 3,000 m deep). In a study of 19 juvenile male northern fur seals on foraging excursions during the breeding season in the Bering Sea, the maximum recorded dive depth was 175 m, with a mean dive depth of 17.5 m (SD=1.5 m) (Sterling and Ream 2004).

2.2.2.1.3 *Zalophus californianus*, California Sea Lion

California sea lions primarily breed on island beaches off southern California, along Baja California, Mexico, and in the Gulf of California (Heath 2002). They eat a variety of prey, including schooling fish, crustaceans, and cephalopods (García-Rodríguez and Aurióles-Gamboa 2004; Melin et al. 2008; Porras-Peters et al. 2008). California sea lions often make shallow dives in some coastal areas for epipelagic prey, and deeper dives in others, such as the Gulf of California, for mesopelagic prey (Costa et al. 2004; Lowry and Carretta 1999; Melin and DeLong 1999).

In order to build a representative depth distribution for California sea lions, data from Figure 4 in Feldkamp et al. (1989) and Figure 1a in Weise (2006) were used. Both Figures show the number of dives to specific depth bins, thus the data from Feldkamp et al. (1989) and Weise (2006) will be used as a proxy for percentage of time spent at depth. Because data indicate differences in diving behavior of males (Weise 2006) and females (Feldkamp et al. 1989; Kuhn 2006; Melin et al. 2008), data from Weise (2006) and Feldkamp et al. (1989) were averaged to create a composite distribution. Weise (2006) reported an average dive depth of 32 m, but that most dives occurred within the depth range of 10 to 20 m. Feldkamp et al. (1989) found that the maximum dive depth for females was roughly 274 m. The mean depths ranged from 31 to 98 m, but the majority of dives went to between 20 and 50 m. In order to account for the time spent at the surface, an activity budget from Table 3 in Thomas et al. (2010) was used, incorporating both male and female sea lions. Based on these numbers, a percent time swimming at the surface (63.62 percent) was added into the average taken from the Feldkamp et al. (1989) and Weise (2006) data. The remaining averages from both studies were redistributed proportionally to account for the remaining 36.38 percent of time. The resulting depth distribution profile includes data from all three studies. The depth distribution for California sea lions is given in Table 29.

Table 29. Percentage of Time at Depth for the California Sea Lion¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–10	68.39	120–130	0.56
10–20	5.92	130–140	0.40
20–30	6.01	140–150	0.46
30–40	4.86	150–160	0.53
40–50	4.00	160–170	0.46
50–60	2.31	170–180	0.33
60–70	1.49	180–190	0.15
70–80	1.02	190–200	0.13
80–90	0.75	200–210	0.22
90–100	0.69	210–220	0.11
100–110	0.73	220–230	0.05
110–120	0.60	230–500	0.22

¹Based on data from Feldkamp et al. (1989), Weise (2006), and Thomas (2010)

While other studies did not include usable depth distributions for California sea lions, they did provide additional information to categorize dive behavior. Mean dive depths reported for female California sea lions range from 45–70 m with maximum dive depths ranging from 104–279 m (Kuhn 2004; Kuhn 2006; Melin et al. 2008). While this data set does not necessarily capture the maximum dive depths reported in several locations (greater than 200 m) (Costa et al. 2004; Feldkamp et al. 1989; Kuhn 2004; Kuhn 2006; Melin et al. 2008), the depth distribution is consistent with studies by Kuhn (2004) and Melin (2008), both of whom demonstrated that the majority of dives (ranging from 55–85 percent) were less than 50 m deep. Similarly, Melin and DeLong (1999) reported that most dives were shallower than 100 m. Therefore, although the depth distribution in Table 29 may be shallower than seen in some locations, it does capture published diving behavior for the species, and is representative for California sea lions.

2.2.2.2 Family Phocidae

2.2.2.2.1 *Cystophora cristata*, Hooded Seal

Hooded seals typically inhabit continental shelf and slope waters of the North Atlantic Ocean. Adults feed on commercial-size fish such as capelin, cod, and redfish, as well as squid and amphipods (Bjørke 2001; Haug et al. 2004; Hauksson and Bogason 1997).

In order to build a representative depth distribution for hooded seals, both the text and data from Figures 2A-L in Folkow and Blix (1999) were used. Folkow and Blix (1999) reported tag data from 16 seals from the Greenland Sea. The seals had an average submersion time of 77 percent. While not an estimate of time spent at depth, it is the closest approximation to a depth distribution in the published literature. Due to the lack of data on time spent at depth, the data from Folkow and Blix (1999) will be used as a proxy for percentage of time spent at depth. Assuming a submersion time of 77 percent gives us a remaining surface time of 23 percent, which has been included as the 0–1 m bin below. The remaining distribution bins were taken from the data in the text of Folkow and Blix (1999), which is also displayed in Figures 2A through 2L. The remaining bins from the Folkow and Blix (1999) depth distribution were redistributed proportionally to account for the remaining 77.0 percent of time. The depth distribution for hooded seals is given in Table 30.

Table 30. Percentage of Time at Depth for the Hooded Seal¹

Depth Bin (m)	% of Time at Depth
0–1	23
1–52	13.09
52–100	6.16
100–600	54.67
600–968	1.925
968–1,016	1.155

¹Based on data from Folkow and Blix (1999)

While other studies did not include usable depth distributions for hooded seals, they did provide additional information to categorize dive behavior. Bajzak et al. (2009) reported dive depths for post-breeding seals consistent with those from Folkow and Blix (1999), although the seals dove to shallower depths during the molting period. Seals from the Gulf of St. Lawrence spent a similar percentage of their time diving, but considerably less during the molting period (Bajzak et al. 2009). However, the depth distribution in from Folkow and Blix (1999) represents dive depths from seals in waters of different depths; therefore this depth distribution represents the best representative depth distribution.

2.2.2.2 *Monachus schauinslandi*, Hawaiian Monk Seal

The range of the Hawaiian monk seal is limited to the central Pacific Ocean, with breeding colonies principally in the northwestern Hawaiian Islands (Gilmartin and Forcada 2002; Johanos and Baker 2002). Hawaiian monk seals primarily consume benthic prey; stomach contents and scat analyses indicate that prey constitute both diurnal and nocturnal species. Reef associated fish and octopus compose a large portion of the diet (DeLong et al. 1984; Goodman-Lowe 1998; Kenyon and Rice 1959). Most seals focus foraging efforts in the top 100 m of the water column, although some seals dive to greater than 300 m deep (Parrish et al. 2002; Parrish et al. 2000; Parrish et al. 2005; Stewart et al. 2006).

In order to build a representative depth distribution for Hawaiian monk seals, data from two Stewart and Yochem (2004a, 2004b) studies were used. Figure 25 from Stewart and Yochem (2004a) and Figure 40 from (2004b) provided time at depth for weaned pups, juveniles, and adults that could be averaged for a representative depth distribution. Stewart and Yochem (2004a, 2004b) presented depth distributions for 18 animals from Kure Atoll and Laysan Island, respectively. The maximum dive depth reported was greater than the recording limit of the tag at 490 m, therefore, 500 m is used as a conservative maximum dive depth. Most dives of all seals were shallower than 40 m and lasted less than 6–8 min (Stewart and Yochem 2004a, 2004b). The depth distribution for Hawaiian monk seals is given in Table 31.

Table 31. Percentage of Time at Depth for the Hawaiian Monk Seal¹

Depth Bin (m)	% of Time at Depth
0–4	33.0
4–20	34.68
20–40	13.2
40–60	5.45
60–80	3.58
80–100	2.08
100–120	2.53
120–140	2.0
140–160	0.75
160–180	0.69
180–200	0.25
200–250	0.38
250–350	0.88
350–500	0.56

¹Based on data from Stewart and Yochem (2004a, 2004b)

While other studies did not include usable depth distributions for Hawaiian monk seals, they did provide additional information to categorize dive behavior. In the representative depth distribution above, 92 percent of the monk seals' time is spent within the top 100 m. This is consistent with previous studies that have demonstrated Hawaiian monk seal foraging occurring at shallow reef sites (DeLong et al. 1984; Littnan et al. 2004; Parrish et al. 2000; Schlexer 1984).

2.2.2.2.3 *Mirounga angustirostris*, Northern Elephant Seal

Northern elephant seals are limited to the North Pacific Ocean, with breeding haul outs located along the western coast of North America from northern California to Baja California, Mexico. Seals utilize deep waters for foraging, traveling north and west from breeding beaches, and traveling as far north as the Gulf of Alaska and Aleutian Islands (Hindell 2002). Elephant seals feed primarily on vertically migrating epipelagic and mesopelagic squid, but eat a variety of prey species, including elasmobranchs, crustaceans, cephalopods, and fish (DeLong and Stewart 1991; Sinclair 1994; Stewart and Huber 1993).

In order to build a representative depth distribution for Northern elephant seals, data from DeLong and Stewart (1991), Hakoyama et al. (1994), Le Boeuf et al. (1986), and Naito et al. (1989) were used. In addition, a surface time from DeLong and Stewart (1991) was incorporated into the data. The text of DeLong and Stewart (1991) states that Northern elephant seals are submerged 86 percent of the time, leaving an above-surface time percentage of 14 percent (DeLong and Stewart 1991). Due to the lack of data on time spent at depth, the data from these sources, citing percentage of dives to specific depth bins will be used as a proxy for percentage of time spent at depth. As the data show sex differences in foraging and diving behavior (DeLong and Stewart 1991; Hakoyama et al. 1994; Le Boeuf et al. 1986; Le Boeuf et al. 1993; Naito et al. 1989), males and females were averaged together to create a representative depth distribution. In males, the data averaged from Figure 1 in DeLong and Stewart (1991) and Figure 4 in Hakoyama et al. (1994) showed that 83 percent of dives took place between the surface to 500 m, most frequently occurring from 300–500 m. In females, the data averaged from Figure 1 in Le Boeuf et al. (1986), Figure 2 in Naito et al. (1989), and Figure 4 in Hakoyama et al. (1994), showed that 85 percent of dives were to depths of 300–700 m, most frequently from 400–600 m. For males, dives deeper than 500 m were unusual, whereas dives over 700 m were rare for females (DeLong and Stewart 1991; Hakoyama et al. 1994; Le Boeuf et al. 1988; Le Boeuf et al. 1986; Naito et al. 1989). Data from each Figure were summed into 100 m bins and averaged together. The

remaining bins of each study were averaged and redistributed proportionally to account for the remaining 86 percent of non-surface time. The resulting depth distribution profile includes data from all studies. The depth distribution for Northern elephant seals is given in Table 32.

Table 32. Percentage of Time at Depth for the Northern Elephant Seal¹

Depth Bin (m)	% of Time at Depth
0–100	21.4
100–200	7.65
200–300	8.86
300–400	13.42
400–500	18.06
500–600	16.34
600–700	9.98
700–800	2.15
800–900	0.77
900–1000	0.52
1,000–1,100	0.52
1,100–1,200	0.43

¹Based on data from DeLong and Stewart (1991), Hakoyama et al. (1994), Le Boeuf et al. (1986), and Naito et al. (1989)

While other studies did not include usable depth distributions for northern elephant seals, they did provide additional information to categorize dive behavior. Robinson et al. (2012) found that the dive depths of most seals showed a clear diel pattern, consistent with targeting vertically migrating prey species. Robinson et al. (2012) found the mean dive depth to be 516 m and a mean dive duration of 23.1 min. Active-bottom dives made up the greatest percentage of dives (54.0 percent) in this study (Robinson et al. 2012). Maximum dive depths for elephant seals covered a wide range of depths, from under 1,000 m (Crocker et al. 2006; Davis et al. 2001; Davis and Weihs 2007; Le Boeuf et al. 1988) to the deepest dive ever recorded by an elephant seal at 1,747 m (Robinson et al. 2012).

2.2.2.2.4 *Halichoerus grypus*, Grey Seal

Grey seals are found in temperate, coastal waters of the North Atlantic Ocean. Grey seals feed on both pelagic (herring and sand lance) and benthic (flatfish) prey (Bowen et al. 1993; Iverson et al. 2004).

In order to build a representative depth distribution for grey seals, data from the text of Harvey et al. (2008), a maximum dive depth from Beck et al. (2003), as well as an activity budget from Thomson et al. (1991), were used. Harvey et al. (2008) presented a distribution of maximum dive depths for a range of age, sex, and seasonal classes, which, while not an exact representation of time spent in each depth range, is a good approximation as grey seals are primarily shallow divers. Due to the lack of data on time spent at depth, the data from Harvey et al. (2008) will be used as a proxy for percentage of time spent at depth. However, the data from Harvey et al. (2008) only consider when the animal is beneath the surface. Thompson et al. (1991) tracked three swimming juvenile grey seals and reported that across three behavioral states (traveling, short trips, and resting), animals spent on average 84.1 percent of their time submerged. Therefore, the average time spent at the surface from Thompson et al. (1991) was combined with the dive distribution data from Harvey et al. (2008) to generate an estimate of the depth distribution for the grey seal. The maximum dive depth of 412 m reported by Beck et al. (2003) is used as the maximum depth. The depth distribution for grey seals is given in Table 33.

Table 33. Percentage of Time at Depth for the Grey Seal¹

Depth Bin (m)	% of Time at Depth
0–40	81.7
40–100	14.2
100–200	3.3
200–412	0.8

¹Based on data from Harvey et al. (2008), Beck et al. (2003), and Thompson et al. (1991)

While other studies did not include usable depth distributions for grey seals, they did provide additional information to categorize dive behavior. Beck et al. (2003) reported that 73.9 percent and 59.4 percent of dives were in the top 60 m during the day and night, respectively, and that 95.9 percent of dives were in the top 120 m. Lidgard et al. (2003) found that 41 percent of dives were within 10 m of the surface.

2.2.2.2.5 *Phoca vitulina*, Harbor Seal

Harbor seals are found in shallow inshore and coastal waters of the Northern Hemisphere (Burns 2002). Prey species include epibenthic and benthic fish (e.g., sandlance, flounder, and herring) and squid (Brown and Mate 1983; Olesiuk 1993; Payne and Selzer 1989).

In order to build a representative depth distribution for harbor seals, data from Table 3 in Womble et al. (2014) and an activity budget from Bowen et al. (1999) were used. Womble et al. (2014) studied the dive behavior of 12 female harbor seals in Alaska. Table 3 shows the percentage of dives to specific depth bins; thus, this data from Womble et al. (2014) will be used as a proxy for percentage of time spent at depth. Seals in this study were found to dive most frequently (81.6 percent) to depths shallower than 50 m. It should be noted that this study cites the percentage of dives to certain depths, as opposed to the percentage of time spent in each depth bin, including time spent in that bin on the way to a deeper bin. However, the Womble et al. (2014) data are the best available estimate of time spent at depth. The activity budget from Bowen et al. (1999) provided a surface time of 85.01 percent. The remaining bins from Womble et al. (2014) were redistributed proportionally to account for the remaining 14.99 percent of non-surface time. The resulting depth distribution profile includes data from both studies. The depth distribution for harbor seals is given in Table 34.

Table 34. Percentage of Time at Depth for the Harbor Seal in the HSTT Study Area¹

Depth Bin (m)	% of Time at Depth
0-4	85.01
4–50	12.22
50–100	2.32
100–150	0.37
150–200	0.05
200–250	0.01
250–300	0

¹Based on data from Womble et al. (2014) and Bowen et al. (1999)

While other studies did not include usable depth distributions for harbor seals, they did provide additional information to categorize dive behavior. Gjertz et al. (2001) reported the maximum depth reached by harbor seals to be within the 200–350 m range, however, additional studies have reported shallower maximum dive depths, reaching less than 100 m off Nova Scotia (Bowen et al. 1999), Svalbard (Jorgensen et al. 2001), and in Prince William sound (Frost et al. 2001). Hastings et al. (2004) observed that less than six percent of harbor seals dive to depths greater than 100 m, although one seal dove to 508 m (Hastings et al. 2004). Harbor seals

near Svalbard, Norway, dove to a maximum depth of 452 m, although 50 percent of dives were shallower than 40 m, and 95 percent were to less than 250 m (Gjertz et al. 2001). Eguchi and Harvey (2005) observed that males dive deeper than females, with males diving to depths shallower than 154 m and females diving to depths shallower than 76 m 95 percent of the time.

2.2.2.2.6 Seal Guild in the AFTT Study Area

Within the AFTT Study Area, seals are often difficult to distinguish when viewing at sea. Many surveyed species are listed as “unidentified.” Therefore, the density data from these surveys have been combined for the two most abundant species of seals in the AFTT Study Area, grey seals (Section 2.2.2.2.4) and harbor seals (Section 2.2.2.2.5). As a result, the depth distributions for the two species from Table 33 and Table 34 were averaged together using the source data from Womble et al. (2014), Harvey et al. (2008), Beck et al. (2003), Bowen et al. (1999) and Thomson et al. (1991). In order to average these two species together to create a representative depth distribution for seals in the AFTT Study Area. While the surface bin for harbor seals (0–50 m) was slightly larger than the surface bin for gray seals (0–40 m), they were averaged together to create a composite depth distribution for these two species. The resulting representative depth distribution for seals in the AFTT Study Area is given in Table 35.

Table 35. Percentage of Time at Depth for Seals in the AFTT Study Area¹

Depth Bin (m)	% of Time at Depth
0–50	89.47
50–100	8.26
100–200	1.86
200–412	0.41

¹Based on Womble et al. (2014), Harvey et al. (2008), Beck et al. (2003), Bowen et al. (1999), and Thomson et al. (1991)

2.2.2.2.7 *Pagophilus groenlandicus*, Harp Seal

Harp seals are found in ice-associated waters of the Arctic (Lavigne 2002). Harp seals forage on a variety of prey, including Arctic cod and capelin, amphipods, and krill (Haug et al. 2004; Lawson et al. 1998; Nilssen et al. 2001).

Given the characteristic shallow diving pattern in the post-breeding to post-molting period (spring), two depth distributions are given in Table 36 for the harp seal. Folkow et al. (2004) characterized diving behavior of 16 adult harp seals throughout the year in two different years. In order to build a representative depth distribution for harp seals during the breeding and molting season, data from Figure 5A in Folkow et al. (2004) were used. In order to build a representative depth distribution for harp seals during the remainder of the year, data from Figure 7C of Folkow et al. (2004) were used. This study reported that after the period of time close to the ice edge, the seals then moved into deeper water in the open ocean, diving up to 492 m and 568 m in the different years. While close to the ice edge during the breeding and molting season in summer, seals spent most of their time in shallow depths, with 70-90 percent of their diving time spent in the top 50 m of the water column (Folkow et al. 2004). A large proportion of the seals (9 out of a total of 14; both years) at some time dove deeper than 450 m, and dives deeper than 300 m represented greater than 12 percent of all dives. There were significant seasonal variations in diving depths, with dives being significantly shallower in spring/summer than in autumn and winter (Folkow et al. 2004). The depth distribution for harp seals is given in Table 36.

Table 36. Percentage of Time at Depth for the Harp Seal¹

Depth Bin (m)	% of Time at Depth	
	Post-breeding to Post-molting (Spring)	Remainder of Year (Summer, Winter, Fall)
0–20	52	32
20–50	29	17
50–100	12	16
100–200	5	18
200–300	2	11
300–400	0	5
400–492	0	1

¹Based on data from Folkow (2004)

While other studies did not include usable depth distributions for harp seals, they did provide additional information to categorize dive behavior. Lydersen and Kovacs (1993) found that lactating females had maximum dive depths of 90 m, which is consistent with the majority of seals in the data above. Pacific cod, one of the dominant prey types found in seals feeding offshore, inhabit the top 300 m of the water column, consistent with the diving depths in Table 36 (Wathne et al. 2000).

2.2.2.8 *Phoca hispida*, Ringed Seal

Ringed seals have a circumpolar distribution within Arctic waters. Primary prey species include arctic cod, amphipods, and crustaceans (Labansen et al. 2007; Wathne et al. 2000).

In order to build a representative depth distribution for ringed seals, data from Figure 3 in Gjertz et al. (2000) were used, along with a surface bin taken from Lydersen (1991). Gjertz et al. (2000) reported the percentage of dives to different depths for seven ringed seals. This data from Gjertz et al. (2000) will be used as a proxy for percentage of time spent at depth. Gjertz et al. (2000) found that 48 percent of all dives were shallower than 20 m and 90 percent of dives were shallower than 100 m. Because percentage of time spent at the surface was not calculated in Gjertz et al. (2000), the Lydersen data (1991) were used to estimate this time. The Lydersen (1991) data are from a single female ringed seal over a six day period. If the percentage of time at the surface (30.3 percent) from Lydersen (1991) is used as an estimate, and the percentage of dives taken as a proxy for the percentage of time, then the following representative depth distribution results. The remaining bins of the Gjertz study were redistributed proportionally to account for the remaining 69.7 percent of non-surface time. The resulting depth distribution profile includes data from both studies. The depth distribution for ringed seals is given in Table 37.

Table 37. Percentage of Time at Depth for the Ringed Seal¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–1	30.3	30–40	4.88
1–4	9.76	40–50	4.18
4–8	13.24	50–100	14.64
8–12	5.23	100–150	5.58
12–16	3.49	150–200	1.39
16–20	1.39	200–300	0.35
20–30	5.58		

¹Based on data from Gjertz et al. (2000) and Lydersen (1991)

These dive depths are consistent with Wathne et al. (2000), who found that ringed seals feed on shallower prey than harp seals, which make 35 percent of their dives to deeper than 100 m.

2.2.3 Sirenians

2.2.3.1 Family Trichechidae

2.2.3.1.1 *Trichechus manatus*, West Indian Manatee

The West Indian manatee is found in coastal and estuarine waters of the mid-Atlantic United States south through the waters of central Brazil (Reynolds III and Powell 2002). Manatees are primarily herbivores, feeding on plants in all parts of the water column (Alves-Stanley et al. 2010; Baugh et al. 1989; Lefebvre et al. 2000). There are occasional reports of manatees feeding on fish and invertebrate prey (Courbis and Worthy 2003; Etheridge et al. 1985; O'Shea et al. 1991). Manatees are limited to shallow water (less than 20 m), due to the light requirements of the vegetation on which they feed (Wells et al. 1999).

In order to build a representative depth distribution for manatees, data from Figure 8 in Nowacek et al. (2002) and Figure 3 in Edwards et al. (2016) were used. Nowacek et al. (2002) reported the percent time at depth for two manatees in a central Belize lagoon. The manatees were located in water that ranged from 1.5–10.5 m deep. The average time spent in portions of the water column was 80 percent in less than 3 m and 57 percent in less than 1.5 m of water. The Edwards et al. (2016) study was conducted during winters in Tampa Bay, Florida from 2002–2005, using tag data from nine manatees. In the area of the study, the water depth averages 4 m, but reaches a maximum of 27 m. Mean dive depth of these manatees was 1.09 m (SD= 0.17) with a maximum dive depth ranging from 6.0–16.2 m. Manatee diving behavior showed a diel pattern, diving deeper during the day and shallower at night (Edwards et al. 2016). The data from Nowacek et al. (2002) and Edwards et al. (2016) were averaged together to create the representative depth distribution for West Indian manatees. The depth distribution for West Indian manatees is given in Table 38.

Table 38. Percentage of Time at Depth for the West Indian Manatee¹

Depth Bin (m)	% of Time at Depth
0–1.0	46.43
1.1–2.0	31.38
2.1–3.0	10.83
3.1–4.0	10.83
4.1–16.2	0.50

¹Based on data from Nowacek et al. (2002) and Edwards et al. (2016)

2.2.4 Sea Turtles

2.2.4.1 Family Dermochelyidae

2.2.4.1.1 *Dermochelys coriacea*, Leatherback Sea Turtle

The leatherback turtle is globally distributed in tropical, subtropical, and warm-temperate waters throughout the year, and throughout cooler temperate waters during warmer months (James et al. 2005a; National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 1993). Adult leatherback turtles forage in temperate and subpolar regions in all oceans and migrate to tropical nesting beaches between 30° N and 20° S (Bleakney 1965; Brongersma 1972; Goff and Lien 1988; Threlfall 1978). The leatherback is typically associated with continental shelf habitats and pelagic environments. To a large extent, the oceanic distribution of leatherbacks may reflect the distribution and abundance of their favored prey, macro-planktonic soft-bodied animals such as jellyfish, salps, and pyrosomes (Wallace et al. 2013). It is

suggested that leatherbacks make scouting dives while transiting as an efficient means for sampling prey density and perhaps also to feed opportunistically at these times (James et al. 2006b; Jonsen et al. 2007).

In order to build a representative depth distribution for leatherback turtles, data from Figure 2 as well as the text of Houghton et al. (2008) were used. Houghton et al. (2008) tagged 13 adult leatherback sea turtles at two sites over the course of four years. The data in Figure 2 accounts only for the percentage of dives made to each depth bin, rather than the percent of time spent in each bin. The text states the percent of dives made to less than 10 m (18.9 percent), which is considered the surface bin. This representative depth distribution was created to account for several behavioral states, including post-nesting, migration, and foraging. The mean dives in this study ranged from 32.5–69.0 m. Houghton et al. (2008) found that 99.6 percent of dives were to depths shallower than 300 m with only 0.4 percent extending to greater depths. These findings support the hypothesis that deep dives are periodically employed to survey the water column for diurnally descending gelatinous prey. The depth distribution for leatherback turtles in the AFTT Study Area is given in Table 39; within the HSTT Study Area, the leatherback sea turtle is encompassed within the sea turtle guild (Section 2.2.4.2.7) and is represented by the distribution found in Table 46.

Table 39. Percentage of Time at Depth for the Leatherback Turtle¹

Depth Bin (m)	% of Time at Depth
0–10	18.93
10–100	65.26
100–200	14.63
200–300	0.818
300–400	0.119
400–500	0.103
500–600	0.069
600–700	0.023
700–800	0.015
800–900	0.015
900–1,000	0.008
1,000–1,100	0.004
1,100–1,200	0
1,200–1,280	0.008

¹Based on data from Houghton et al. (2008)

While other studies did not include usable depth distributions for leatherback turtles, they did provide additional information to categorize dive behavior. The leatherback is the deepest diving sea turtle, with a recorded maximum depth of 1,280 m (Doyle et al. 2008), though most dives are much shallower, usually less than 250 m (Hays et al. 2004; Sale et al. 2006). Fossette et al. (2007) reported that eighty percent of the leatherback’s time at sea is spent diving, which is in agreement with the roughly 20 percent of time in the surface bin of the representative depth distribution. Dodge et al. (2014) found that over 90% of the time was spent in the top 100 m of the water column, and 25% of time was spent at the surface. Similarly in the Atlantic, Hays et al. (2004) determined that leatherbacks spent 71–94 percent of their diving time at depths from 70 to 110 m. Daytime foraging dives off of the Canadian Atlantic coast during summer ranged between 5.5 and 97 m with a median depth of 21.5 m (Heaslip et al. 2012). Leatherbacks dive deeper and longer in the lower latitudes versus the higher (Dodge et al. 2014; James et al. 2005a; James et al. 2005b), where they are known to dive to waters with temperatures just above freezing (James et al. 2006a; Jonsen et al. 2007). James et al. (2006b) noted that dives in higher latitudes are punctuated by longer surface intervals and much more time at the surface; individuals spend up to 50 percent of their time at or near the surface in northern foraging areas, perhaps in part to thermoregulate (i.e., bask). While transiting, leatherbacks make longer and

deeper dives (Jonsen et al. 2007). During inter-nesting periods, tag data has revealed that dives are likely constrained by bathymetry adjacent to nesting sites (Hays et al. 2006; Myers and Hays 2006).

2.2.4.2 Family Cheloniidae

2.2.4.2.1 *Chelonia mydas*, Green Sea Turtle

The green turtle has a global distribution, occurring throughout tropical and, to a lesser extent, subtropical waters (Marine Turtle Specialist Group 2004) and is generally distributed between 30° N and 30° S. Green turtles are highly migratory and undertake complex movements and migrations through geographically disparate habitats, with the longest migrations occurring between foraging habitats and nesting beaches. Hatchlings swim to offshore areas where they float passively in major current systems. Juveniles leave the pelagic habitat and recruit to protected lagoons and open coastal areas that are rich in seagrass or marine algae (Bresette et al. 2006), the main prey items of green turtles. Green turtles will spend the majority of their lives in these coastal areas (Bjorndal and Bolten 1988; National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 1991). A small number of green turtles appear to remain in open ocean habitats for extended periods, perhaps never recruiting to coastal foraging sites (National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 2007a; Pelletier et al. 2003). The optimal developmental habitats for late juveniles and foraging habitats for adults are warm, shallow waters (3–5 m in bottom depth) with abundant submerged aquatic vegetation and in close proximity to nearshore reefs or rocky areas (Ernst et al. 1994).

In order to build a representative depth distribution for green sea turtles, data from Figures 2 and 4 from Godley et al. (2002) and a maximum dive depth from Rice and Balazs (2008) were used. Technical limitations of the tag used in the Godley et al. (2002) study did not allow for the precise dive depth to be recorded at depths greater than 45 m. The maximum dive depth in other literature for the green turtle is 138 m, recorded from an adult female turtle migrating between nesting grounds on French Frigate Shoals in the Northwest Hawaiian Islands and foraging grounds off Laniakea, Oahu, HI (Rice and Balazs 2008). This value was used to represent the maximum depth for the dive distribution profile. The percentage of time that green turtles spent in each behavioral state (e.g., traveling vs. foraging) was not reported in the literature used for the development of the depth distribution. As a result, the time spent in each behavioral state was assumed to be equal and was averaged for a composite dive depth distribution. The depth distribution for green sea turtles is given in Table 40.

Table 40. Percentage of Time at Depth for the Green Turtle^{1,2}

Depth Bin (m)	% of Time at Depth
0–5	59.23
6–10	16.98
11–15	11.68
16–20	6.78
21–25	2.61
26–30	1.39
31–35	0.73
36–40	0.26
41–45	0.06
45–138	0.28

¹Based on data from Godley et al. (2002) and Rice and Balzacs (2008)

²This depth distribution is also representative of the Kemp’s ridley sea turtle

While other studies did not include usable depth distributions for green turtles, they did provide additional information to categorize dive behavior. Blanco et al. (2012) reported that four turtles near Costa Rica spent 46% of their time at the surface, and while the majority of their time was in the top 10 m, the deepest dive was to 110 m. Seventy percent of the dives of migrating and foraging turtles near Brazil were to less than 30 m depth (Chambault et al. 2015). Hatase (2006) observed that green turtles dive to a maximum of 80 m in areas of the open ocean, where depths are greater than 200 m, while green turtles migrating between the northwestern and main Hawaiian Islands reached a maximum depth greater than 138 m at night (the deepest dives ever recorded for a green turtle), but only 4 m during the day (Rice and Balazs 2008). In their coastal habitat, green turtles typically make dives shallower than 30 m (Cheng et al. 2013; Godley et al. 2002; Hatase et al. 2006; Hays et al. 2000; Hochscheid et al. 1999) and often not exceeding 17.5 m (Ballorain et al. 2013; Hays et al. 2004; Rice and Balazs 2008). Green turtles are known to both forage and rest at depths of 20–50 m (Balazs 1980; Brill et al. 1995).

2.2.4.2.2 *Lepidochelys kempii*, Kemp’s Ridley Sea Turtle

The Kemp’s ridley has one of the smallest ranges of all marine turtle species. The distribution of the Kemp’s ridley population is most concentrated in the Gulf of Mexico, with year round occurrence throughout the Gulf and southern Atlantic coasts of Florida, and seasonal occurrence along the Atlantic coast as far north as Nova Scotia, Canada (Lazell Jr. 1980; Morreale et al. 1992). Habitats frequently utilized by Kemp’s ridley turtles in U.S. waters include warm-temperate to subtropical sounds, bays, estuaries, tidal passes, shipping channels, and beachfront waters where their preferred food, the blue crab, is known to exist (Landry and Costa 1999; Lutcavage and Musick 1985; Seney and Musick 2005).

While past studies have described Kemp’s ridley diving behavior, results were only generalized accounts of diving behavior due to the limited capabilities of the VHF radio, sonic, and satellite telemetry equipment used in these studies. Only a couple of studies exist for which satellite tags were deployed that had the ability to record detailed time-depth information (i.e., the Renaud and Williams (1997) study in Texas and the Sasso and Witzell (2006) study in Florida). Unfortunately, the locations in which they were deployed were all extremely shallow habitats (0.3–6 m), and dive behavior was probably depth-limited, calling into question the reliability of the data. General migration and movement data indicate that Kemp’s ridley turtles generally utilize waters less than 50 m deep as adults, and even shallower waters as juveniles. Byles and Plotkin (1994) noted that 18 adult females stayed in waters less than 50 m in depth during post-nesting movements in the Gulf of Mexico after nesting in Tamulipas, Mexico. This was supported by Renaud (1995) which reported that juvenile Kemp’s ridleys occupy coastal waters of less than 20 m in depth and adults remain in offshore areas

of depths shallower than 50 m. Data compiled in Shaver et al. (2005) for 11 adult male in the Gulf of Mexico (Tamulipas, Mexico) showed that the majority of locations were recorded in waters of 37 m or less. Shaver and Rubio (2007) studied 28 adult female Kemp's ridley turtles and noted similar results after nesting in North Padre and Mustang Island, TX. Lastly, Seney and Landry (2008) found that 6 adult females spent 80 percent of their time in locations with waters less than 10 m deep after nesting in Galveston, TX. The submergence time of Kemp's ridley turtles varies seasonally; dives are longest during the winter (greater than 30 min), and 15 min the remainder of the year (Renaud and Williams 2005). Over a 12 hour period, Kemp's ridley turtles spend as long as 96 percent of their time submerged (Byles 1989; Gitschlag 1996; Renaud and Williams 2005; Sasso and Witzell 2006).

Due to the lack of available data on the diving behavior of the Kemp's ridley sea turtle, it will be represented by a surrogate species: the green sea turtle (Section 2.2.4.2.1). Green sea turtles inhabit similar developmental habitats in the Gulf of Mexico and Florida waters as juvenile and adult Kemp's ridley sea turtles. Furthermore, green turtles are a relatively shallow diving species, similar to the Kemp's ridley. The depth distribution for the Kemp's ridley sea turtle can be found in Table 40.

2.2.4.2.3 *Lepidochelys olivacea*, Olive Ridley Sea Turtle

Olive ridley sea turtles are globally distributed and have a large range in tropical and subtropical regions of the Pacific, Indian, and South Atlantic oceans; they are generally found between 40°N and 40°S. Most olive ridley turtles lead a primarily pelagic existence. The Pacific population migrates throughout the Pacific, from their nesting grounds in Mexico and Central America to the North Pacific (National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 2007c). The olive ridley is the most abundant sea turtle in the open ocean waters of the eastern tropical Pacific Ocean (Pitman 1990). While olive ridleys are primarily a pelagic species and are known to migrate great distances in the Pacific, there are few pelagic records of this species in the northwest Atlantic, and little is known about their migration patterns (National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 2007c). A few reports of olive ridley turtles in the North Atlantic, as far north as the Grand Banks, indicate that the species does traverse the entire Atlantic Ocean, although their occurrence north of the equator is believed rare (National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 2007c).

In order to build a representative depth distribution for olive ridley sea turtles in the AFTT Study Area, data from the text and Figure 3 in Polovina et al. (2003), the same data shown in Figure 5 in Polovina et al. (2004), as well as a maximum dive depth from Swimmer et al. (2006) were used. The depth distribution data given in both Polovina et al. studies are based on data recorded from two satellite tracked olive ridley turtles, which were tagged after being caught in commercial long-line fishing gear southwest of the Hawaiian Islands (Polovina et al. 2004; Polovina et al. 2003). The data found that the tagged olive ridley sea turtles spent about 20 percent of their time in the top meter and that 70 percent of the dives were no deeper than 50 m (Polovina et al. 2004; Polovina et al. 2003). Technical limitations of the tags used in the studies did not allow for the precise dive depth to be recorded at depths greater than 150 m (Polovina et al. 2004; Polovina et al. 2003). The maximum dive depth in other literature for the olive ridley turtle is 288 m, recorded from an adult turtle tagged after being incidentally caught in long-line fishing gear off the Pacific coast of Costa Rica (Swimmer et al. 2006). This value was used to represent the maximum depth for the dive distribution profile. The depth distribution for olive ridley sea turtles is given in Table 41.

Table 41. Percentage of Time at Depth for the Olive Ridley Turtle¹

Depth (m)	% of Time at Depth	Depth (m)	% of Time at Depth
0–1	20.00	81–90	3.00
1–10	5.00	91–100	2.50
11–20	8.50	101–110	1.50
21–30	14.00	111–120	1.00
31–40	13.50	121–130	1.50
41–50	10.00	131–140	0.50
51–60	7.00	141–150	1.00
61–70	5.50	151–288	1.00
71–80	4.50		

¹Based on data from Polovina et al. (2003), Polovina et al. (2004), and Swimmer et al. (2006)

While other studies did not include usable depth distributions for olive ridley sea turtles, they did provide additional information to categorize dive behavior. Olive ridley turtles can dive and feed at considerable depths (80–300 m) (Eckert 1995), although only about 5–10 percent of their time is spent at depths greater than 100 m (Eckert et al. 1986; Polovina et al. 2003). In the eastern tropical Pacific Ocean, at least 25 percent of their total dive time is spent in the permanent thermocline, located between 20–100 m (Parker et al. 2003). Olive ridleys spend considerable time at the surface basking, presumably to speed their metabolism and digestion after a deep dive (Spotila 2004). In the open ocean of the eastern Pacific, olive ridley sea turtles are often seen near flotsam, possibly feeding on associated fish and invertebrates (Pitman 1992). Olive ridleys in the shallow (less than 40 m depth) coastal waters near Oman routinely made use of the entire water column during dives (Rees et al. 2012). The average dive durations for adult females and males are reported to be 54.3 and 28.5 min, respectively [Plotkin, as cited in Lutcavage and Lutz (1997)]. In a separate study by McMahan et al. (2007), olive ridleys exhibited exceptionally long dive durations (greater than two hours), allowing this species to exploit deeper benthic prey.

2.2.4.2.4 AFTT Hardshell Turtle Guild

In the AFTT Study Area, individual species densities were available for the leatherback (Section 2.2.4.1.1), loggerhead (Section 2.2.4.2.6), and Kemp’s ridley (Section 2.2.4.2.2) sea turtles (Navy 2012). For green, hawksbill and olive ridley sea turtles, it was concluded that the number of sightings for these species was not sufficient to allow spatial modeling and density calculation as individual species (U.S. Department of the Navy In Prep.-b). When sightings for these three species of chelonid sea turtles were pooled with the “unidentified” sightings of all hardshell species, a density was calculated for the new group labeled the “AFTT hardshell turtle guild”. Given what is known about sea turtle occurrence along the east coast of the U.S. and in the Gulf of Mexico, it is likely that the majority of the “hardshell turtle” sightings were green turtles. However, as this density number could account for green, hawksbill, or the rarer olive ridley turtle, an average depth distribution (Table 42) was created from the individual depth distributions for these three species. The green sea turtle depth distribution can be found in Section 2.2.4.2.1; the hawksbill sea turtle depth distribution can be found in Section 2.2.4.2.5; and the olive ridley sea turtle depth distribution can be found in Section 2.2.4.2.3. Because each species of sea turtle had a different depth range (green sea turtles diving to a maximum of 138 m, hawksbill sea turtles diving to a maximum of 91 m, and olive ridley sea turtles diving to a much deeper depth of 288 m) the bottom depth bin for hardshell turtles is very large. Each bin over 40 m for each individual species was summed, then averaged with the other species. With data combined in this way, the representative depth distribution in Table 42 represents the AFTT hardshell turtle guild in the family Cheloniidae.

Table 42. Percent of Time at Depth for Hardshell Turtles in the AFTT Study Area¹

Depth Bin (m)	% of Time at Depth
0–10	59.59
11–20	12.82
21–30	7.56
31–40	6.03
41–288	14.01

¹Based on depth distributions for green (Table 40), hawksbill (Table 43), and olive ridley sea turtles (Table 41)

2.2.4.2.5 *Eretmochelys imbricata*, Hawksbill Sea Turtle

The hawksbill is the most tropical of the world’s sea turtles, occurring in all oceans but rarely above or below 30° N and 30° S (Witzell 1983). Hawksbill nesting occurs in at least 70 countries, although much of it now at only low densities. Juveniles and adults share the same foraging areas, including tropical, nearshore waters associated with coral reefs, hard bottoms, or estuaries with mangroves (Musick and Limpus 1997). Coral reefs are optimal hawksbill habitat for juveniles, subadults, and adults (National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 1998). Hawksbills are also found around rocky outcrops and high-energy shoals—optimum sites for sponge growth, their preferred prey item—as well as mangrove-fringed bays and estuaries where coral reefs are absent (National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 2007b). While hawksbills are known to occasionally migrate large distances, possibly in the open ocean, this is the most coastal of all marine turtles. There is very little available information on hawksbills in the pelagic environments of the Atlantic and Pacific Oceans.

In order to build a representative depth distribution for hawksbill sea turtles, data from Figure 6 in Van Dam and Diez (1996) and Figure 7 in Blumenthal et al. (2009) were used. Van Dam and Diez (1996) collected data from four immature hawksbills on the foraging and resting dive behavior for resident juvenile hawksbill turtles of Mona Island, Puerto Rico. Van Dam and Diez (1996) reported foraging dives at the study site in Puerto Rico ranging from 19 to 26 min at depths of 8–10 m. Foraging dives of immature hawksbills are shorter in length, ranging from 8.6 to 14 min in duration (van Dam and Diez 1996), with a mean and maximum depth of 5 m and 20 m, respectively (Blumenthal et al. 2009; Lutcavage and Lutz 1997; van Dam and Diez 1996). A maximum dive duration of 73.5 min has been reported for a female hawksbill in the U.S. Virgin Islands (van Dam and Diez 1996). The data recorded in Blumenthal et al. (2009) represent an average of data from eighteen immature hawksbills captured in Blood Bay, Cayman Islands in 2005. Blumenthal et al. (2009) collected time and depth data that recorded for 8 days for each turtle, but the individuals’ general movements were tracked for an average of 37 ± 69 days (range 11–316 days). Both datasets used to derive the depth distribution for hawksbill turtles were from juvenile turtles. Ideally, data from adult specimens are preferred since dive capacity and habitat use are influenced by body size (Schreer and Kovacs 1997). However, individuals of the same species in the same habitat can vary in body length by a factor of four (Diez and van Dam 2002; McGowan et al. 2008) and in body weight by a factor of 20 (Storch et al. 2005). To date, there have been no studies where a large number of time depth recorders have been deployed on turtles across a wide range of body sizes, enabling investigation of scaling in dive capacity and habitat use (Blumenthal et al. 2009). As a result, the data obtained from studies with juvenile turtles represents the best available data and should be interpreted as generally representative across the entire hawksbill population. The depth distribution for hawksbill sea turtles is given in Table 43.

Table 43. Percentage of Time at Depth for the Hawksbill Turtle^{1,2}

Depth Bin (m)	% of Time at Depth
0–2	11.31
3–10	66.25
11–20	11.49
21–30	4.68
31–40	3.59
41–50	2.04
51–91	0.65

¹Based on data from Van Dam and Diez (1996) and Blumenthal et al. (2009)

²Used to create depth distributions for the AFTT hardshell turtle guild (Table 42) and the HSTT sea turtle guild (Table 46)

While other studies did not include usable depth distributions for hawksbill turtles, they did provide additional information to categorize dive behavior. In general, studies have found that hawksbills may have one of the longest routine dive times of all sea turtles. Starbird et al.(1999) reported that interesting females at Buck Island, U.S. Virgin Islands, averaged 56.1 min dives, longer than those reported in Puerto Rico by Van Dam and Diez (1996). Turtles in Eastern Tropical Pacific spent 89% of their time in waters less than 10 m deep, with the majority of time in the 5-10 m depth bin, and dove to a maximum depth of 40 m (Gaos et al. 2012). Changes in water temperature have an effect on the behavioral ecology of hawksbill turtles, with an increase in nocturnal dive duration with decreasing water temperatures during the winter (Storch et al. 2005).

2.2.4.2.6 *Caretta caretta*, Loggerhead Sea Turtle

Loggerhead turtles are widely distributed in subtropical and temperate waters (Dodd Jr. 1988). The loggerhead turtle occurs worldwide in habitats ranging from coastal estuaries to waters far beyond the continental shelf (Dodd Jr. 1988). The species may be found hundreds of miles out to sea, as well as in inshore areas such as bays, lagoons, salt marshes, creeks, ship channels, and the mouths of large rivers. Results from tagging data of juvenile loggerheads in both the eastern and western North Atlantic suggest that the location of currents and associated frontal eddies are important to the foraging ecology of the pelagic stage of this species (McClellan et al. 2007). Tagging data revealed that migratory routes may be coastal or may involve crossing deep ocean waters; an oceanic route may be taken even when a coastal route is an option (Schroeder et al. 2003). Their large heads support powerful jaws and enable them to feed on hard-shelled prey, such as whelks and conch (Erhart et al. 2003).

In order to build a representative depth distribution for loggerhead turtles in the HSTT Study Area, data from Figure 2c in Howell et al. (2010) and the text of Sakamoto et al. (1990) were used. The data recorded in Howell et al. (2010) are an average of data from fourteen adult loggerheads captured incidentally by long-line vessels in the central North Pacific Ocean during 2002–2004. The turtles were tracked for a period ranging from 51 to 578 days. Technical limitations of the tags used in the Howell et al. (2010 study did not allow for the precise dive depth to be recorded at depths greater than 150 m. The maximum dive depth in other literature for the loggerhead turtle is 233 m, recorded from a female turtle tagged off the coast of Japan (Sakamoto et al. 1990). This value was used to represent the maximum depth for the representative depth distribution below. The depth distribution for loggerhead sea turtles in the HSTT Study Area is given in Table 44.

Table 44. Percent of Time at Depth for the Loggerhead Turtle in the HSTT study area¹

Depth (m)	% of Time at Depth	Depth (m)	% of Time at Depth
0–1	19.25	31–40	0.25
2 –5	43.75	41–50	0.25
6–10	13.00	51–60	0.25
11–15	9.00	61–80	0.25
16–20	9.00	81–100	0.25
21–25	3.00	101–150	0.25
26–30	1.25	150 – 233	0.25

¹Based on Howell et al. (2010) and Sakamoto et al. (1990)

In order to build a representative depth distribution for loggerhead sea turtles in the AFTT Study Area, data from Figures 5.12 through 5.16 in Arendt et al. (2009) were used. The data recorded by Arendt et al. (2009) are an average of data from five male loggerhead turtles satellite tagged after their capture during trawl surveys in Cape Canaveral, FL. The turtles were tracked for an average of 144 days with a range of 7 to 366 days of recordings. Arendt et al. (2009) found that diving behavior of adult male loggerheads in water depths greater than 25 m was characterized by frequent, short (less than 15 min) duration dives and that the majority of time was spent in the upper 5 m of the water column. The depth distribution for loggerhead sea turtles in the AFTT Study Area is given in Table 45.

Table 45. Percentage of Time at Depth for the Loggerhead Turtle in the AFTT Study Area¹

Depth Bin (m)	% of Time at Depth
surface	32.74
0–5	21.67
6–10	6.27
11–15	12.68
16–20	3.66
21–25	5.40
26–50	16.97
51–75	0.28
76–100	0.17
101–127	0.17

¹Based on data from Arendt et al. (2009)

While other studies did not include usable depth distributions for loggerhead turtles, they did provide additional information to categorize dive behavior. Most studies found that, on average, loggerhead turtles spend over 90 percent of their time underwater (Byles 1988; Narazaki et al. 2006; Renaud and Carpenter 1994) and remain at depths shallower than 100 m (Hawkes et al. 2006; Houghton et al. 2002; McClellan et al. 2007; Narazaki et al. 2006; Polovina et al. 2003). Routine dive depths are typically shallower than 30 m (Hochscheid et al. 2010; Houghton et al. 2002), although dives of up to 233 m were recorded for a post-nesting female loggerhead off Japan (Sakamoto et al. 1990), and time-depth recorder-tagged turtles near Brazil recorded occasional dives in the 200-300 m depth bin (Barcelo et al. 2013). Dives can last anywhere from 4 to 120 minutes (Bentivegna et al. 2003; Byles 1988; Dodd and Byles 2003; Renaud and Carpenter 1994; Sakamoto et al. 1990), though approximately 80 percent have a duration of about 2 minutes (Howell et al. 2010).

2.2.4.2.7 HSTT Sea Turtle Guild

This guild includes all sea turtles in Hawaiian waters including green, hawksbill, olive ridley, loggerhead and leatherback sea turtles. In the Hawaii Range Complex (HRC) region of the HSTT Study Area, three species of sea turtles are considered native: green, hawksbill, and leatherback turtles. Two other species, the loggerhead and olive ridley sea turtles, are sometimes observed in Hawaiian waters. Because it is difficult to distinguish between species during surveys, the majority of sighting records were those of “unidentified” sea turtles. The number of species-specific sighting records for individuals were not sufficient to allow for spatial modeling and density calculation. While some hawksbill and green sea turtles were observed, loggerhead, leatherback, and olive ridley were not positively identified during the collection of Navy monitoring data (Hanser et al. 2012). Although not identified during the surveys, loggerhead, leatherback, and olive ridley sea turtles do occur in Hawaiian waters and could still be among the unidentified turtle sightings. Thus, within the HRC region of the HSTT Study Area, the sea turtle guild represents hawksbill, olive ridley, green, loggerhead, and leatherback sea turtles. By considering all of these species sightings together, a more complete result can be provided for sea turtles as a guild.

The combining of sea turtle species in density calculation leads to some generalizations when a depth distribution is created for them. While some species spend the majority of time in surface waters (e.g., green, hawksbill, loggerhead), others spend more time diving below the upper 10 m (e.g., leatherback, olive ridley). Because the leatherback sea turtle is such a deep diver, this representative depth distribution for the HSTT sea turtle guild includes their maximum dive depth of 1,280 m. However, most of the other turtle species spend the majority of their time in the upper 50 m of the water column. In order to separate each individual species depth distribution from larger bins into 10 m bins, it was assumed that, within the larger bin, equal amounts of time were spent in each 10 m increment of that bin. For example, leatherbacks spent 65.26 percent of their time in an 11–100 m bin, according to Houghton et al. (2008). For the purpose of averaging this together with the other species, we have separated the 11–100 m bin into nine 10 m bins, with the leatherback spending 7.25 percent of their time in each 10 m section of that bin. In this manner, each species depth distribution was separated into 10 m bins (for depths up to 150 m, after which the bins increased in size) then averaged together and given equal weight. With data combined in this way, the representative depth distribution in Table 46 represents the HSTT sea turtle guild.

Table 46. Percentage of Time at Depth for the HSTT Sea Turtle Guild¹

Depth (m)	% of Time at Depth	Depth (m)	% of Time at Depth
0–10	54.74	101–110	0.83
11–20	12.74	111–120	0.73
21–30	6.83	121–130	0.83
31–40	5.12	131–140	0.63
41–50	3.92	141–150	0.74
51–60	2.94	151–300	0.74
61–70	2.61	301–400	0.02
71–80	2.41	401–500	0.02
81–90	2.14	501–1280	0.01
91–100	2.01		

¹Based on depth distributions of the green (Table 40), hawksbill (Table 43), leatherback (Table 39), loggerhead (Table 44), and olive ridley sea turtles (Table 41)

2.3 Conclusions

The recommended static depth distributions are provided for 59 marine animal species occurring within the AFTT and HSTT Study Areas. These distributions, especially those that rely on surrogates, should be updated periodically as new data become available. Also, for most species, only a single depth distribution is presented; ideally, each species should have multiple distributions available, depending on the behavior and age/sex class of the animals being modeled, as well as the geographic location and season in which the simulation occurs. More detailed depth distribution data will permit improved realism for the scenarios being modeled.

3 MARINE MAMMAL GROUP SIZE INFORMATION

3.1 Introduction

Many marine mammals are known to travel and feed in groups. The Navy Acoustic Effects Model (NAEMO) accounts for this behavior by incorporating species-specific group sizes into the modeled animal distributions, and accounting for statistical uncertainty around the group size estimate. Methods for determining group size varied between the Atlantic Fleet Training and Testing (AFTT) and Hawaii-Southern California Training and Testing (HSTT) Study Areas, based on data availability and the recommendations of the research groups that provided density information.

Within the AFTT Study Area, group size data were collected through a comprehensive and systematic review of available visual survey data and relevant literature. Journal articles, books, technical reports, cruise reports, funding agency reports, theses, dissertations, and raw data from individual researchers, theses, and dissertations were assessed for this report. Odontocetes are more likely to occur in large groups at sea than other types of marine mammals (Acevedo-Gutierrez 2002), and are the only group for which group size data were gathered within the AFTT Study Area. Within the HSTT Study Area, visual survey data were primarily used in addition to literature, and therefore group size for some mysticetes were also included, but pinniped, sirenians, and sea turtles were not included for either Study Area.

Group size data for AFTT and HSTT includes mean group size, standard deviation, and a range (the minimum and maximum number of animals reported). If a range of means was given in any particular study, the maximum value was used to represent the mean as a conservative estimate of animals present. Minimum and maximum group sizes also were determined for each species. The standard deviations are incorporated in NAEMO by randomly and repetitively selecting a value from the poisson or lognormal distribution defined by the mean group size and standard deviation provided. For HSTT survey data, the Southwest Fisheries Science Center specified which species' group size followed a poisson distribution and which followed a lognormal.

3.2 Atlantic Fleet Training and Testing Study Area Group Sizes

Data from all relevant sources were pooled based on individual species occurring within the AFTT Study Area (Table 47). In some instances where data were lacking, data from multiple areas were combined or the geographically closest data were applied. The short- and long-beaked common dolphins (*Delphinus delphis* and *D. capensis*, respectively) only were differentiated into two specific species within the last decade. Therefore, data for both species of common dolphin were pooled, and herein both species are referred to as common dolphins (*Delphinus* spp.). Additionally, pygmy and dwarf sperm whales (*Kogia breviceps* and *K. simus*, respectively) often are indistinguishable at sea. Subsequently data for these species were combined, and the species herein are referred to as *Kogia* spp.

Table 47. Mean Group Size, Standard Deviation, and Ranges for Odontocetes in the Atlantic Fleet Training and Testing Study Area

Common and Scientific Names	Mean Group Size	SD	Range	References
Sperm whale <i>Physeter macrocephalus</i>	4.47	5.33	1–32	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; Gero et al. 2009; Griffin 1999; Hooker 1998; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b,

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Common and Scientific Names	Mean Group Size	SD	Range	References
				1998c, 1999b, 2002)
Dwarf and pygmy sperm whales <i>Kogia</i> spp.	1.52	0.82	1–4	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; Griffin 1999; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
Atlantic spotted dolphin <i>Stenella frontalis</i>	28.44	10.42	1–121	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1998c, 1999b, 2002)
Atlantic white-sided dolphin <i>Lagenorhynchus acutus</i>	15.86	17.06	1–2500	(Gowans and Whitehead 1995; Hooker et al. 1999; Kingsley and Reeves 1998; National Marine Fisheries Service (NMFS) 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1999a; Simard et al. 2006; Weinrich et al. 2001)
Bottlenose dolphin <i>Tursiops truncatus</i>	14.22	7.59	1–350	(Beck and Rice 2003; Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; Gowans and Whitehead 1995; Griffin 1999; Hooker et al. 1999; Kenney 1990; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
Clymene dolphin <i>Stenella clymene</i>	80.00	75.54	2–165	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; National Marine Fisheries Service (NMFS) 1992b, 1998b, 1999b, 2002)
Common dolphins <i>Delphinus</i> spp.	27.87	29.13	1–700	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; Gowans and Whitehead 1995; Griffin 1999; Hooker et al. 1999; Kenney 1990; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
False killer whale <i>Pseudorca crassidens</i>	5.54	1.71	1–23	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
Fraser's dolphin <i>Lagenodelphis hosei</i>	130.04	61.19	7–1000	(Anderson 2005; Ballance et al. 2001; Clark et al. 2012; De Vos et al. 2012; Dulau-Drouot et al. 2008; Findlay et al. 1992; Kiszka et al. 2007; Weir et al. 2008)
Killer whale <i>Orcinus orca</i>	2.50	0.71	2–3	(National Marine Fisheries Service (NMFS) 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a)
Long-finned pilot whale <i>Globicephala melas</i>	10.21	1.10	1–316	(Anderson and Siegismund 1994; Bloch et al. 2003; de Stephanis et al. 2008; Gowans and

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Common and Scientific Names	Mean Group Size	SD	Range	References
				Whitehead 1995; Hooker et al. 1999; Kingsley and Reeves 1998)
Melon-headed whale <i>Peponocephala electra</i>	23.26	33.85	2–400	(Gero and Whitehead 2006; Maze-Foley and Mullin 2006; Mullin et al. 2004; Watkins et al. 1997)
Pantropical spotted dolphin <i>Stenella attenuata</i>	24.55	23.41	4–145	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
Pygmy killer whale <i>Feresa attenuata</i>	9.17	5.42	3–13	(Clarke and Norman 2005; National Marine Fisheries Service (NMFS) 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a)
Risso's dolphin <i>Grampus griseus</i>	8.47	1.13	1–85	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; Griffin 1999; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
Rough-toothed dolphin <i>Steno bredanensis</i>	5.50	3.54	2–20	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
Short-finned pilot whale <i>Globicephala macrorhynchus</i>	15.37	4.78	1–135	(Caldwell et al. 1971; Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; National Marine Fisheries Service (NMFS) 1992b, 1998b, 1999b, 2002; Ottensmeyer and Whitehead 2003; Payne and Heinemann 1993)
Spinner dolphin <i>Stenella longirostris</i>	27.60	25.84	1–225	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
Striped dolphin <i>Stenella coeruleoalba</i>	45.59	37.50	1–400	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; Gowans and Whitehead 1995; Griffin 1999; Hooker et al. 1999; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
White-beaked dolphin <i>Lagenorhynchus albirostris</i>	8.72	0.35	2–20	(Kingsley and Reeves 1998; Simard and Gowans 2008; Simard et al. 2006)
Harbor porpoise <i>Phocoena phocoena</i>	2.46	1.30	1–25	(Johnston et al. 2005; Kingsley and Reeves 1998; National Marine Fisheries Service (NMFS) 1992a, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1997b, 1998a, 1998c; Simard et al. 2006)

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Common and Scientific Names	Mean Group Size	SD	Range	References
Blainville's beaked whale <i>Mesoplodon densirostris</i>	3.31	1.06	1–10	(Carrillo 2003; Johnson et al. 2004; Ritter 2001; Ritter and Brederlau 1999)
Cuvier's beaked whale <i>Ziphius cavirostris</i>	2.77	0.64	1–6	(Azzellino et al. 2008; Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; National Marine Fisheries Service (NMFS) 1992b, 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a, 1998b, 1999b, 2002)
Gervais' beaked whale <i>Mesoplodon europaeus</i>	3.50	0.05	1–10	(Carrillo 2003; Palka 2012)
Northern bottlenose whale <i>Hyperoodon ampullatus</i>	3.82	1.90	1–14	(Department of the Navy (DoN) 1995, 1998, 1999, 2000, 2001, 2002; Gowans et al. 2001; National Marine Fisheries Service (NMFS) 1992b, 1998b, 1999b, 2002; Reeves et al. 1993; Whitehead and Wimmer 2005)
Sowerby's beaked whale <i>Mesoplodon bidens</i>	2.17	1.04	1–3	(Hooker et al. 1999; National Marine Fisheries Service (NMFS) 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1997b, 1998c; New et al. 2013; Palka 2012)
True's beaked whale <i>Mesoplodon mirus</i>	1.75	0.96	1–3	(Brereton et al. 2004; Kiszka et al. 2007; National Marine Fisheries Service (NMFS) 1995a, 1995b, 1995c, 1995d, 1995e, 1997a, 1998a; New et al. 2013; Tove 1995)

3.3 Hawaii-Southern California Training and Testing Study Area Group Sizes

In order to create a set of group size data for marine mammals in HSTT, survey information was used. For the majority of species, group size estimates were derived from the same datasets used to develop habitat-based density models for the Central Pacific and California Current Ecosystem (Becker et al. 2016; Forney et al. 2015). For island-associated stocks in Hawaii, group size estimates were taken from peer reviewed literature and personal communications with marine mammal experts. Group Size Information

Data from all relevant sources were pooled based on individual species occurring at the following two components of the HSTT Study Area: the Southern California portion of the HSTT Study Area and Hawaii Range Complex (HRC). The average of all mean group sizes reported by the survey data for each species were determined, as well as standard deviations associated with these means. Minimum and maximum group sizes also were determined for each species at each location. In some instances where data were lacking, data from multiple areas were combined or the geographically closest data were applied. For example, pygmy and dwarf sperm whales (*Kogia breviceps* and *K. simus*, respectively) often are indistinguishable at sea, so data for these species were combined, and the species herein are referred to as *Kogia* spp.

Group size information, including means, SDs, and ranges is presented for the Southern California portion of the HSTT Study Area (Table 48) and HRC Study Area (Table 49).

Table 48. Mean Group Size, Standard Deviation, and Range for Cetaceans in the Southern California Portion of the HSTT Study Area

Common and Scientific Name	Mean Group Size	SD	Range	References
Minke whale <i>Balaenoptera acutorostrata</i>	1.14	0.53	1–2	(Becker et al. 2016; Forney et al. 2015)
Sei whale <i>Balaenoptera borealis</i>	1.78	1	1–5	(Becker et al. 2016; Forney et al. 2015)
Bryde's whale <i>Balaenoptera edeni</i>	2.00	0	2–2	(Becker et al. 2016; Forney et al. 2015)
Blue whale <i>Balaenoptera musculus</i>	1.68	0.90	1–5	(Becker et al. 2016; Forney et al. 2015)
Fin whale <i>Balaenoptera physalus</i>	2.23	2.99	1–29.5	(Becker et al. 2016; Forney et al. 2015)
Gray whale <i>Eschrichtius robustus</i>	2.07	1.85	1–4.2	(Becker et al. 2016; Forney et al. 2015)
Humpback whale <i>Megaptera novaeangliae</i>	1.70	0.65	1–3	(Becker et al. 2016; Forney et al. 2015)
<i>Kogia</i> spp.	1.37	0.46	1–3	(Becker et al. 2016; Forney et al. 2015)
Sperm whale <i>Physeter macrocephalus</i>	11.59	20.95	1–109.2	(Becker et al. 2016; Forney et al. 2015)
Long-beaked common dolphin <i>Delphinus capensis</i>	254.57	316.38	1–2,150	(Becker et al. 2016; Forney et al. 2015)

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Common and Scientific Name	Mean Group Size	SD	Range	References
Short-beaked common dolphin <i>Delphinus delphis</i>	161.62	236.83	1.5–1,910	(Becker et al. 2016; Forney et al. 2015)
Pygmy killer whale <i>Feresa attenuata</i>	16.30	10.41	2–35.3	(Becker et al. 2016; Forney et al. 2015)
Short-finned pilot whale <i>Globicephala macrorhynchus</i>	37.65	16.22	26.4–61.7	(Becker et al. 2016; Forney et al. 2015)
Risso's dolphin <i>Grampus griseus</i>	18.40	21.24	1–148.5	(Becker et al. 2016; Forney et al. 2015)
Pacific white-sided dolphin <i>Lagenorhynchus obliquidens</i>	25.85	32.43	1–194.4	(Becker et al. 2016; Forney et al. 2015)
Northern right whale dolphin <i>Lissodelphis borealis</i>	13.41	12.24	2–59.9	(Becker et al. 2016; Forney et al. 2015)
Killer whale <i>Orcinus orca</i>	9.00	13.85	1–47.7	(Becker et al. 2016; Forney et al. 2015)
Bottlenose dolphin, CA/OR/WA <i>Tursiops truncatus</i>	15.88	17.16	1–90	(Becker et al. 2016; Forney et al. 2015)
Bottlenose dolphin, CA coastal stock <i>Tursiops truncatus</i>	9.10	9.80	1–50	(Becker et al. 2016; Forney et al. 2015)
Dall's porpoise <i>Phocoenoides dalli</i>	3.97	2.17	1–8.8	(Becker et al. 2016; Forney et al. 2015)
Mesoplodon beaked whale guild <i>Mesoplodon spp.</i>	2.40	1.02	1–4.5	(Becker et al. 2016; Forney et al. 2015)
Baird's beaked whale <i>Berardius bairdii</i>	8.08	4.35	5–14.5	(Becker et al. 2016; Forney et al. 2015)
Cuvier's beaked whale <i>Ziphius cavirostris</i>	2.04	1.26	1–7	(Becker et al. 2016; Forney et al. 2015)
Striped dolphin <i>Stenalla coeruleoalba</i>	55.97	64.48	1.2–348.8	(Becker et al. 2016; Forney et al. 2015)

Table 49. Mean Group Size, Standard Deviation, and Range for Cetaceans in the Hawaiian Region Study Area

Common and Scientific Name [Stock]	Group Size	SD	Range	References
Minke whale <i>Balaenoptera acutorostrata</i>	1.00	0	1–2	(Becker et al. 2016; Forney et al. 2015)
Sei whale <i>Balaenoptera borealis</i>	2.65	1.18	1–4.5	(Becker et al. 2016; Forney et al. 2015)
Bryde's whale <i>Balaenoptera edeni</i>	1.86	1.67	1–10.2	(Becker et al. 2016; Forney et al. 2015)
Blue whale <i>Balaenoptera musculus</i>	1.43	0.72	1–2.5	(Becker et al. 2016; Forney et al. 2015)
Fin whale <i>Balaenoptera physalus</i>	2.13	0.78	1–3.2	(Becker et al. 2016; Forney et al. 2015)
Humpback whale <i>Megaptera novaeangliae</i>	1.82	1.43	1–8	(Becker et al. 2016; Forney et al. 2015)
Pygmy sperm whale <i>Kogia breviceps</i>	1.00	0	1–3	(Becker et al. 2016; Forney et al. 2015)
Dwarf sperm whale <i>Kogia sima</i>	1.49	0.71	1–3	(Becker et al. 2016; Forney et al. 2015)
Sperm whale <i>Physeter macrocephalus</i>	8.79	9.30	1–53.8	(Becker et al. 2016; Forney et al. 2015)
Pygmy killer whale <i>Feresa attenuata</i>	16.30	10.41	2–35.3	(Becker et al. 2016; Forney et al. 2015)
Short-finned pilot whale <i>Globicephala macrorhynchus</i>	24.28	19.07	1–131.4	(Becker et al. 2016; Forney et al. 2015)
Risso's dolphin <i>Grampus griseus</i>	17.23	19.94	1–99.8	(Becker et al. 2016; Forney et al. 2015)
Fraser's dolphin <i>Lagenodelphis hosei</i>	302.98	352.88	26.3–1535	(Becker et al. 2016; Forney et al. 2015)
Killer whale <i>Orcinus orca</i>	5.71	3.88	1–17	(Becker et al. 2016; Forney et al. 2015)
False killer whale <i>Pseudorca crassidens</i> [Hawaiian Pelagic and Northwestern Hawaiian Islands]	11.18	12.13	1–52	Personal communications with Baird (Baird 2015)
False killer whale <i>Pseudorca crassidens</i> [Main Hawaiian Islands Insular]	17.88	10.26	1–41	(Becker et al. 2016; Forney et al. 2015)
Striped dolphin <i>Stenella coeruleoalba</i>	51.33	38.10	1–186.7	(Becker et al. 2016; Forney et al. 2015)
Bottlenose dolphin <i>Tursiops truncatus</i> [Oahu]	9.93	10.83	1–40	Personal communications with Baird (Baird 2015)
Bottlenose dolphin	5.71	4.56	1–18	Personal communications

Marine Species Dive Distribution and Group Size Parameters Technical Report
 May 2017

Common and Scientific Name [Stock]	Group Size	SD	Range	References
<i>Tursiops truncatus</i> [4-Islands Region]				with Baird (Baird 2015)
Bottlenose dolphin <i>Tursiops truncatus</i> [Kauai and Niihau]	10.23	9.99	1–45	Personal communications with Baird (Baird 2015)
Bottlenose dolphin <i>Tursiops truncatus</i> [Hawaii Islands]	9.59	9.59	1–60	(Becker et al. 2016; Forney et al. 2015)
Bottlenose dolphin <i>Tursiops truncatus</i> [Hawaii Pelagic]	17.55	18.38	1–81	(Becker et al. 2016; Forney et al. 2015)
Rough-toothed dolphin <i>Steno bredanensis</i>	18.99	17.54	1–70	(Becker et al. 2016; Forney et al. 2015)
Longman's beaked whale <i>Indopacetus pacificus</i>	28.77	29.20	3.3–88.5	(Becker et al. 2016; Forney et al. 2015)
Blainville's beaked whale <i>Mesoplodon densirostris</i>	2.51	1.56	1–6	(Becker et al. 2016; Forney et al. 2015)
Cuvier's beaked whale <i>Ziphius cavirostris</i>	2.33	2.12	1–12.8	(Becker et al. 2016; Forney et al. 2015)
Spinner dolphin <i>Stenella longirostris</i> [Hawaii Pelagic]	134.94	219.38	1–2237	(Becker et al. 2016; Forney et al. 2015)
Spinner dolphin <i>Stenella longirostris</i> [Hawaii Island]	49.00	68.93	5–250	(Tyne et al. 2014)
Spinner dolphin <i>Stenella longirostris</i> [Kauai and Niihau]	38.70	35.10	1–185	(Baird et al. 2013)
Spinner dolphin <i>Stenella longirostris</i> [Oahu and 4-Islands Region]	38.70	35.10	1–185	(Baird et al. 2013)
Pantropical spotted dolphin <i>Stenella attenuata</i> [Hawaii Pelagic]	106.95	143.65	1–1400	(Becker et al. 2016; Forney et al. 2015)
Pantropical spotted dolphin <i>Stenella attenuata</i> [Hawaii Island]	65.40	49.60	1–350	(Baird et al. 2013)
Pantropical spotted dolphin <i>Stenella attenuata</i> [Oahu]	65.40	49.60	1–350	(Baird et al. 2013)
Pantropical spotted dolphin <i>Stenella attenuata</i> [4-Islands Region]	65.40	49.60	1–350	(Baird et al. 2013)
Melon-headed whale <i>Peponocephala electra</i> [Hawaiian Islands]	251.8	167.6	1–800	(Baird et al. 2013)
Melon-headed whale <i>Peponocephala electra</i> [Kohala Resident]	251.8	167.6	1–447	(Becker et al. 2016; Forney et al. 2015)

4 REFERENCES

- Acevedo-Gutierrez, A. (2002). Group Behavior. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 537-544). San Diego, CA: Academic Press.
- Acevedo-Gutiérrez, A., Brennan, B., Rodriguez, P., & Thomas, M. (1997). Resightings and Behavior of False Killer Whales (*Pseudorca crassidens*) in Costa Rica. *Marine Mammal Science*, 13(2), 307-314.
- Acevedo-Gutiérrez, A., Croll, D. A., & Tershy, B. R. (2002). High Feeding Costs Limit Dive Time in the Largest Whales. *The Journal of Experimental Biology*, 205, 1747-1753.
- Aguilar, A. (2002). Fin Whale (*Balaenoptera physalus*). In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 435-438). San Diego, CA: Academic Press.
- Aguilar de Soto, N., Johnson, M., Madsen, P. T., Tyack, P. L., Bocconcelli, A., & Fabrizio Borsani, J. (2006). DOES INTENSE SHIP NOISE DISRUPT FORAGING IN DEEP-DIVING CUVIER'S BEAKED WHALES (ZIPHIUS CAVIROSTRIS)? *Marine Mammal Science*, 22(3), 690-699.
- Alonso, M. K., Pedraza, S. N., Schiavini, A. C. M., Goodall, R. N. P., & Crespo, E. A. (1999). Stomach Contents of False Killer Whales (*Pseudorca crassidens*) Stranded on the Coasts of the Strait of Magellan, Tierra del Fuego. *Marine Mammal Science*, 15(3), 712-724.
- Alves-Stanley, C. D., Worthy, G. A. J., & Bonde, R. K. (2010). Feeding Preferences of West Indian Manatees in Florida, Belize, and Puerto Rico as Indicated by Stable Isotope Analysis. *Marine Ecology Progress Series*, 402, 255-267.
- Alves, F., Dinis, A., & Cascão, I. (2010). Bryde's Whale (*Balaenoptera brydei*) Stable Associations and Dive Profiles: New Insights into Foraging Behavior. *Marine Mammal Science*, 26(1), 202-212.
- Amano, M., & Yoshioka, M. (2003). Sperm Whale Diving Behavior Monitored Using a Suction-Cup-Attached TDR Tag. *Marine Ecology Progress Series*, 258, 291-295.
- Anderson, L. W., & Siegismund, H. R. (1994). Genetic Evidence for Migration of Males between Schools of Long-Finned Pilot Whale *Globicephala melas*. *Marine Ecology Progress Series*, 105, 1-7.
- Anderson, R. C. (2005). Observations of Cetaceans in the Maldives, 1990-2002. *The Journal of Cetacean Research and Management*, 7(2), 119-135.
- Anderson, R. C., Clark, R., Madsen, P. T., Johnson, C., Kiszka, J., & Breysse, O. (2006). Observations of Longman's Beaked Whale (*Indopacetus pacificus*) in the Western Indian Ocean. *Aquatic Mammals*, 32(2), 223-231.
- Andrade, A. L. V., Pinedo, M. C., & Barreto, A. S. (2001). Gastrointestinal Parasites and Prey Items from a Mass Stranding of False Killer Whales, *Pseudorca crassidens*, in Rio Grande do Sul, Southern Brazil. *Revista Brasileira de Biologia*, 61(1), 55-61.
- Aoki, K., Amano, M., Mori, K., Kouroggi, A., Kubodera, T., & Miyazaki, N. (2012). Active hunting by deep-diving sperm whales: 3D dive profiles and maneuvers during bursts of speed. *Marine Ecology Progress Series*, 444, 289-301.

- Aoki, K., Amano, M., Yoshioka, M., Mori, K., Tokuda, D., & Miyazaki, N. (2007). Diel Diving Behavior of Sperm Whales off Japan. *Marine Ecology Progress Series*, 349, 277-287.
- Aoki, K., Sakai, M., Miller, P. J. O., Visser, F., & Sato, K. (2013). Body contact and synchronous diving in long-finned pilot whales. *Behavioural Processes*, 99, 12-20. doi: DOI: 10.1016/j.beproc.2013.06.002.
- Archer II, F. I. (2002). Striped Dolphin: *Stenella coeruleoalba*. In: Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1201-1203). San Diego, CA: Academic Press.
- Arendt, M., Byrd, J., Segars, A., Maier, P., Schwenter, J., Burgess, D., . . . Blanvillain, G. (2009). *Examination of Local Movement and Migratory Behavior of Sea Turtles during Spring and Summer Along the Atlantic Coast off the Southeastern United States*. Charleston, South Carolina: South Carolina Department of Natural Resources.
- Arranz, P., de Soto, N. A., Madsen, P. T., Brito, A., Bordes, F., & Johnson, M. P. (2011). Following a foraging fish-finder: diel habitat use of Blainville's beaked whales revealed by echolocation. *PloS one*, 6(12), e28353.
- Aurioles-Gamboa, D., & Camacho-Ríos, F. J. (2007). Diet and Feeding Overlap of Two Otariids, *Zalophus californianus* and *Arctocephalus townsendi*: Implications to Survive Environmental Uncertainty. *Aquatic Mammals*, 33(3), 315-326.
- Azzellino, A., Gaspari, S., Airoidi, S., & Nani, B. (2008). Habitat Use and Preferences of Cetaceans along the Continental Slope and the Adjacent Pelagic Waters in the Western Ligurian Sea. *Deep-Sea Research I*, 55, 296–323.
- Baird, R. W. (1994). *Foraging Behaviour and Ecology of Transient Killer Whales (Orcinus Orca)*. Doctor of Philosophy, Simon Fraser University.
- Baird, R. W. (2009). *A Review of False Killer Whales in Hawaiian Waters: Biology, Status, and Risk Factors*. (Order No. E40475499). Olympia, WA: United States Marine Mammal Commission. p. 41.
- Baird, R. W. (2015, 12 November 2015). *Personal Communication*. Personal communication via to Blackstock, S.,
- Baird, R. W., Borsani, J. F., Hanson, M. B., & Tyack, P. L. (2002). Diving and Night-Time Behavior of Long-Finned Pilot Whales in the Ligurian Sea. *Marine Ecology Progress Series*, 237, 301-305.
- Baird, R. W., & Hanson, M. B. (1998). A Preliminary Analysis of the Diving Behavior of Dall's Porpoise in the Transboundary Waters of British Columbia and Washington. In: Hill, P. S., Jones, B. & DeMaster, D. P. (Eds.), *Marine Mammal Protection Act and Endangered Species Act Implementation Program 1997* (pp. 99-110). Silver Spring, MD: National Marine Fisheries Service.
- Baird, R. W., Hanson, M. B., & Dill, L. M. (2005a). Factors Influencing the Diving Behaviour of Fish-eating Killer Whales: Sex Differences and Diel and Interannual Variation in Diving Rates. *Canadian Journal of Zoology*, 83, 257-267.

- Baird, R. W., Ligon, A. D., & Hooker, S. K. (2000). *Sub-Surface and Night-Time Behavior of Humpback Whales off Maui, Hawaii: A Preliminary Report*. Paia, HI: Hawaii Wildlife Fund. p. 18.
- Baird, R. W., Ligon, A. D., Hooker, S. K., & Gorgone, A. M. (2001). Subsurface and Nighttime Behaviour of Pantropical Spotted Dolphins in Hawai'i. *Canadian Journal of Zoology*, 79(6), 988-996.
- Baird, R. W., Martin, S. W., Webster, D. L., & Southall, B. L. (2014). *Assessment of Modeled Received Sound Pressure Levels and Movements of Satellite-Tagged Odontocetes Exposed to Mid-Frequency Active Sonar at the Pacific Missile Range Facility: February 2011 Through February 2013*. DTIC Document.
- Baird, R. W., McSweeney, D. J., Webster, D. L., Gorgone, A. M., & Ligon, A. D. (2003). *Studies of Odontocete Population Structure in Hawaiian Waters: Results of a Survey Through the Main Hawaiian Islands in May and June 2003*. (Contract Number AB133F-02-CN-0106). Seattle, WA: National Oceanic and Atmospheric Administration, Western Administrative Support Center. p. 24.
- Baird, R. W., Webster, D. L., Mahaffy, S. D., Schorr, G. S., Aschettino, J. M., & Gorgone, A. M. (2013). *Movement and spatial use of Odontocetes in the western main Hawaiian Islands: results of a three-year study off O'ahu and Kaua'i*. Monterey, California, Naval Postgraduate School.
- Baird, R. W., Webster, D. L., McSweeney, D. J., Ligon, A. D., & Schorr, G. S. (2005b). *Diving Behavior and Ecology of Cuvier's (Ziphius cavirostris) and Blainville's Beaked Whales (Mesoplodon densirostris) in Hawai'i*. La Jolla, California: National Marine Fisheries Service (NMFS). p. 24.
- Baird, R. W., Webster, D. L., McSweeney, D. J., Ligon, A. D., Schorr, G. S., & Barlow, J. (2006). Diving Behaviour of Cuvier's (*Ziphius cavirostris*) and Blainville's (*Mesoplodon densirostris*) Beaked Whales in Hawai'i. *Canadian Journal of Zoology*, 84, 1120-1128.
- Baird, R. W., Webster, D. L., Schorr, G. S., McSweeney, D. J., & Barlow, J. (2008). Diel Variation in Beaked Whale Diving Behavior. *Marine Mammal Science*, 24(3), 630-642.
- Bajzak, C. E., Côté, S. D., Hammill, M. O., & Stenson, G. (2009). Intersexual Differences in the Postbreeding Foraging Behaviour of the Northwest Atlantic Hooded Seal. *Marine Ecology Progress Series*, 385, 285-294.
- Baker, J. D. (2007). Post-weaning Migration of Northern Fur Seal *Callorhinus ursinus* Pups from the Pribilof Islands, Alaska. *Marine Ecology Progress Series*, 341, 243-255.
- Baker, J. D., & Donohue, M. J. (2000). Ontogeny of Swimming and Diving in Northern Fur Seal (*Callorhinus ursinus*) Pups. *Canadian Journal of Zoology*, 78, 100-109.
- Balazs, G. H. (1980). *Synopsis of Biological Data on the Green Turtle in the Hawaiian Islands*. (NOAA Technical Memorandum NMFS). Honolulu, HI: National Marine Fisheries Service. p. 141.
- Balcomb, K. C., III. (1989). Baird's Beaked Whale -- *Berardius bairdii* Stejneger, 1883: Arnoux's Beaked Whale -- *Berardius arnuxii* Duvernoy, 1851. In: Ridgway, S. H. & Harrison, R.

- (Eds.), *Handbook of Marine Mammals: Volume 4: River Dolphins and the Larger Toothed Whales* (Vol. 4, pp. 261-288). New York: Academic Press.
- Ballance, L. T., Anderson, R. C., Pitman, R. L., Stafford, K., Shaan, A., Waheed, Z., & Brownell, R. L., Jr. (2001). Cetacean Sightings Around the Republic of the Maldives, April 1998. *Journal of Cetacean Research and Management*, 3(2), 213-218.
- Ballorain, K., Bourjea, J., Ciccione, S., Kato, A., Hanuise, N., Enstipp, M., . . . Georges, J.-Y. (2013). Seasonal diving behaviour and feeding rhythms of green turtles at Mayotte Island. *Marine Ecology Progress Series*, 483, 289-302. doi: 10.3354/meps10301.
- Barlow, J. (2010). *Cetacean Abundance in the California Current Estimated from a 2008 Ship-Based Line-Transsect Survey*. (NMFS-SWFSC-456). U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service.
- Barlow, J., & Forney, K. A. (2007). Abundance and Density of Cetacean in the California Current Ecosystem. *Fishery Bulletin*, 105(4), 509-526.
- Barlow, J., Tyack, P. L., Johnson, M. P., Baird, R. W., Schorr, G. S., Andrews, R. D., & de Soto, N. A. (2013). Trackline and point detection probabilities for acoustic surveys of Cuvier's and Blainville's beaked whales. *The Journal of the Acoustical Society of America*, 134(3), 2486-2496.
- Barrett-Lennard, L. G., Ford, J. K. B., & Heise, K. A. (1996). The Mixed Blessing of Echolocation: Differences in Sonar Use by Fish-Eating and Mammal-Eating Whales. *Animal Behaviour*, 51, 553-565.
- Baugh, T. M., Valade, J. A., & Zoodsma, B. J. (1989). Manatee Use of *Spartina alterniflora* in Cumberland Sound. *Marine Mammal Science*, 5(1), 88-90.
- Baumgartner, M., & Fratantoni, D. M. (2008). Diel periodicity in both sei whale vocalization rates and the vertical migration of their copepod prey observed from ocean gliders. *Limnol Oceanogr*, 53, 2197-2209.
- Baumgartner, M., Lysiak, N. S. J., Schuman, C., Urban-Rich, J., & Wenzel, F. (2011). Diel vertical migration behavior of *Calanus finamarchicus* and its influence on right and sei whale occurrence. *Mar Ecol Prog Ser*, 423, 167-184.
- Baumgartner, M. F. (1997). The Distribution of Risso's Dolphin (*Grampus griseus*) with Respect to the Physiography of the Northern Gulf of Mexico. *Marine Mammal Science*, 13(4), 614-638.
- Baumgartner, M. F., & Mate, B. R. (2003). Summertime Foraging Ecology of North Atlantic Right Whales. *Marine Ecology Progress Series*, 264, 123-135.
- Beatson, E. (2007). The Diet of Pygmy Sperm Whales, *Kogia breviceps*, Stranded in New Zealand: Implications for Conservation. *Reviews in Fish Biology and Fisheries*, 17, 295-303.
- Beck, B. M., & Rice, C. D. (2003). Serum Antibody Levels Against Select Bacterial Pathogens in Atlantic Bottlenose Dolphins, *Tursiops truncatus*, from Beaufort NC USA and Charleston Harbor, Charleston, SC, USA. *Marine Environmental Research*, 55, 161-179.

- Beck, C. A., Bowen, W. D., McMillan, J. I., & Iverson, S. J. (2003). Sex Differences in the Diving Behaviour of a Size-Dimorphic Capital Breeder: The Grey Seal. *Animal Behaviour*, 66(4), 777-789.
- Becker, E. A., Forney, K. A., Fiedler, P. C., Barlow, J., Chivers, S. J., Edwards, C. A., . . . Redfern, J. V. (2016). Moving Towards Dynamic Ocean Management: How Well Do Modeled Ocean Products Predict Species Distributions? *Remote Sensing*, 8(2), 149.
- Belcher, R. L., & Lee, T. E. J. (2002). *Arctocephalus townsendi*. *Mammalian Species*, 700, 1-5.
- Benoit-Bird, K. J., & Au, W. W. L. (2003). Prey Dynamics Affect Foraging by a Pelagic Predator (*Stenella longirostris*) Over a Range of Spatial and Temporal Scales. *Behavioral Ecology and Sociobiology*, 53, 364-373.
- Bentivegna, F., Hochscheid, S., & Minucci, C. (2003). Seasonal Variability in Voluntary Dive Duration of the Mediterranean Loggerhead Turtle, *Caretta caretta*. *Scientia Marina*, 67(3), 371-375.
- Best, P. B. (2001). Distribution and Population Separation of Bryde's Whale *Balaenoptera edeni* off Southern Africa. *Marine Ecology Progress Series*, 220, 277-289.
- Bjorge, A., & Tolley, K. A. (2002). Harbor Porpoise. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 549-551). San Diego, CA: Academic Press.
- Bjørke, H. (2001). Predators of the Squid *Gonatus fabricii* (Lichtenstein) in the Norwegian Sea. *Fisheries Research*, 52, 113-120.
- Bjorndal, K. A., & Bolten, A. B. (1988). Growth Rates of Immature Green Turtles, *Chelonia mydas*, of Feeding Grounds in the Southern Bahamas. *Copeia*, 1988(3), 555-564.
- Black, N. A. (1994). *Behavior and Ecology of Pacific White-sided Dolphins (Lagenorhynchus obliquidens) in Monterey Bay, California*. (Master's thesis), San Francisco State University.
- Blanco, C., Aznar, J., & Raga, J. A. (1995). Cephalopods in the Diet of Striped Dolphin *Stenella coeruleoalba* from the Western Mediterranean during an Epizootic in 1990. *Journal of Zoology: Proceedings of the Zoological Society of London*, 237, 151-158.
- Blanco, G. S., Morreale, S. J., Seminoff, J. A., Paladino, F. V., Piedra, R., & Spotila, J. R. (2012). Movements and diving behavior of internesting green turtles along Pacific Costa Rica. *Integrative Zoology*. doi: 10.1111/j.1749-4877.2012.00298.x.
- Bleakney, J. S. (1965). Reports of Marine Turtles from New England and Eastern Canada. *Canadian Field-Naturalist*, 79, 120-128.
- Blix, A. S., & Folkow, L. P. (1995). Daily Energy Expenditure in Free Living Minke Whales. *Acta Physiologica Scandinavica*, 153(1), 61-66.
- Bloch, D., Heide-Jorgensen, M. P., Stefansson, E., Mikkelsen, B., Ofstad, L. H., Dietz, R., & Andersen, L. W. (2003). Short-term Movements of Long-finned Pilot Whales *Globicephala melas* Around the Faroe Islands. *Wildlife Biology*, 9(1), 47-58.
- Bloodworth, B. E., & Odell, D. K. (2008). *Kogia breviceps* (Cetacea: Kogiidae). *Mammalian Species*, 819, 1-12.

- Blumenthal, J. M., Austin, T. J., Bothwell, J. B., Broderick, A. C., Ebanks-Petrie, G., Olynik, J. R., . . . Godley, B. J. (2009). Diving Behavior and Movements of Juvenile Hawksbill Turtles *Eretmochelys imbricata* on a Caribbean Coral Reef. *Coral Reefs*, 28, 55-65.
- Bowen, W. D., Boness, D. J., & Iverson, S. J. (1999). Diving Behaviour of Lactating Harbour Seals and Their Pups During Maternal Foraging Trips. *Canadian Journal of Zoology*, 77, 978-988.
- Bowen, W. D., Lawson, J. W., & Beck, B. (1993). Seasonal and Geographic Variation in the Species Composition and Size of Prey Consumed by Grey Seals (*Halichoerus grypus*) on the Scotian Shelf. *Canadian Journal of Fisheries and Aquatic Sciences*, 50(8), 1768-1778.
- Brereton, T., Wall, D., Cermeño, P., Vasquez, A., Curtis, D., & Williams, A. (2004). *Cetacean Monitoring in North-West European Waters 2001*. The Atlantic Research Coalition. p. 28.
- Bresette, M., Singewald, D. A., & DeMaye, E. (2006). *Recruitment of Post-Pelagic Green Turtles (Chelonia mydas) to Nearshore Reefs on Florida's East Coast*. Paper presented at the Twenty-sixth Annual Symposium on Sea Turtle Biology and Conservation: Book of Abstracts, Athens, Greece.
- Brill, R. W., Balazs, G. H., Holland, K. N., Chang, R. K. C., Sullivan, S., & George, J. C. (1995). Daily Movements, Habitat Use, and Submergence Intervals of Normal and Tumor-Bearing Juvenile Green Turtles (*Chelonia mydas* L.) Within a Foraging Area in the Hawaiian Islands. *Journal of Experimental Marine Biology and Ecology*, 185, 203-218.
- Brongersma, L. D. (1972). European Atlantic Turtles. In *Zoologische Verhandelingen* (Vol. 121, pp. 42-109). Leiden, Netherlands: E.J. Brill.
- Brown, R. F., & Mate, B. R. (1983). Abundance, Movements, and Feeding Habits of Harbor Seals, *Phoca vitulina*, at Netarts and Tillamook Bays, Oregon. *Fishery Bulletin*, 81, 291-301.
- Brownell Jr., R. L., Ralls, K., Baumann-Pickering, S., & Poole, M. M. (2009). Behavior of Melon-Headed Whales, *Peponocephala electra*, near Oceanic Islands. *Marine Mammal Science*, 25(3), 639-658.
- Brownell, R. L., Jr., Walker, W. A., & Forney, K. A. (1999). Pacific White-sided Dolphin -- *Lagenorhynchus obliquidens* Gill, 1865. In Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals. Volume 6: The Second Book of Dolphins and the Porpoises* (Vol. 6, pp. 57-84). New York: Academic Press.
- Buckland, S. T., Bloch, D., Cattanaach, K. L., Gunnlaugsson, T., Hoydal, K., Lens, S., & Sigurjónsson, J. (1993). Distribution and Abundance of Long-finned Pilot Whales in the North Atlantic, Estimated from NASS-87 and NASS-89 Data. In Donovan, G. P., Lockyer, C. H. & Martin, A. R. (Eds.), *Biology of Northern Hemisphere Pilot Whales* (Vol. 14, pp. 33-49). Cambridge, UK: International Whaling Commission.
- Budylenko, G. A. (1978). On Sei Whale Feeding in the Southern Ocean. *Reports of the International Whaling Commission*, 28, 379-385.
- Burns, J. J. (2002). Harbor Seal and Spotted Seal. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 552-560). San Diego, CA: Academic Press.

- Byles, B. A. (1989). *Satellite Telemetry of Kemp's Ridley Sea Turtle, Lepidochelys kempii, in the Gulf of Mexico*. Paper presented at the Proceedings of the Ninth Annual Workshop on Sea Turtle Conservation and Biology.
- Byles, R. A. (1988). *Behavior and Ecology of Sea Turtles from Chesapeake Bay, Virginia*. (Ph.D. diss.), College of William and Mary, Williamsburg, VA.
- Byles, R. A., & Plotkin, P. T. (1994). *Comparison of the Migratory Behavior of the Congeneric Sea Turtles Lepidochelys olivaceas and L. kempii*. Paper presented at the Thirteenth Annual Symposium on Sea Turtle Biology and Conservation, Jekyll Island, GA.
- Cacchione, D. A., Drake, D. E., Field, M. E., & Tate, G. B. (1987). Sea-Floor Gouges caused by Migrating Gray Whales off Northern California. *Continental Shelf Research*, 7(6), 553-560.
- Calambokidis, J., Schorr, G. S., Steiger, G. H., Francis, J., Bakhtiari, M., Marshall, G., . . . Robertson, K. (2008). Insights into the Underwater Diving, Feeding, and Calling Behavior of Blue Whales from a Suction-Cup-Attached Video-Imaging Tag (Cittercam). *Marine Technology Society Journal*, 41(4), 19-29.
- Caldwell, D. K., Neuhauser, H., Caldwell, M. C., & Coolidge, H. W. (1971). Recent Records of Marine Mammals from the Coasts of Georgia and South Carolina. *Cetology*, 5, 1-12.
- Cañadas, A., & Hammond, P. S. (2008). Abundance and Habitat Preferences of the Short-beaked Common Dolphin *Delphinus delphis* in the Southwestern Mediterranean: Implications for Conservation. *Endangered Species Research*, 4, 309-331.
- Cañadas, A., Sagarminaga, R., & García-Tiscar, S. (2002). Cetacean Distribution Related with Depth and Slope in the Mediterranean Waters off Southern Spain. *Deep-Sea Research I*, 49, 2053-2073.
- Cardinale, M., Casini, M., Arrhenius, F., & Håkansson, N. (2003). Diel spatial distribution and feeding activity of herring (*Clupea harengus*) and sprat (*Sprattus sprattus*) in the Baltic Sea. *Aquatic Living Resources*, 16, 283-292.
- Carrillo, M. (2003, 10 -13 March 2003). *Presence and Distribution of the Ziphiidae Family in the South West Coast of Tenerife. Canary Islands*. Paper presented at the 17th Annual Conference of the European Cetacean Society, Canary Island, Spain.
- Cheng, I. J., Bentivegna, F., & Hochscheid, S. (2013). The behavioural choices of green turtles nesting at two environmentally different islands in Taiwan. *Journal of Experimental Marine Biology and Ecology*, 440, 141-148. doi: 10.1016/j.jembe.2012.12.002.
- Clapham, P. J. (2002). Humpback Whale. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 589-592). San Diego, CA: Academic Press.
- Clark, R. A., Johnson, C. M., Johnson, G., Payne, R., Kerr, I., Anderson, R. C., . . . Madsen, P. T. (2012). Cetacean sightings and acoustic detections in the offshore waters of the Maldives during the northeast monsoon seasons of 2003 and 2004. *J. Cetacean Res. Manage*, 12, 227-234.
- Clarke, J. T., & Norman, S. A. (2005). Results and Evaluation of US Navy Shock Trial Environmental Migration of Marine Mammals and Sea Turtles. *The Journal of Cetacean Research and Management*, 7(1), 43-50.

- Clarke, M. R. (2003). Production and control of sound by the small sperm whales, *Kogia breviceps* and *K. sima* and their implications for other Cetacea. *Journal of the Marine Biological Association of the UK*, *83*, 241-263. doi: 10.1017/S0025315403007045h.
- Clarke, M. R., & Pascoe, P. L. (1985). The Stomach Contents of a Risso's Dolphin (*Grampus griseus*) Stranded at Thurlestone, South Devon. *Journal of Marine Biological Association of the United Kingdom*, *65*, 663-665.
- Clarke, T. A. (1973). Some aspects of the ecology of lanternfishes (Myctophidae) in the Pacific Ocean near Hawai'i. *Fisheries Bulletin*, *71*, 127-138.
- Cooper, C. (1993). Diving Behaviour of Harbour Porpoises in the Lower Bay of Fundy. *Project Summary, Industry Services and Native Fisheries*, *42*, 4.
- Costa, D. P., Kuhn, C., Weise, M., Shaffer, S., & Arnould, J. P. Y. (2004). When Does Physiology Limit the Foraging Behavior of Freely Diving Mammals. *International Congress Series*, *1275*, 259-366.
- Couperus, A. S. (1997). Interactions Between Dutch Midwater Trawl and Atlantic White-sided Dolphins (*Lagenorhynchus acutus*) Southwest of Ireland. *Journal of Northwest Atlantic Fishery Science*, *22*, 209-218.
- Courbis, S. S., & Worthy, G. A. J. (2003). Opportunistic Carnivory by Florida Manatees (*Trichechus manatus latirostris*). *Aquatic Mammals*, *29*(1), 104-107.
- Craddock, J. E., Polloni, P. T., Hayward, B., & Wenzel, F. (2009). Food Habits of Atlantic White-sided Dolphins (*Lagenorhynchus acutus*) off the Coast of New England. *Fishery Bulletin*, *107*(3), 384-394.
- Crocker, D. E., Costa, D. P., Boeuf, B. J. L., Webb, P. M., & Houser, D. S. (2006). Impact of El Nino on the Foraging Behavior of Female Northern Elephant Seals. *Marine Ecology Progress Series*, *309*, 1-10.
- Croll, D. A., Acevedo-Gutierrez, A., Tershy, B. R., & Urban-Ramirez, J. (2001). The Diving Behavior of Blue and Fin Whales: Is Dive Duration Shorter than Expected Based on Oxygen Stores? *Comparative Biochemistry and Physiology*, *129*, 797-809.
- Cruikshank, R. A., & Brown, S. G. (1981). Recent Observations and Some Historical Records of Southern Right-Whale Dolphins *Lissodelphis peronii*. *Fisheries Bulletin*, *15*, 109-121.
- Dalebout, M. L., Ross, G. J. B., Baker, C. S., Anderson, R. C., Best, P. B., Cockcroft, V. G., . . . Pitman, R. L. (2003). Appearance, Distribution and Genetic Distinctiveness of Longman's Beaked Whale, *Indopacetus pacificus*. *Marine Mammal Science*, *19*(3), 421-461.
- Darling, J. D., Keogh, K. E., & Steeves, T. E. (1998). Gray Whale (*Eschrichtius robustus*) Habitat Utilization and Prey Species off Vancouver Island, B.C. *Marine Mammal Science*, *14*(4), 692-720.
- Davis, R. W., & Fargion, G. S. (1996). *Distribution and Abundance of Cetaceans in the North-Central and Western Gulf of Mexico, Final Report. Volume II: Technical Report*. (OCS Study MMS 96-0027). New Orleans, LA: U.S. Dept of the Interior, Minerals Management Service. p. 356.

- Davis, R. W., Fargion, G. S., May, N., Leming, T. D., Baumgartner, M. F., Evans, W. E., . . . Mullin, K. D. (1998). Physical Habitat of Cetaceans Along the Continental Slope in the North-Central and Western Gulf of Mexico. *Marine Mammal Science*, 14(3), 490-507.
- Davis, R. W., Fuiman, L. A., Williams, T. M., & Le Boeuf, B. J. (2001). Three-Dimensional Movements and Swimming Activity of a Northern Elephant Seal. *Comparative Biochemistry and Physiology Part A Molecular & Integrative Physiology*, 129A(4), 759-770.
- Davis, R. W., & Weihs, D. (2007). Locomotion in Diving Elephant Seals: Physical and Physiological Constraints. *Philosophical Transactions of the Royal Society of Britain*, 362, 2141-2150.
- Davis, R. W., Worthy, G. A. J., Würsig, B., Lynn, S. K., & Townsend, F. I. (1996). Diving Behavior and At-Sea Movements of an Atlantic Spotted Dolphin in the Gulf of Mexico. *Marine Mammal Science*, 12(4), 569-581.
- de Stephanis, R., Cornulier, T., Verborgh, P., Sierra, J. S., Gimeno, N. P., & Guinet, C. (2008). Summer Spatial Distribution of Cetaceans in the Strait of Gibraltar in Relation to the Oceanographic Context. *Marine Ecology Progress Series*, 353, 275-288.
- De Vos, A., Clark, R., Johnson, C., Johnson, G., Kerr, I., Payne, R., & MADSEN, P. (2012). Cetacean sightings and acoustic detections in the offshore waters of Sri Lanka: March–June 2003. *J. Cetacean Res. Manage*, 12(2), 185-193.
- DeLong, R. L., Kooyman, G. L., Gilmartin, W. G., & Loughlin, T. R. (1984). Hawaiian Monk Seal Diving Behavior. *Acta Zoologica Fennica*, 172, 129-131.
- DeLong, R. L., & Stewart, B. S. (1991). Diving Patterns of Northern Elephant Seal Bulls. *Marine Mammal Science*, 7(4), 369-384.
- Department of the Navy (DoN). (1995). *Aerial Census Survey Report of Marine Mammals and Sea Turtles within Candidate Test Sites off Norfolk, Virginia and Mayport, Florida. Surveys 1-6*. Prepared for the Southern Division, Naval Facilities Engineering Command by Continental Shelf Associates, Inc.
- Department of the Navy (DoN). (1998). *Aerial Census Survey Report of Marine Mammals and Sea Turtles within the Candidate Test Area off Mayport, Florida: Surveys 1-5: May-September 1997*. Jupiter, FL: Continental Shelf Associates, Inc. p. 47.
- Department of the Navy (DoN). (1999). *Aerial Survey Report for Mayport Testing Area, June and August 1999*. Jupiter, FL: Continental Shelf Associates Inc. p. 20.
- Department of the Navy (DoN). (2000). *Data Report for the 1997-1998 Year 2 Aerial Surveys of Northern Right Whales and other Listed Species in Atlantic Waters from Charleston, South Carolina to Cape Canaveral, Florida*.
- Department of the Navy (DoN). (2001). *Data Report for the 1998-1999 Year 3 Aerial Surveys of Northern Right Whales and other Listed Species in Atlantic Waters from Charleston, South Carolina to Cape Canaveral, Florida*.
- Department of the Navy (DoN). (2002). *Summary Report on Aerial Surveys (1996/97, 1997/98, 1998/99) of Northern Right Whales and Other Listed Species in Atlantic Waters from Charleston, South Carolina to Cape Canaveral, Florida*. Jupiter, FL: Continental Shelf Associates, Inc. p. 78.

- Desportes, G., & Mouritsen, R. (1993). Preliminary Results on the Diet of Long-finned Pilot Whales off the Faroe Islands. In: Donovan, G. P., Lockyer, C. H. & Martin, A. R. (Eds.), *Biology of Northern Hemisphere Pilot Whales* (Vol. 14, pp. 306-324). Cambridge, UK: International Whaling Commission.
- Dietz, R., Teilmann, J., Jorgensen, M.-P. H., & Jensen, M. V. (2002). *Satellite Tracking of Humpback Whales in West Greenland*. (NERI Technical Report No. 411). Roskilde, Denmark: National Environmental Research Institute. p. 40.
- Diez, C. E., & van Dam, R. P. (2002). Habitat Effect on Hawksbill Turtle Growth Rates on Feeding Grounds at Mona and Monito Islands, Puerto Rico. *Marine Ecology Progress Series*, 234, 301-309.
- Dodd, C. K. J., & Byles, R. (2003). Post-Nesting Movements and Behavior of Loggerhead Sea Turtles (*Caretta caretta*) Departing from East-Central Florida Nesting Beaches. *Chelonian Conservation and Biology*, 4(3), 530-536.
- Dodd Jr., C. K. (1988). *Synopsis of the Biological Data on the Loggerhead Sea Turtle *Caretta caretta* (Linnaeus 1758)*. (Biological Report 88(14)). Washington, DC: US Department of the Interior, Fish and Wildlife Service. p. 110.
- Dodge, K. L., Galuardi, B., Miller, T. J., & Lutcavage, M. E. (2014). Leatherback turtle movements, dive behavior, and habitat characteristics in ecoregions of the Northwest Atlantic Ocean. *PLoS One*, 9(3), e91726.
- Doksaeter, L., Olsen, E., Nottestad, L., & Ferno, A. (2008). Distribution and Feeding Ecology of Dolphins along the Mid-Atlantic Ridge between Iceland and the Azores. *Deep Sea Research II*, 55, 243-253.
- Dolar, M. L. L., Suarez, P., Ponganis, P. J., & Kooyman, G. L. (1999). Myoglobin in Pelagic Small Cetaceans. *Journal of Experimental Biology*, 202(3), 227-236.
- Dolar, M. L. L., Walker, W. A., Kooyman, G. L., & Perrin, W. F. (2003). Comparative Feeding Ecology of Spinner Dolphins (*Stenella longirostris*) and Fraser's Dolphins (*Lagenodelphis hosei*) in the Sulu Sea. *Marine Mammal Science*, 19(1), 1-19.
- Dolphin, W. F. (1987a). Dive Behavior and Estimated Energy Expenditure of Foraging Humpback Whales in Southeast Alaska [USA]. *Canadian Journal of Zoology*, 65(2), 354-362.
- Dolphin, W. F. (1987b). Prey Densities and Foraging of Humpback Whales, *Megaptera novaeangliae*. *Experientia*, 43, 468-471.
- Domenici, P., Batty, R. S., Simila, T., & Ogam, E. (2000). Killer Whales (*Orcinus orca*) Feeding on Schooling Herring (*Clupea harengus*) Using Underwater Tail-Slaps: Kinematic Analyses of Field Observations. *The Journal of Experimental Biology*, 203, 283-294.
- Donahue, M. A., & Perryman, W. L. (2002). Pygmy Killer Whale. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1009-1010). San Diego, CA: Academic Press.
- Doniol-Valcroze, T., Lesage, V., Girard, J., & Michaud, R. (2011). Optimal foraging theory predicts diving and feeding strategies of the largest marine predator. *Behavioral Ecology*, 22(4), 880-888. doi: 10.1093/beheco/arr038.

- Dorsey, E. M., Stern, S. J., Hoelzel, A. R., & Jacobsen, J. (1990). Minke Whales (*Balaenoptera acutorostrata*) from the West Coast of North America: Individual Recognition and Small-scale Site Fidelity. *Reports of the International Whaling Commission*(Special Issue 12), 357-368.
- Doyle, T. K., Houghton, J. D. R., O'Suilleabháin, P. F., Hobson, V. J., Marnell, F., Davenport, J., & Hays, G. C. (2008). Leatherback Turtles Satellite-Tagged in European Waters. *Endangered Species Research*, 4, 23-31.
- Dulau-Drouot, V., Boucaud, V., & Rota, B. (2008). Cetacean Diversity off La Reunion Island (France). *Journal of the Marine Biological Association of the United Kingdom*, 88(6), 1263-1272.
- Dunham, J. S., & Duffus, D. A. (2002). Diet of Gray Whales (*Eschrichtius robustus*) in Clayoquot Sound, British Columbia, Canada. *Marine Mammal Science*, 18(2), 419-437.
- Eckert, K. L. (1995). Anthropogenic Threats to Sea Turtles. In: Bjorndal, K. A. (Ed.), *Biology and Conservation of Sea Turtles* (Revised Edition ed., pp. 611-612). Washington, DC: Smithsonian Institution Press.
- Eckert, S. A., Nellis, D. W., Eckert, K. L., & Kooyman, G. L. (1986). Diving Patterns of Two Leatherback Sea Turtles (*Dermochelys coriacea*) During Internesting Intervals at Sandy Point, St. Croix, U.S. Virgin Islands. *Herpetologica*, 42(3), 381-388.
- Edwards, H. H., Martin, J., Deutsch, C. J., Muller, R. G., Koslovsky, S. M., Smith, A. J., & Barlas, M. E. (2016). Influence of Manatees' Diving on Their Risk of Collision with Watercraft. *PLoS one*, 11(4).
- Eguchi, T., & Harvey, J. T. (2005). Diving Behavior of the Pacific Harbor Seal (*Phoca vitulina richardii*) in Monterey Bay, California. *Marine Mammal Science*, 21(2), 283-295.
- Erhart, L. M., Bagley, D. A., & Redfoot, W. E. (2003). *Loggerhead Turtles in the Atlantic Ocean: Geographic Distribution, Abundance, and Population Status*. . Washington, D.C.: Smithsonian Books.
- Ernst, C. H., Barbour, R. W., & Lovich, J. E. (1994). *Turtles of the United States and Canada*. Washington DC: Smithsonian Institution Press.
- Etheridge, K., Rathbun, G. B., Powell, J. A., & Kochman, H. I. (1985). Consumption of Aquatic Plants by the West Indian Manatee. *Journal of Aquatic Plant Management*, 23, 21-25.
- Evans, W. E. (1974). Telemetry of Temperature and Depth Data From a Free Ranging Yearling California Gray Whale, *Eschrichtius robustus*. *Marine Fisheries Review*, 36(4), 52-58.
- Evans, W. E. (1975). *Distribution, Differentiation of Populations, and Other Aspects of the Natural History of Delphinus delphis Linnaeus in the Northeastern Pacific*. Doctor of Philosophy (Ph.D. Dissertation), University of California, Los Angeles, Los Angeles.
- Evans, W. E. (1994). Common Dolphin, White-Bellied Porpoise--*Delphinus delphis* Linnaeus, 1758. In: Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals. Volume 5: The First Book of Dolphins* (Vol. 5, pp. 191-224). New York: Academic Press.
- Feldkamp, S. D., DeLong, R. L., & Antonelis, G. A. (1989). Diving Patterns of California Sea Lions, *Zalophus californianus*. *Canadian Journal of Zoology*, 67, 872-883.

- Fernandez, R., Santos, M. B., Carrillo, M., Tejedor, M., & Pierce, G. J. (2009). Stomach Contents of Cetaceans Stranded in the Canary Islands 1996–2006. *Journal of the Marine Biological Association of the United Kingdom*, 89(5), 873-883.
- Findlay, K. P., Best, P. B., Ross, G. J. B., & Cockcroft, V. G. (1992). The Distribution of Small Odontocete Cetaceans off the Coasts of South Africa and Namibia. *South African Journal of Marine Science*, 12, 237-270.
- Fiscus, C. H., Rice, D. W., & Wolman, A. A. (1989). *Cephalopods from the Stomachs of Sperm Whales taken off California*. (NOAA Technical Report NMFS 83). National Oceanic and Atmospheric Administration, National Marine Fisheries Service. pp. 1-12.
- Fitch, J. E., & Brownell, R. L., Jr. (1968). Fish Otoliths in Cetacean Stomachs and Their Importance in Interpreting Feeding Habits. *Journal of the Fisheries Research Board of Canada*, 25(12), 2561-2574.
- Flinn, R. D., Trites, A. W., Gregr, E. J., & Perry, R. I. (2002). Diets of Fin, Sei, and Sperm Whales in British Columbia: An Analysis of Commercial Whaling Records, 1963-1967. *Marine Mammal Science*, 18(3), 663-679.
- Folkow, L. P., & Blix, A. S. (1999). Diving Behaviour of Hooded Seals (*Cystophora cristata*) in the Greenland and Norwegian Seas. *Polar Biology*, 22(1), 61-74.
- Folkow, L. P., Nordoy, E. S., & Blix, A. S. (2004). Distribution and Diving Behaviour of Harp Seals (*Pagophilus groenlandicus*) from the Greenland Sea Stock. *Polar Biology*, 27, 281-298.
- Ford, J. K. B. (2002). Killer Whale. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 669-676). San Diego, CA: Academic Press.
- Ford, J. K. B., Ellis, G. M., Barrett-Lennard, L. G., Morton, A. B., Palm, R. S., & III, K. C. B. (1998). Dietary Specialization in Two Sympatric Populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. *Canadian Journal of Zoology*, 76, 1456-1471.
- Forney, K. A. (2007). *Preliminary Estimates of Cetacean Abundance along the U.S. West Coast and within Four National Marine Sanctuaries during 2005*. (NOAA-TM-NMFS-SWFSC-406). Santa Cruz, CA: National Oceanic and Atmospheric Administration. p. 36.
- Forney, K. A., & Barlow, J. (1998). Seasonal Patterns in the Abundance and Distribution of California Cetaceans, 1991-1992. *Marine Mammal Science*, 14(3), 460-489.
- Forney, K. A., Becker, E. A., Foley, D. G., Barlow, J., & Oleson, E. M. (2015). Habitat-based models of cetacean density and distribution in the central North Pacific. *Endangered Species Research*, 27(1), 1.
- Fossette, S., Ferraroli, S., Tanaka, H., Ropert-Coudert, Y., Arai, N., & Sato, K. (2007). Dispersal and Dive Patterns in Gravid Leatherback Turtles during the Nesting Season in French Guiana. *Marine Ecology Progress Series*, 338, 233-247.
- Friedlaender, A. S., Hazen, E. L., Nowacek, D. P., Halpin, P. N., Ware, C., Weinrich, M. T., . . . Wiley, D. (2009a). Diel Changes in Humpback Whale *Megaptera novaeangliae* Feeding Behavior in Response to Sand Lance *Ammodytes* spp. Behavior and Distribution. *Marine Ecology Progress Series*, 395, 91-100.

- Friedlaender, A. S., Lawson, G. L., & Halpin, P. N. (2009b). Evidence of Resource Partitioning Between Humpback and Minke Whales Around the Western Antarctic Peninsula. *Marine Mammal Science*, 25(2), 402-415.
- Frost, K. J., Simpkins, M. A., & Lowry, L. F. (2001). Diving Behavior of Subadult and Adult Harbor Seals in Prince William Sound, Alaska. *Marine Mammal Science*, 17(4), 813-834.
- Gallo-Reynoso, J.-P., & Figueroa-Carranza, A.-L. (1995). Occurrence of Bottlenose Whales in the Waters of Isla Guadalupe, Mexico. *Marine Mammal Science*, 11(4), 573-575.
- Gallo-Reynoso, J. P., Figueroa-Carranza, A.-L., & Le Boeuf, B. (2008). Foraging behavior of lactating Guadalupe fur seal females. *Avances en el Estudio de los Mamíferos de México, C. Lorenzo, E. Espinoza and J. Ortega (eds.). Publicaciones Especiales*, 2, 595-614.
- Gannon, D. P., Read, A. J., Craddock, J. E., Fristrup, K. M., & Nicolas, J. R. (1997). Feeding Ecology of Long-Finned Pilot Whales *Globicephala melas* in the Western North Atlantic. *Marine Ecology Progress Series*, 148(1), 1-10.
- García-Rodríguez, F. J., & Aurióles-Gamboa, D. (2004). Spatial and Temporal Variation In the Diet of the California Sea Lion (*Zalophus californianus*) in the Gulf of California, Mexico. *Fishery Bulletin*, 102(1), 47-62.
- Gelatt, T., & Lowry, L. (2008). *Callorhinus ursinus*. Retrieved from <http://www.iucnredlist.org/> as accessed on 11 August 2010.
- Gentry, R. L. (2002). Northern Fur Seal. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 813-817). San Diego, CA: Academic Press.
- Gentry, R. L., Kooyman, G. L., & Goebel, M. E. (1986). Feeding and Diving Behavior of Northern Fur Seals. In Gentry, R. L. & Kooyman, G. L. (Eds.), *Fur Seals: Maternal Strategies on Land and at Sea* (pp. 61-78). Princeton: Princeton University Press.
- Gero, S., Engelhaupt, D., Rendell, L., & Whitehead, H. (2009). Who Cares? Between-Group Variation in Alloparental Caregiving in Sperm Whales. *Behavioral Biology*, 20(4), 838-843.
- Gero, S., & Whitehead, H. (2006). *Opportunistic Sightings of Small Cetaceans off the Leeward Shore of the Commonwealth of Dominica*. ((SC/58/SM1)). International Whaling Commission. p. 7.
- Gilmartin, W. G., & Forcada, J. (2002). Monk Seals. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 756-759). San Diego, CA: Academic Press.
- Gitschlag, G. R. (1996). Migration and Diving Behavior of Kemp's Ridley (Garman) Sea Turtles Along the U.S. Southeastern Atlantic Coast. *Journal of Experimental Marine Biology and Ecology*, 205, 115-135.
- Gjertz, I., Kovacs, K. M., Lydersen, C., & Wiig, O. (2000). Movements and Diving of Adult Ringed Seals (*Phoca hispida*) in Svalbard. *Polar Biology*, 23(9), 651-656.
- Gjertz, I., Lydersen, C., & Wiig, O. (2001). Distribution and Diving of Harbour Seals (*Phoca vitulina*) in Svalbard. *Polar Biology*, 24(3), 209-214.

- Godley, B. J., Richardson, S., Broderick, A. C., Coyne, M. S., Glen, F., & Hays, G. C. (2002). Long-Term Satellite Telemetry of the Movements and Habitat Utilisation by Green Turtles in the Mediterranean. *Ecography*, *25*, 352-362.
- Goebel, M. E., Bengtson, J. L., DeLong, R. L., Gentry, R. L., & Loughlin, T. R. (1991). Diving Patterns and Foraging Locations of Female Northern Fur Seals. *U S National Marine Fisheries Service Fishery Bulletin*, *89*(2), 171-180.
- Goff, G. P., & Lien, J. (1988). Atlantic Leatherback Turtles, *Dermochelys coriacea*, in Cold Water off Newfoundland and Labrador. *Canadian Field-Naturalist*, *102*(1), 1-5.
- Goldbogen, J. A., Calambokidis, J., Croll, D. A., Harvey, J. T., Newton, K. M., Oleson, E. M., . . . Shadwick, R. E. (2008). Foraging Behavior of Humpback Whales: Kinematic and Respiratory Patterns Suggest a High Cost for a Lunge. *The Journal of Experimental Biology*, *211*, 3712-3719.
- Goldbogen, J. A., Calambokidis, J., Croll, D. A., McKenna, M. F., Oleson, E., Potvin, J., . . . Tershy, B. R. (2012). Scaling of lunge-feeding performance in rorqual whales: Mass-specific energy expenditure increases with body size and progressively limits diving capacity. *Functional Ecology*, *26*, 216-226. doi: 10.1111/j.1365-2435.2011.01905.x.
- Goldbogen, J. A., Calambokidis, J., Friedlaender, A. S., Francis, J., DeRuiter, S. L., Stimpert, A. K., . . . Southall, B. L. (2013). Underwater acrobatics by the world's largest predator: 360° rolling manoeuvres by lunge-feeding blue whales. *Biology Letters*, *9*, 20120986. doi: 10.1098/rsbl.2012.0986.
- Goldbogen, J. A., Calambokidis, J., Shadwick, R. E., Oleson, E. M., McDonald, M. A., & Hildebrand, J. A. (2006). Kinematics of Foraging Dives and Lunge-Feeding in Fin Whales. *The Journal of Experimental Biology*, *209*, 1231-1244.
- Goodman-Lowe, G. D. (1998). Diet of the Hawaiian Monk Seal (*Monachus schauinslandi*) from the Northwestern Hawaiian Islands During 1991 to 1994. *Marine Biology*, *132*, 535-546.
- Gowans, S. (2002). Bottlenose Whale. In: Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 128-129). San Diego, CA: Academic Press.
- Gowans, S., & Whitehead, H. (1995). Distribution and Habitat Partitioning by Small Odontocetes in the Gully, a Submarine Canyon on the Scotian Shelf. *Canadian Journal of Zoology*, *73*, 1599-1608.
- Gowans, S., Whitehead, H., & Hooker, S. K. (2001). Social Organization in Northern Bottlenose Whales, *Hyperoodon ampullatus*: Not Driven by Deep-Water Foraging? *Animal Behaviour*, *62*(2), 369-377.
- Green, G. A., Brueggeman, J. J., Grotefendt, R. A., Bowlby, C. E., Bonnell, M. L., & K.C. Balcomb III. (1992). *Cetacean Distribution and Abundance off Oregon and Washington, 1989-1990*. Los Angeles, California: Minerals Management Service. pp. 1-1 to 1-100.
- Griffin, R. B. (1999). Sperm Whale Distributions and Community Ecology Associated with a Warm-Core Ring off Georges Bank. *Marine Mammal Science*, *15*(1), 33-51.
- Griffin, R. B. (2005). *Movement Patterns and Diving Behavior of Atlantic Spotted Dolphins (Stenella frontalis) in Relation to Oceanographic Features: A Study Using Remotely-*

- deployed Suction-cup Attached Tags*. (Mote Marine Laboratory Technical Report Number 1005). Sarasota, FL: Mote Marine Laboratory. p. 102.
- Gross, A., Kiszka, J., Canneyt, O. V., Richard, P., & Ridoux, V. (2009). A Preliminary Study of Habitat and Resource Partitioning among Co-occurring Tropical Dolphins around Mayotte, Southwest Indian Ocean. *Estuarine, Coastal and Shelf Science*, 84, 367-374.
- Guerrero, J. A. (1989). *Feeding Behavior of Gray Whales in Relation to Patch Dynamics of their Benthic Prey*. Master of Science, San José State University, San José, CA.
- Hain, J. H. W., Carter, G. R., Kraus, S. D., Mayo, C. A., & Winn, H. E. (1982). Feeding Behavior of the Humpback Whale, *Megaptera novaeangliae*, in the Western North Atlantic. *Fishery Bulletin*, 80(2), 259-268.
- Hakoyama, H., Le Boeuf, B. J., Naito, Y., & Sakamoto, W. (1994). Diving behavior in relation to ambient water temperature in northern elephant seals. *Canadian Journal of Zoology*, 72, 643-651.
- Hall, J. D. (1970). *Conditioning Pacific White Stripped Dolphins, Lagenorhynchus obliquidene, For Open Ocean Release*. (Naval Undersea Center Technical Publication 200). San Diego, CA: Naval Undersea Research and Development Center. p. 13.
- Hanser, S., Becker, E. A., & Kumar, A. (2012). *Pacific Navy Marine Species Density Database Technical Report*.
- Harvey, V., Côté, S. D., & Hammill, M. O. (2008). The Ecology of 3-D Space Use in a Sexually Dimorphic Mammal. *Ecography*, 31, 371-380.
- Hastings, K. K., Frost, K. J., Simpkins, M. A., Pendleton, G. W., Swain, U. G., & Small, R. J. (2004). Regional Differences in Diving Behavior of Harbor Seals in the Gulf of Alaska. *Canadian Journal of Zoology*, 82, 1755-1773.
- Hatase, H., Sata, K., Yamaguchi, M., Takahashi, K., & Tsukamoto, K. (2006). Individual Variation in Feeding Habitat Use by Adult Female Green Sea Turtles (*Chelonia mydas*): Are They Obligately Neritic Herbivores. *Oecologia*, 149, 52-64.
- Haug, T., Gjøsaeter, H., Lindstrøm, U., & Nilssen, K. T. (1995). Diet and Food Availability for North-East Atlantic Minke Whales (*Balaenoptera acutorostrata*), During the Summer of 1992. *ICES Journal of Marine Science*, 52(1), 77-86.
- Haug, T., Nilssen, K. T., & Lindblom, L. (2004). Feeding Habits of Harp and Hooded Seals in Drift Ice Waters along the East Coast of Greenland in Summer and Winter. *Polar Research*, 23(1), 35-42.
- Hauksson, E., & Bogason, V. (1997). Comparative Feeding of Grey (*Halichoerus grypus*) and Common Seals (*Phoca vitulina*) in Coastal Waters of Iceland, with a Note on the Diet of Hooded (*Cystophora cristata*) and Harp Seals (*Phoca groenlandica*). *Journal of Northwest Atlantic Fishery Science*, 22, 125-135.
- Hawkes, L. A., Broderick, A. C., Coyne, M. S., Godfrey, M. H., Lopez-Jurado, L.-F., Lopez-Suarez, P., . . . Godley, B. J. (2006). Phenotypically Linked Dichotomy in Sea Turtle Foraging Requires Multiple Conservation Approaches. *Current Biology*, 16, 990-995.

- Hays, G. C., Adams, C. R., Broderick, A. C., Godley, B. J., Lucas, D. J., Metcalfe, J. D., & Prior, A. A. (2000). The Diving Behaviour of Green Turtles at Ascension Island. *Animal Behaviour*, *59*, 577-586.
- Hays, G. C., Hobson, V. J., Metcalfe, J. D., Righton, D., & Sims, D. W. (2006). Flexible Foraging Movements of Leatherback Turtles Across the North Atlantic Ocean. *Ecology*, *87*(10), 2647-2656.
- Hays, G. C., Houghton, J. D. R., Isaacs, C., King, R. S., Lloyd, C., & Lovell, P. (2004). First Records of Oceanic Dive Profiles for Leatherback Turtles, *Dermochelys coriacea*, Indicate Behavioural Plasticity Associated with Long-Distance Migration. *Animal Behaviour*, *67*, 733-743.
- Hazen, E. L., Friedlaender, A. S., Thompson, M. A., Ware, C. R., Weinrich, M. T., Halpin, P. N., & Wiley, D. N. (2009). Fine-Scale Prey Aggregations and Foraging Ecology of Humpback Whales *Megaptera novaeangliae*. *Marine Ecology Progress Series*, *395*, 75-89.
- Heaslip, S. G., Iverson, S. J., Bowen, W. D., & James, M. C. (2012). Jellyfish support high energy intake of leatherback sea turtles (*Dermochelys coriacea*): video evidence from animal-borne cameras. *PLoS ONE*, *7*(3), e33259. doi: 10.1371/journal.pone.0033259.
- Heath, C. B. (2002). California, Galapagos, and Japanese Sea Lions. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 180-186). San Diego, CA: Academic Press.
- Heide-Jorgensen, M. P., Bloch, D., Stefansson, E., Mikkelsen, B., Ofstad, L. H., & Dietz, R. (2002). Diving Behaviour of Long-Finned Pilot Whales *Globicephala melas* Around the Faroe Islands. *Wildlife Biology*, *8*(4), 307-313.
- Heise, K. (1997). Diet and Feeding Behaviour of Pacific White-sided Dolphins (*Lagenorhynchus obliquidens*) as Revealed Through the Collection of Prey Fragments and Stomach Content Analyses. *Reports of the International Whaling Commission*, *47*, 807-815.
- Herman, J. S., Kitchener, A. C., Baker, J. R., & Lockyer, C. (1994). The Most Northerly Record of Blainville's Beaked Whale, *Mesoplodon densirostris*, from the Eastern Atlantic. *Mammalia*, *58*(4), 657-661.
- Herzing, D. L., & Elliser, C. R. (2013). Nocturnal feeding of Atlantic spotted dolphins in the Bahamas. *Marine Mammal Science*.
- Heyning, J. E. (1989). Cuvier's Beaked Whale -- *Ziphius cavirostris* G. Cuvier, 1823. In Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals: Volume 4: River Dolphins and the Larger Toothed Whales* (Vol. 4, pp. 289-308). New York: Academic Press.
- Heyning, J. E., & Perrin, W. F. (1994). Evidence for Two Species of Common Dolphins (Genus *Delphinus*) from the Eastern North Pacific. *Contributions in Science*, *442*, 1-35.
- Hickmott, L. S. (2005). *Diving Behaviour and Foraging Behaviour and Foraging Ecology of Blainville's and Cuvier's Beaked Whales in the Northern Bahamas*. Master of Research in Environmental Biology (Master's), University of St. Andrews, Scotland.
- Hindell, M. A. (2002). Elephant Seals: *Mirounga angustirostris* and *M. leonina*. In Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 370-373). San Diego, CA: Academic Press.

- Hobson, K. A., & Sease, J. L. (1998). Stable Isotope Analyses of Tooth Annuli Reveal Temporal Dietary Records: An Example Using Steller Sea Lions. *Marine Mammal Science*, 14(1), 116-129.
- Hobson, K. A., Sease, J. L., Merrick, R., L., & Piatt, J., F. (1997). Investigating Trophic Relationships of Pinnipeds in Alaska and Washington Using Stable Isotope Ratios of Nitrogen and Carbon. *Marine Mammal Science*, 13(1), 114-132.
- Hochscheid, S., Bentivegna, F., Hamza, A., & Hays, G. C. (2010). When surfacers do not dive: multiple significance of extended surface times in marine turtles. *the Journal of Experimental Biology*, 213, 1328-1337.
- Hochscheid, S., Godley, B. J., Broderick, A. C., & Wilson, R. P. (1999). Reptilian Diving: Highly Variable Dive Patterns in the Green Turtle *Chelonia mydas*. *Marine Ecology Progress Series*, 185, 101-112.
- Hooker, S., Fahlman, A., Aguilar de Soto, N., Bernaldo de Quirós, Y., Brubakk, A., Costidis, A., . . . Garner, M. (2011). Deadly Diving? Physiological and Behavioural Management of Decompression Stress in Diving Mammals.
- Hooker, S. K. (1998). Extensive Scarring Observed on Female or Juvenile Male Sperm Whales off the Galápagos Islands. *Mammalia*, 62(1), 134-139.
- Hooker, S. K., & Baird, R. W. (1999). Deep-Diving Behaviour of the Northern Bottlenose Whale, *Hyperoodon ampullatus* (Cetacea: Ziphiidae). *Proceedings of the Royal Society of London: Series B Biological Sciences*, 266(1420), 671-676.
- Hooker, S. K., Iverson, S. J., Ostrom, P., & Smith, S. C. (2001). Diet of Northern Bottlenose Whales Inferred from Fatty-Acid and Stable-Isotope Analyses of Biopsy Samples. *Canadian Journal of Zoology*, 79, 1442-1454.
- Hooker, S. K., Whitehead, H., & Gowans, S. (1999). Marine Protected Area Design and the Spatial and Temporal Distribution of Cetaceans in a Submarine Canyon. *Conservation Biology*, 13(3), 592-602.
- Horwood, J. (2002). Sei Whale. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1069-1071). San Diego, CA: Academic Press.
- Houghton, J. D. R., Broderick, A. C., Godley, B. J., Metcalfe, J. D., & Hays, G. C. (2002). Diving Behaviour During the Internesting Interval for Loggerhead Turtles *Caretta caretta* Nesting in Cyprus. *Marine Ecology Progress Series*, 227, 63-70.
- Houghton, J. D. R., Doyle, T. K., Davenport, J., Wilson, R. P., & Hays, G. C. (2008). The Role of Infrequent and Extraordinary Deep Dives in Leatherback Turtles (*Dermochelys coriacea*). *The Journal of Experimental Biology*, 211, 2566-2575.
- Howell, E. A., Dutton, P. H., Polvina, J. J., Bailey, H., Parker, D. M., & Balazs, G. H. (2010). Oceanographic Influences on the Dive Behavior of Juvenile Loggerhead Turtles (*Caretta caretta*) in the North Pacific Ocean. *Marine Biology*, 157, 1011-1026.
- Iverson, S. J., Field, C., Bowen, W. D., & Blanchard, W. (2004). Quantitative Fatty Acid Signature Analysis: A New Method of Estimating Predator Diets. *Ecological Monographs*, 74(2), 211-235.

- James, M. C., Davenport, J., & Hays, G. C. (2006a). Expanded Thermal Niche for a Diving Vertebrate: A Leatherback Turtle Diving into Near-Freezing Water. *Journal of Experimental Marine Biology and Ecology*, 335, 221-226.
- James, M. C., Eckert, S. A., & Myers, R. A. (2005a). Migratory and Reproductive Movements of Male Leatherback Turtles (*Dermochelys coriacea*). *Marine Biology*, 147, 845-853.
- James, M. C., Myers, R. A., & Ottensmeyer, C. A. (2005b). Behaviour of Leatherback Sea Turtles, *Dermochelys coriacea*, During the Migratory Cycle. *Proceedings of the Royal Society of London, Part B*, 272, 1547-1555.
- James, M. C., Ottensmeyer, C. A., Eckert, S. A., & Myers, R. A. (2006b). Changes in Diel Diving Patterns Accompany Shifts between Northern Foraging and Southward Migration in Leatherback Turtles. *Canadian Journal of Zoology*, 84, 754-765.
- Jefferson, T. A. (1988). *Phocoenoides dalli*. *Mammalian Species*, 319, 1-7.
- Jefferson, T. A. (2002a). Clymene Dolphin *Stenella clymene*. In: Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of marine mammals* (pp. 234-236). San Diego: Academic Press.
- Jefferson, T. A. (2002b). Dall's Porpoise: *Phocoenoides dalli*. In: Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 308-310). San Diego, CA: Academic Press.
- Jefferson, T. A., & Barros, N. B. (1997). *Peponocephala electra*. *Mammalian Species*, 553, 1-6.
- Jefferson, T. A., Fertl, D., Bolaños-Jiménez, J., & Zerbini, A. N. (2009). Distribution of Common Dolphins (*Delphinus spp.*) in the Western Atlantic Ocean: A Critical Re-examination. *Marine Biology*, 156, 1109-1124.
- Jefferson, T. A., & Newcomer, M. W. (1993a). *Lissodelphis borealis*. *Mammalian Species*, 425, 1-6.
- Jefferson, T. A., Newcomer, M. W., Leatherwood, S., & Van Waerebeek, K. (1994). Right Whale Dolphins--*Lissodelphis borealis* (Peale, 1848) and *Lissodelphis peronii* (Lacépède, 1804). In: Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals. Volume 5: The First Book of Dolphins* (Vol. 5, pp. 335-362). New York: Academic Press.
- Jefferson, T. A., & Newcomer, W. W. (1993b). *Lissodelphis borealis*. *Mammalian Species*, 425, 1-6.
- Jefferson, T. A., Webber, M. A., & Pitman, R. L. (2008). *Marine Mammals of the World: A Comprehensive Guide to Their Identification*. San Diego, CA: Academic Press.
- Jensen, F. H., Perez, J. M., Johnson, M., Soto, N. A., & Madsen, P. T. (2011). Calling under pressure: short-finned pilot whales make social calls during deep foraging dives. *Proceedings of the Royal Society Biological Sciences*, 278, 3017-3025. doi: 10.1098/rspb.2010.2604.
- Johanos, T. C., & Baker, J. D. (2002). *The Hawaiian Monk Seal in the Northwestern Hawaiian Islands, 2000*. (NOAA-TM-NMFS-SWFSC-340). Honolulu, Hawaii: Honolulu Laboratory, SWFSC, NMFS, NOAA.

- Johnson, M., Madsen, P., Zimmer, W. M. X., de Soto, N. A., & Tyack, P. (2004). Beaked Whales Echolocate On Prey. *Proceedings of the Royal Society of London B (Suppl.)*, 4.
- Johnson, M. P., & Aguilar de Soto, N. (2008a). Mesoplodon densirostris El Hierro, Canary Islands, Spain 10/11/2003 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 11 October 2003 from <http://hdl.handle.net/1912/2367>.
- Johnson, M. P., & Aguilar de Soto, N. (2008b). Mesoplodon densirostris El Hierro, Canary Islands, Spain 10/25/2003 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 25 October 2003 from <http://hdl.handle.net/1912/2366>.
- Johnson, M. P., & Sturlese, A. (2008a). Ziphius cavirostris Liguria, Italy 9/17/2003 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 17 September 2003
- Johnson, M. P., & Sturlese, A. (2008b). Ziphius cavirostris Liguria, Italy 9/20/2003 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 20 September 2003
- Johnston, D. W., Westgate, A. J., & Read, A. J. (2005). Effects of Fine-Scale Oceanographic Features on the Distribution and Movements of Harbour Porpoises *Phocoena phocoena* in the Bay of Fundy. *Marine Ecology Progress Series*, 295, 279-293.
- Jones, M. L., & Swartz, S. L. (2002). Gray Whale (*Eschrichtius robustus*). In Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 524-536). San Diego, CA: Academic Press.
- Jonsen, I. D., Myers, R. A., & James, M. C. (2007). Identifying Leatherback Turtle Foraging Behavior from Satellite Telemetry using a Switching State-Space Model. *Ecology Progress Series*, 337, 255-264.
- Jorgensen, C., Lydersen, C., Brix, O., & Kovacs, K. M. (2001). Diving development in nursing harbour seal pups. *Journal of Experimental Biology*, 204, 3993-4004.
- Joyce, G. G., Sigurjonsson, J., & Vikingsson, G. (1990). Radio Tracking a Minke Whale (*Balaenoptera acutorostrata*) in Icelandic Waters for the Examination of Dive-time Patterns. *Reports of the International Whaling Commission*, 40, 357-361.
- Kajimura, H., & Loughlin, T. R. (1988). Marine Mammals in the Oceanic Food Web of the Eastern Subarctic Pacific. *Bulletin of the Ocean Research Institute, University of Tokyo*, 26, 187-223.
- Kasuya, T. (1986). Distribution and Behavior of Baird's Beaked Whales off the Pacific Coast of Japan. *The Scientific Reports of the Whales Research Institute*, 37, 61-83.
- Kasuya, T. (2002). Giant Beaked Whales. In Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 519-522). San Diego, CA: Academic Press.
- Kato, H. (2002). Bryde's Whales. In Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 171-177). San Diego, CA: Academic Press.
- Kawakami, T. (1980). A Review of Sperm Whale Food. *Scientific Reports of the Whales Research Institute*, 32, 199-218.
- Kenney, R. D. (1990). Bottlenose Dolphins off the Northeastern United States. In Leatherwood, S. & Reeves, R. R. (Eds.), *The Bottlenose Dolphin* (pp. 369-386). New York: Academic Press, Inc.

- Kenney, R. D. (2002). North Atlantic, North Pacific, and Southern Right Whales (*Eubalaena glacialis*, *E. japonica*, and *E. australis*). In: Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals*. San Diego, CA: Academic Press.
- Kenney, R. D., Mayo, C. A., & Winn, H. E. (2001). Migration and Foraging Strategies at Varying Spatial Scales in Western North Atlantic Right Whales: A Review of Hypotheses. *Journal of Cetacean Research and Management, Special Issue 2*, 251-260.
- Kenyon, K. W., & Rice, D. W. (1959). Life History of the Hawaiian Monk Seal. *Pacific Science, XIII*, 215-252.
- Kingsley, M. C. S., & Reeves, R. R. (1998). Aerial Surveys of Cetaceans in the Gulf of St. Lawrence in 1995 and 1996. *Canadian Journal of Zoology, 76*, 1529-1550.
- Kinze, C. C. (2002). White-Beaked Dolphin. In: Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1332-1334). San Diego, CA: Academic Press.
- Kiszka, J., Macleod, K., Canneyt, O. V., Walker, D., & Ridoux, V. (2007). Distribution, Encounter Rates, and Habitat Characteristics of Toothed Cetaceans in the Bay of Biscay and Adjacent Waters from Platform-of-Opportunity Data. *Oxford Journals*, 1033-1043.
- Klatsky, L. J. (2004). *Movement and dive behavior of bottlenose dolphins (Tursiops truncatus) near the Bermuda Pedestal*. (masters), San Diego State University;Biology.
- Kooyman, G. L., Gentry, R. L., & Urquhart, D. L. (1976). Northern Fur Seal Diving Behavior: A New Approach to its Study. *Science, 193*(4251), 411-412.
- Kuhn, C. (2004). *Assessing the Feeding Behavior of California sea lions*. Santa Cruz, CA: University of California. p. 9.
- Kuhn, C. E. (2006). *Measuring At-Sea Feeding to Understand the Foraging Behavior of Pinnipeds*. PhD, University of California Santa Cruz.
- Kuhn, C. E., Johnson, D. S., Ream, R. R., & Gelatt, T. S. (2009). Advances in the Tracking of Marine Species: Using GPS Locations to Evaluate Satellite Track Data and a Continuous-Time Movement Model. *Marine Ecology Progress Series, 393*, 97-109.
- Kvadsheim, P. H., Miller, P. J., Tyack, P. L., Sivle, L. D., Lam, F.-P. A., & Fahlman, A. (2012). Estimated tissue and blood N₂ levels and risk of decompression sickness in deep-, intermediate-, and shallow-diving toothed whales during exposure to naval sonar. *Frontiers in physiology, 3*.
- Labansen, A. L., Lydersen, C., Haug, T., & Kovacs, K. M. (2007). Spring Diet of Ringed Seals (*Phoca hispida*) from Northwestern Spitsbergen, Norway. *ICES Journal of Marine Science, 64*, 1246-1256.
- Laerm, J., Wenzel, F., Craddock, J. E., Weinand, D., McGurk, J., Harris, M. J., . . . Barros, N. B. (1997). New Prey Species for Northwestern Atlantic Humpback Whales. *Marine Mammal Science, 13*(4), 705-711.
- Laidre, K. L., Heide-Jørgensen, M. P., Heagerty, P., Cossio, A., Bergström, B., & Simon, M. (2010). Spatial Associations between Large Baleen Whales and their Prey in West Greenland. *Marine Ecology Progress Series, 402*, 269-284.

- Lammers, M. O. (2004). Occurrence and Behavior of Hawaiian Spinner Dolphins (*Stenella longirostris*) Along Oahu's Leeward and South Shores. *Aquatic Mammals*, 30(2), 237-250.
- Lander, M. E., Gulland, F. M. D., & DeLong, R. L. (2000). Satellite Tracking a Rehabilitated Guadalupe Fur Seal (*Arctocephalus townsendi*). *Aquatic Mammals*, 26(2), 137-142.
- Landry, A. M., & Costa, D. (1999). Status of Sea Turtle Stocks in the Gulf of Mexico with Emphasis on the Kemp's Ridley. In: Kumpf, H., Steidinger, K. & Sherman, K. (Eds.), *The Gulf of Mexico Large Marine Ecosystem: Assessment, Sustainability, and Management* (pp. 248-268). New York: Blackwell Science.
- Lavigne, D. M. (2002). Harp Seal. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 560-562). San Diego, CA: Academic Press.
- Lawson, J. W., Anderson, J. T., Dalley, E. L., & Stenson, G. B. (1998). Selective Foraging by Harp Seals *Phoca groenlandica* in Nearshore and Offshore Waters of Newfoundland, 1993 and 1994. *Marine Ecology Progress Series*, 163, 1-10.
- Lazell Jr., J. D. (1980). New England Waters: Critical Habitat for Marine Turtles. *Copeia*, 1980(2), 290-295.
- Le Boeuf, B. J., Costa, D. P., Huntley, A. C., & Feldkamp, S. D. (1988). Continuous, Deep Diving in Female Northern Elephant Seals, *Mirounga angustirostris*. *Canadian Journal of Zoology*, 66(2), 446-458.
- Le Boeuf, B. J., Costa, D. P., Huntley, A. C., Kooyman, G. L., & Davis, R. W. (1986). Pattern and depth of dives in Northern elephant seals, *Mirounga angustirostris*. *The Zoological Society of London*, 208, 1-7.
- Le Boeuf, B. J., Crocker, D. E., Blackwell, S. B., Morris, P. A., & Thorson, P. H. (1993). Sex Differences in Diving and Foraging Behavior of Northern Elephant Seals. In: Boyd, I. L. (Ed.), *Marine Mammals: Advances in Behavioural and Population Biology* (Vol. 66, pp. 149-178). Oxford: Clarendon Press.
- Leatherwood, S., & Reeves, R. R. (1983). *The Sierra Club Handbook of Whales and Dolphins*. San Francisco: Sierra Club Books.
- Leatherwood, S., & Walker, W. A. (1979). The Northern Right Whale Dolphin *Lissodelphis borealis* Peale in the Eastern North Pacific. In: Winn, H. E. & Olla, B. L. (Eds.), *Behavior of Marine Animals: Current Perspectives in Research* (pp. 85-141). New York, NY: Plenum Press.
- LeDuc, R. G., Perrin, W. F., & Dizon, A. E. (1999). Phylogenetic Relationships Among the Delphinid Cetaceans Based on Full Cytochrome *b* Sequences. *Marine Mammal Science*, 15(3), 619-648.
- Lefebvre, L. W., Reid, J. P., Kenworthy, W. J., & Powell, J. A. (2000). Characterizing Manatee Habitat Use and Seagrass Grazing in Florida and Puerto Rico: Implications for Conservation and Management. *Pacific Conservation Biology*, 5, 289-298.
- Lidgard, D. C., Boness, D. J., Bowen, W. D., & McMillan, J. I. (2003). Diving Behaviour During the Breeding Season in the Terrestrially Breeding Male Grey Seal: Implications for Alternative Mating Tactics. *Canadian Journal of Zoology*, 81, 1025-1033.

- Ligon, A. D., & Baird, R. W. (2001, 28 November-3 December 2001). *Diving Behaviour of False Killer Whales off Maui and Lana'i, Hawaii*. Paper presented at the Abstracts, Fourteenth Biennial Conference on the Biology of Marine Mammals, Vancouver, British Columbia.
- Linnenschmidt, M., Akamatsu, T., & Miller, A. (2013). Biosonar, dive, and foraging activity of satellite tracked harbor porpoises (*Phocoena phocoena*). *Marine Mammal Science*, 29(2), E77-E97. doi: 10.1111/j.1748-7692.2012.00592.x.
- Littnan, C. L., Baker, J. D., Parrish, F. A., & Marshall, G. J. (2004). Effects of Video Camera Attachment on the Foraging Behavior of Immature Hawaiian Monk Seals. *Marine Mammal Science*, 20(2), 345-352.
- Ljungblad, D. K., Moore, S. E., Clarke, J., & Bennett, J. C. (1987). *Distribution, Abundance, Behavior and Bioacoustics of Endangered Whales in the Alaskan Beaufort and Eastern Chukchi Seas, 1979-86*. (1177). San Diego, CA: Naval Ocean Systems Center (NOSC). p. 390.
- Lodi, L., & Hetzel, B. (1999). Rough-Toothed Dolphin, *Steno bredanensis*, Feeding Behaviors in Ilha Grande Bay, Brazil. *Biociências*, 7(1), 29-42.
- Louella, M., & Dolar, L. (2002). Fraser's Dolphin. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 485-487). San Diego, CA: Academic Press.
- Lowry, M. S., & Carretta, J. V. (1999). Market Squid (*Loligo opalescens*) in the Diet of California Sea Lions (*Zalophus californianus*) in Southern California (1981-1995). *CalCOFI Reports*, 40, 196-207.
- Lutcavage, M., & Musick, J. A. (1985). Aspects of the Biology of Sea Turtles in Virginia. *Copeia*, 1985(2), 449-456.
- Lutcavage, M. E., & Lutz, P. L. (1997). Diving Physiology. In: Lutz, P. L. & Musick, J. A. (Eds.), *The Biology of Sea Turtles* (Vol. I, pp. 277-296). New York: CRC Press.
- Lydersen, C. (1991). Monitoring Ringed Seal (*Phoca hispida*) Activity by Means of Acoustic Telemetry. *Canadian Journal of Zoology*, 69, 1178-1182.
- Lydersen, C., & Kovacs, K. M. (1993). Diving Behaviour of Lactating Harp Seal, *Phoca groenlandica*, Females from the Gulf of St Lawrence, Canada. *Animal Behavior*, 46, 1213-1221.
- MacLeod, C. D., Santos, M. B., & Pierce, G. J. (2003a). Review of Data on Diets of Beaked Whales: Evidence of Niche Separation and Geographic Segregation. *Journal of the Marine Biological Association of the United Kingdom*, 83, 651-665.
- MacLeod, K., Simmonds, M. P., & Murray, E. (2003b). Summer Distribution and Relative Abundance of Cetacean Populations off North-West Scotland. *Journal of the Marine Biological Association of the United Kingdom*, 83, 1187-1192.
- Madsen, P. T., Kerr, I., & Payne, R. (2004a). Echolocation Clicks of Two Free-Ranging, Oceanic Delphinids with Different Food Preferences: False Killer Whales *Pseudorca crassidens* and Risso's Dolphins *Grampus griseus*. *The Journal of Experimental Biology*, 207, 1811-1823.

- Madsen, P. T., Payne, R., & Kerr, I. (2004b). Source Parameter Estimates of Echolocation Clicks from Wild Pygmy Killer Whales (*Feresa attenuata*) (L). *Journal of Acoustical Society of America*, 116(4), 1909 - 1912.
- Malcolm, C. D., & Duffus, D. A. (2000). Comparison of Subjective and Statistical Methods of Dive Classification Using Data from a Time-Depth Recorder Attached to a Gray Whale (*Eschrichtius robustus*). *Journal of Cetacean Research and Management*, 2(3), 177-182.
- Malcolm, C. D., Duffus, D. A., & Wischniowski, S. G. (1995). Small Scale Behaviour of Large Scale Subjects: Diving Behaviour of a Gray Whale (*Eschrichtius robustus*). *Western Geography*, 5/6, 35-44.
- Maravilla-Chavez, M. O., & Lowry, M. S. (1999). Incipient breeding colony of Guadalupe fur seals at isla Benito del este, Baja California, Mexico. *Marine Mammal Science*, 15(1), 239-241.
- Marine Turtle Specialist Group. (2004). *Marine Turtle Specialist Group Review: 2004 Global Status Assessment, Green Turtle (Chelonia mydas)*. The World Conservation Union (IUCN) Species Survival Commission, Red List Programme. p. 71.
- Martin, A. R., & Clarke, M. R. (1986). The Diet of Sperm Whales (*Physeter macrocephalus*) Captured Between Iceland and Greenland. *Journal of the Marine Biological Association of the United Kingdom*, 66, 779-790.
- Mate, B. R., Lagerquist, B. A., Winsor, M. H., Geraci, J. R., & Prescott, J. H. (2005). Movements and dive habits of a satellite-monitored longfinned pilot whale (*Globicephala melas*) in the northwest Atlantic. *Marine Mammal Science*, 21(1), 136-144.
- Mate, B. R., & Nieukirk, S. (1992, 14-15 April 1992). *Satellite-Monitored Movements of Right Whales*. Paper presented at the The Right Whale in the Western North Atlantic: A Science and Management Workshop, Silver Spring, Maryland.
- Mate, B. R., Palacios, D. M., Baker, C. S., Lagerquist, B. A., Irvine, L. M., Follett, T., . . . Winsori, M. H. (2016). *Baleen (Blue & Fin) Whale Tagging and Analysis in Support of Marine Mammal Monitoring Across Multiple Navy Training Areas. Preliminary Summary*. (Contract No. N62470-15-8006, Monitoring report). p. 60.
- Mate, B. R., Stafford, K. M., Nawojchik, R., & Dunn, J. L. (1994). Movements and Dive Behavior of a Satellite-Monitored Atlantic White-Sided Dolphin (*Lagenorhynchus acutus*) in the Gulf of Maine. *Marine Mammal Science*, 10(1), 116-121.
- Maze-Foley, K., & Mullin, K. D. (2006). Cetaceans of the Oceanic Northern Gulf of Mexico: Distributions, Group Sizes and Interspecific Associations. *Journal of Cetacean Research and Management*, 8(2), 203-213.
- McAlpine, D. F., Murison, L. D., & Hoberg, E. P. (1997). New Records for the Pygmy Sperm Whale, *Kogia breviceps* (Physeteridae) from Atlantic Canada with Notes on Diet and Parasites. *Marine Mammal Science*, 13(4), 701-704.
- McClellan, C. M., Hudson, C. G., & Read, A. J. (2007). *Use of Oceanic Habitats by Loggerhead Sea Turtles (Caretta caretta)*. Paper presented at the Twenty-Seventh Annual Symposium on Sea Turtle Biology and Conservation, Myrtle Beach, SC.

- McGowan, A., Broderick, A. C., Frett, G., Gore, S., Hastings, M., Pickering, A., . . . Godley, B. J. (2008). Down But Not Out: Marine Turtles of the British Virgin Islands. *Animal Conservation*, *11*, 92-103.
- McMahon, C. R., Bradshaw, C. J. A., & Hays, G. C. (2007). Satellite Tracking Reveals Unusual Diving Characteristics for a Marine Reptile, the Olive Ridley Turtle *Lepidochelys olivacea*. *Marine Ecology Progress Series*, *329*, 239-252.
- Mead, J. G. (1989). Beaked Whales of the Genus -- *Mesoplodon*. In. Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals* (Vol. 4: River Dolphins and the Larger Toothed Whales, pp. 349-430). London, United Kingdom: Academic Press.
- Mead, J. G., & Potter, C. W. (1990). Natural History of Bottlenose Dolphins Along the Central Atlantic Coast of the United States. In. Leatherwood, S. & Reeves, R. R. (Eds.), *The Bottlenose Dolphin* (pp. 165-195). New York: Academic Press, Inc.
- Melin, S. R., & DeLong, R. L. (1999). *At-Sea Distribution and Diving Behavior of California Sea Lion Females from San Miguel Island, California*. Seattle, WA: Alaska Fisheries Science Center, National Marine Fisheries Service, National Marine Mammal Laboratory. pp. 407-412.
- Melin, S. R., DeLong, R. L., & Siniff, D. B. (2008). The Effects of El Nino on the Foraging Behavior of Lactating California Sea Lions (*Zalophus californianus californianus*) During the Nonbreeding Season. *Canadian Journal of Zoology*, *86*, 192-206.
- Mignucci-Giannoni, A. A., Toyos-González, G. M., Pérez-Padilla, J., Rodríguez-López, M. A., & Overing, J. (1999). Mass Stranding of Pygmy Killer Whales (*Feresa attenuata*) in the British Virgin Islands. *Journal of the Marine Biological Association of the United Kingdom*, *80*, 759-760.
- Miller, P., Antunes, R., Alves, A. C., Wensveen, P., Kvadsheim, P., Kleivane, L., . . . Tyack, P. (2011). *The 3S Experiments: Studying the Behavioural Effects of Naval Sonar on Killer Whales (*Orcinus orca*), Sperm Whales (*Physeter macrocephalus*), and Long-finned Pilot Whales (*Globicephala melas*) in Norwegian Waters*.
- Miller, P. J. O. M., Shapiro, A. D., & Deecke, V. B. (2010). The diving behavior of mammal-eating killer whales (*Orcinus orca*): variations with ecological not physical factors. *Canadian Journal of Zoology*, *88*, 1103-1112.
- Minamikawa, S., Iwasaki, T., & Kishiro, T. (2007). Diving Behaviour of a Baird's Beaked Whale, *Berardius bairdii*, in the Slope Water Region of the western North Pacific: First Dive Records using a Data Logger. *Fisheries Oceanography*, *16*(6), 573-577.
- Minamikawa, S., Iwasaki, T., Tanaka, Y., Ryono, A., Noji, S., Sato, H., . . . Kato, H. (2003, 17-21 March 2003). *Diurnal Pattern of Diving Behavior in Striped Dolphins, *Stenella coeruleoalba**. Paper presented at the International Symposium on Bio-logging Science, Tokyo, Japan.
- Mintzer, V. J., Gannon, D. P., Barros, N. B., & Read, A. J. (2008). Stomach contents of mass-stranded short-finned pilot whales (*Globicephala macrorhynchus*). *Marine Mammal Science*, *24*(2), 290-302.

- Miyazaki, N., Kuramochi, T., & Amano, M. (1991). Pacific White-sided Dolphins (*Lagenorhynchus obliquidens*) Off Northern Hokkaido. *Memoirs of the National Science Museum, Tokyo*, 24, 131-139.
- Mizroch, S. A., Rice, D. W., & Breiwick, J. M. (1984). The Sei Whale, *Balaenoptera borealis*. *Marine Fisheries Review*, 46(4), 25-29.
- Moore, J. C. (1972). More Skull Characters of the Beaked Whale *Indopacetus pacificus*. *Fieldiana: Zoology*, 62(1), 1-19.
- Morreale, S. J., Meylan, A. B., Sadove, S. S., & Standora, E. A. (1992). Annual Occurrence and Winter Mortality of Marine Turtles in New York Waters. *Journal of Herpetology*, 26(3), 301-308.
- Morton, A. (2000). Occurrence, Photo-Identification and Prey of Pacific White-Sided Dolphins (*Lagenorhynchus obliquidens*) in the Broughton Archipelago, Canada 1984-1998. *Marine Mammal Science*, 16(1), 80-93.
- Mullin, K. D., Hoggard, W., & Hansen, L. J. (2004). Abundance and Seasonal Occurrence of Cetaceans in Outer Continental Shelf and Slope Waters of the North-Central and Northwestern Gulf of Mexico. *Gulf of Mexico Science*, 22(1), 62 - 73.
- Musick, J. A., & Limpus, C. J. (1997). Habitat Utilization and Migration in Juvenile Sea Turtles. In Lutz, P. L. & Musick, J. A. (Eds.), *The Biology of Sea Turtles* (Vol. I, pp. 137-164). New York: CRC Press.
- Myers, A. E., & Hays, G. C. (2006). Do Leatherback Turtles *Dermochelys coriacea* Forage During the Breeding Season? A Combination of Data-Logging Devices Provide New Insights. *Marine Ecology Progress Series*, 322, 259-267.
- Naito, Y., Le Boeuf, B. J., Asaga, T., & Huntley, A. C. (1989). Long Term Diving Records of an Adult Female Northern Elephant Seal. *Antarctic Record*, 33(1), 1-9.
- Narazaki, T., Sato, K., & Miyazaki, N. (2006, 03-08 April 2006). *Fine-Scale Diving Behaviour of Migrating Turtles Revealed by Auto Releasing Logger System*. Paper presented at the Twenty-sixth Annual Symposium on Sea Turtle Biology and Conservation, Island of Crete, Greece.
- National Marine Fisheries Service. (2015, 16 January 2015). Bottlenose Dolphin (*Tursiops truncatus*). *Protected Resources-Species* Retrieved Retrieved 16 January 2015 from <http://www.nmfs.noaa.gov/pr/species/mammals/dolphins/bottlenose-dolphin.html> as accessed on 2 September 2016.
- National Marine Fisheries Service (NMFS). (1992a). *Cruise Report of the Harbor Porpoise Survey - 1992, AJ92-01*. Woods Hole, Massachusetts: National Marine Fisheries Service (NMFS). p. 13 pp.
- National Marine Fisheries Service (NMFS). (1992b). *Cruise Results, NOAA Ship Oregon II, Cruise No. 92-01 (198), 3 January-11 February 1992*. (Unpublished Cruise Report). Pascagoula, MS: National Marine Fisheries Service, Southeast Fisheries Science Center. p. 22.
- National Marine Fisheries Service (NMFS). (1995a). *Cruise Results, R/V Abel-J Cruise AJ 95-01, (Part I) Summer Marine Mammal Survey*. Woods Hole, Massachusetts: National Marine Fisheries Service (NMFS).

- National Marine Fisheries Service (NMFS). (1995b). *Cruise Results, R/V Abel-J Cruise No. AJ 95-01 (Part II), Summer Marine Mammal Survey*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 4.
- National Marine Fisheries Service (NMFS). (1995c). *Cruise Results, R/V Abel-J Cruise No. AJ 95-02 (Leg 2), Marine Mammal Abundance Survey*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 7.
- National Marine Fisheries Service (NMFS). (1995d). *Cruise Results, R/V Pelican, Cruise No. PE 95-01, Marine Mammal Abundance Survey, Leg 1*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 6.
- National Marine Fisheries Service (NMFS). (1995e). *Cruise Results, R/V Pelican, Cruise No. PE 95-02, Marine Mammal Survey*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 7.
- National Marine Fisheries Service (NMFS). (1997a). *Cruise Results, R/V Abel-J Cruise No. AJ 97-01, Sea Mount Marine Mammal Survey*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 16.
- National Marine Fisheries Service (NMFS). (1997b). *Cruise Results, R/V Delaware II, Cruise No. DEL 97-05, Marine Mammal Distribution Survey*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 34.
- National Marine Fisheries Service (NMFS). (1998a). *Cruise Results, NOAA Charter R/V Abel-J, Cruise Numbers AJ 98-01 & AJ 98-02, Pelagic Cetacean Abundance Survey-Legs I and II*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 11.
- National Marine Fisheries Service (NMFS). (1998b). *Cruise Results, NOAA Ship Relentless, Cruise RS 98-01 (3), Summer Atlantic Ocean Marine Mammal Survey*. (Unpublished Cruise Report). Pascagoula, MS: National Marine Fisheries Service, Southeast Fisheries Science Center. p. 20.
- National Marine Fisheries Service (NMFS). (1998c). *Cruise Results, R/V Delaware II, Cruise No. DEL 98-04, Mid-Atlantic Marine Mammal Distribution Survey*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 30.
- National Marine Fisheries Service (NMFS). (1999a). *Cruise Results, R/V Delaware II, Cruise No. DE 99-03, Monitoring the Impacts of the Harbor Porpoise Take Reduction Plan*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 7.
- National Marine Fisheries Service (NMFS). (1999b). *Cruise Results: Summer Atlantic Ocean Marine Mammal Survey: NOAA Ship Oregon II Cruise OT 99-05 (236)*. Pascagoula, MS: United States Department of Commerce (USDoC). p. 32.
- National Marine Fisheries Service (NMFS). (2002). *Cruise Results, NOAA Ship Delaware II, Cruise No. DE-02-06, Joint Deepwater Systematics and Marine Mammal Survey*. (Unpublished Cruise Report). Woods Hole, MA: National Marine Fisheries Service, Northeast Fisheries Science Center. p. 35.

- National Marine Fisheries Service (NMFS), & United States Fish and Wildlife Service (USFWS). (1991). *Recovery Plan for U.S. Population of Atlantic Green Turtle*. (National Marine Fisheries Service (NMFS)). Washington, DC. p. 52.
- National Marine Fisheries Service (NMFS), & United States Fish and Wildlife Service (USFWS). (1993). *Recovery Plan for Hawksbill Turtles in the U.S. Caribbean Sea, Atlantic Ocean, and Gulf of Mexico*. St Petersburg, FL: National Marine Fisheries Service (NMFS). p. 55.
- National Marine Fisheries Service (NMFS), & United States Fish and Wildlife Service (USFWS). (1998). *Recovery Plan for U.S. Pacific Populations of the Hawksbill Turtle (Eretmochelys imbricata)*. (National Marine Fisheries Service (NMFS)). Silver Spring, MD. p. 83.
- National marine Fisheries Service (NMFS), & United States Fish and Wildlife Service (USFWS). (2007a). *Green Sea Turtle (Chelonia mydas) 5-Year Review: Summary and Evaluation*. U.S. Department of Commerce. p. 105.
- National Marine Fisheries Service (NMFS), & United States Fish and Wildlife Service (USFWS). (2007b). *Hawksbill Sea Turtle (Eretmochelys imbricata) 5-Year Review: Summary and Evaluation*. Silver Spring, MD: National Marine Fisheries Service. p. 90.
- National Marine Fisheries Service (NMFS), & United States Fish and Wildlife Service (USFWS). (2007c). *Olive Ridely Sea Turtle (Lepidochelys olivacea) 5-Year Review: Summary and Evaluation*. Silver Spring, MD: National Marine Fisheries Service. p. 64.
- Naval Undersea Center (NUC) Code 14. (1971). *A Synopsis of Marine Animal Underwater Sounds in Eight Geographic Areas*. (Special Report ADA068875). San Diego, CA: Naval Undersea Research and Development Center. p. 97.
- Navy, D. o. t. (2012). *Commander Task Force 20, 4th, and 6th Fleet Navy marine species density database*. Norfolk, VA: Naval Facilities Engineering Command Atlantic.
- Nawojchik, R., Aubin, D. J. S., & Johnson, A. (2003). Movements and Dive Behavior of Two Stranded, Rehabilitated Long-Finned Pilot Whales (*Globicephala melas*) in the Northwest Atlantic. *Marine Mammal Science*, 19(1), 232-239.
- Nerini, M. (1984). A Review of Gray Whale Feeding Ecology. In: Jones, M. L., Swartz, S. L. & Leatherwood, S. (Eds.), *The Gray Whale (Eschrichtius robustus)* (pp. 423-448). Orlando, FL: Academic Press.
- Neumann, D. R., & Orams, M. B. (2003). Feeding Behaviours of Short-beaked Common Dolphins, *Delphinus delphis*, in New Zealand. *Aquatic Mammals*, 29(1), 137-149.
- New, L. F., Moretti, D. J., Hooker, S. K., Costa, D. P., & Simmons, S. E. (2013). Using energetic models to investigate the survival and reproduction of beaked whales (family Ziphiidae). *PLoS one*, 8(7), e68725.
- Nilssen, K. T., Haug, T., & Lindblom, C. (2001). Diet of Weaned Pups and Seasonal Variations in Body Condition of Juvenile Barents Sea Harp Seals *Phoca groenlandica*. *Marine Mammal Science*, 17(4), 926-936.
- Niño-Torres, C. A., Gallo-Reynoso, J. P., Galván-Magaña, F., Escobar-Briones, E., & Macko, S. A. (2006). Isotopic Analysis of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, AND $\delta^{34}\text{S}$ "A Feeding Tale" in Teeth of the Longbeaked Common Dolphin, *Delphinus capensis*. *Marine Mammal Science*, 22(4), 831-846.

- Noren, S. R., & Williams, T. M. (2000). Body Size and Skeletal Muscle Myoglobin of Cetaceans: Adaptations for Maximizing Dive Duration. *Comparative Biochemistry and Physiology Part A*, 126, 181-191.
- Nøttestad, L., Fernö, A., & Axelsen, B. E. (2002). Digging in the Deep: Killer Whales' Advanced Hunting Tactic. *Polar Biology*, 25, 939-941.
- Nottestad, L., Ferno, A., Mackinson, S., Pitcher, T., & Misund, O. A. (2002). How Whales Influence Herring School Dynamics in a Cold-Front Area of the Norwegian Sea. *ICES Journal of Marine Science*, 59(2), 393-400.
- Nowacek, D. P., & McGregor, A. E. (2010, April 15, 2010). *Right Whale Dive Data*. Personal communication via E-mail to Watwood, S. L.,
- Nowacek, S. M., Nowacek, D. P., Johnson, M. P., Shorter, K. A., Powell, J. A., & Wells, R. S. (2002). *Manatee Behavioral Responses to Vessel Approaches: Results of Digital Acoustic Data Logger Tagging of Manatees in Belize*. (Florida Fish and Wildlife Conservation Commission Contract 01078). Sarasota, FL: Mote Marine Laboratory: Center for Marine Mammal and Sea Turtle Research. p. 61.
- O'Shea, T. J., Rathbun, G. B., Bonde, R. K., Buergelt, C. D., & Odell, D. K. (1991). An Epizootic of Florida Manatees Associated with a Dinoflagellate Bloom. *Marine Mammal Science*, 7(2), 165-179.
- Odell, D. K., & McClune, K. M. (1999). False Killer Whale -- *Pseudorca crassidens* (Owen, 1846). In: Ridgway, S. H. & Harrison, S. R. (Eds.), *Handbook of Marine Mammals* (Vol. 6: The Second Book of Dolphins and the Porpoises, pp. 213-244). New York, NY: Academic Press.
- Ohizumi, H., Kuramochi, T., Amano, M., & Miyazaki, N. (2000). Prey Switching of Dall's Porpoise *Phocoenoides dalli* with Population Decline of Japanese Pilchard *Sardinops melanostictus* Around Hokkaido, Japan. *Marine Ecology Progress Series*, 200, 265-275.
- Ohizumi, H., Yoshioka, M., Mori, K., & Miyazaki, N. (1998). Stomach Contents of Common Dolphins (*Delphinus delphis*) in the Pelagic Western North Pacific. *Marine Mammal Science*, 14(4), 835-844.
- Øien, N. (1990). Sightings Surveys in the Northeast Atlantic in July 1988: Distribution and Abundance of Cetaceans. *Reports of the International Whaling Commission*, 40, 499-511.
- Olesiuk, P. F. (1993). Annual Prey Consumption by Harbor Seals (*Phoca vitulina*) in the Strait of Georgia, British Columbia. *Fishery Bulletin*, 91(3), 491-515.
- Oleson, E. M., Calambokidis, J., Burgess, W. C., McDonald, M. A., LeDuc, C. A., & Hildebrand, J. A. (2007). Behavioral Context of Call Production by Eastern North Pacific Blue Whales. *Marine Ecology Progress Series*, 330, 269-284.
- Omura, H. (1962). Further Information on Bryde's Whale from the Coast of Japan. *Scientific Reports of the Whales Research Institute*, 16, 7-18.
- Ostrom, P. H., Lien, J., & Macko, S. A. (1993). Evaluation of the Diet of Sowerby's Beaked Whale, *Mesoplodon bidens*, Based on Isotopic Comparisons Among Northwestern Atlantic Cetaceans. *Canadian Journal of Zoology*, 71, 858-861.

- Otani, S., Naito, Y., Kato, A., & Kawamura, A. (2000). Diving Behavior and Swimming Speed of a Free-Ranging Harbor Porpoise, *Phocoena phocoena*. *Marine Mammal Science*, 16(4), 811-814.
- Otani, S., Naito, Y., Kawamura, A., Kawasaki, M., Nishiwaki, S., & Kato, A. (1998). Diving Behavior and Performance of Harbor Porpoises, *Phocoena phocoena*, in Funka Bay, Hokkaido, Japan. *Marine Mammal Science*, 14(2), 209-220.
- Ottensmeyer, C. A., & Whitehead, H. (2003). Behavioural Evidence for Social Units in Long-Finned Pilot Whales. *Canadian Journal of Zoology*, 81, 1327-1338.
- Palka, D. (2012). Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2011 line transect survey.
- Palka, D., Read, A., & Potter, C. (1997). Summary of Knowledge of White-Sided Dolphins (*Lagenorhynchus acutus*) from US and Canadian Atlantic Waters. *Reports of the International Whaling Commission*, 47, 729-734.
- Panigada, S., Zanardelli, M., Canese, S., & Jahoda, M. (1999, 28 November - 03 December 1999). *Deep Diving Performances of Mediterranean Fin Whales*. Paper presented at the 13th Biennial Conference on the Biology of Marine Mammals, Wailea, HI.
- Parker, D. M., Dutton, P. H., Kopitsky, K., & Pitman, R. L. (2003). *Movement and Dive Behavior Determined by Satellite Telemetry for Male and Female Olive Ridley Turtles in the Eastern Tropical Pacific*. Paper presented at the Twenty-Second Annual Symposium on Sea Turtle Biology and Conservation.
- Parks, S. E., Warren, J. D., Stamieszkin, K., Mayo, C. A., & Wiley, D. (2011). Dangerous dining: surface foraging of North Atlantic right whales increases risk of vessel collisions. *Biol. Lett.* doi: 10.1098.
- Parrish, F. A., Abernathy, K., Marshall, G. J., & Buhleier, B. M. (2002). Hawaiian Monk Seals (*Monachus schauinslandi*) Foraging in Deep-water Coral Beds. *Marine Mammal Science*, 18(1), 244-258.
- Parrish, F. A., Craig, M. P., Ragen, T. J., Marshall, G. J., & Buhleier, B. M. (2000). Identifying Diurnal Foraging Habitat of Endangered Hawaiian Monk Seals Using a Seal-Mounted Video Camera. *Marine Mammal Science*, 16(2), 392-412.
- Parrish, F. A., Marshall, G. J., Littnan, C. L., Heithaus, M., Canja, S., Becker, B., . . . Antonelis, G. A. (2005). Foraging of Juvenile Monk Seals at French Frigate Shoals, Hawaii. *Marine Mammal Science*, 21(1), 93-107.
- Payne, P. M., & Heinemann, D. W. (1993). The Distribution of Pilot Whales (*Globicephala* spp.) in Shelf/Shelf-edge and Slope Waters of the Northeastern United States, 1978-1988. In: Donovan, G. P., Lockyer, C. H. & Martin, A. R. (Eds.), *Biology of Northern Hemisphere Pilot Whales* (Vol. Special Issue 14, pp. 50-68). Cambridge, UK: International Whaling Commission.
- Payne, P. M., & Selzer, L. A. (1989). The Distribution, Abundance, and Selected Prey of the Harbor Seal, *Phoca vitulina concolor*, in Southern New England. *Marine Mammal Science*, 5(2), 173-192.

- Pelletier, D., Roos, D., & Ciccione, S. (2003). Oceanic Survival and Movements of Wild and Captive-reared Immature Green turtles (*Chelonia mydas*) in the Indian Ocean. *Aquatic Living Resources*, 16, 35-41.
- Perrin, W. F., Caldwell, D. K., & Caldwell, M. C. (1994a). Atlantic Spotted Dolphin--*Stenella frontalis* (G. Cuvier, 1829). In. Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals* (Vol. 5: The First Book of Dolphins, pp. 173-190). New York, NY: Academic Press.
- Perrin, W. F., & Hohn, A. A. (1994). Pantropical Spotted Dolphin--*Stenella attenuata*. In. Ridgway, S. H. & Harrison, S. R. (Eds.), *Handbook of Marine Mammals* (Vol. 5: The First Book of Dolphins, pp. 71-98). New York, NY: Academic Press.
- Perrin, W. F., Mitchell, E. D., Mead, J. G., Caldwell, D. K., & Bree, P. J. H. v. (1981). *Stenella clymene*, a Rediscovered Tropical Dolphin of the Atlantic. *Journal of Mammalogy*, 62(3), 583-598.
- Perrin, W. F., Wilson, C. E., & Archer II, F. I. (1994b). Striped Dolphin--*Stenella coeruleoalba* (Meyen, 1833). In. Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals* (Vol. 5: The First Book of Dolphins, pp. 129-159). New York, NY: Academic Press.
- Perryman, W. L., Au, D. W. K., Leatherwood, S., & Jefferson, T. A. (1994). Melon-Headed Whale--*Peponocephala electra* Gray 1846. In. Ridgway, S. H. & Harrison, S. R. (Eds.), *Handbook of Marine Mammals* (Vol. 5: The First Book of Dolphins, pp. 363-386). New York, NY: Academic Press.
- Perryman, W. L., & Foster, T. C. (1980). *Preliminary Report on Predation by Small Whales, Mainly the False Killer Whale, Pseudorca crassidens, on Dolphins (Stenella spp. and Delphinus delphis) in the Eastern Tropical Pacific*. (Administrative Report LJ-80-05). La Jolla, CA: National Marine Fisheries Service, Southeast Fisheries Science Center. p. 10.
- Pitman, R. L. (1990, 20-24 February 1990). *Pelagic Distribution and Biology of Sea Turtles in the Eastern Tropical Pacific*. Paper presented at the Tenth Annual Workshop on Sea Turtle Biology and Conservation, Hilton Head Island, SC.
- Pitman, R. L. (1992, 26 February - 02 March 1991). *Sea Turtle Associations with Flotsam in the Eastern Tropical Pacific*. Paper presented at the Eleventh Annual Workshop on Sea Turtle Biology and Conservation, Jekyll Island, Georgia.
- Pitman, R. L. (2002a). Indo-Pacific Beaked Whale. In. Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 615-616). San Diego, CA: Academic Press.
- Pitman, R. L. (2002b). Mesoplodont Whales. In. Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 738-742). San Diego, CA: Academic Press.
- Pitman, R. L., & Ensor, P. (2003). Three Forms of Killer Whales (*Orcinus orca*) in Antarctic Waters. *Journal of Cetacean Research and Management*, 5(2), 131-139.
- Pitman, R. L., Palacios, D. M., Brennan, P. L. R., Brennan, B. J., Balcomb III, K. C., & Miyashita, T. (1999). Sightings and Possible Identity of a Bottlenose Whale in the Tropical Indo-Pacific: *Indopacetus pacificus*? *Marine Mammal Science*, 15(2), 531-549.

- Pitman, R. L., & Stinchcomb, C. (2002). Rough-Toothed Dolphins (*Steno bredanensis*) as Predators of Mahimahi (*Coryphaena hippurus*). *Pacific Science*, 56(4), 447-450.
- Plön, S. (2004). *The Status and Natural History of Pygmy (Kogia breviceps) and Dwarf (K. sima) Sperm Whales off Southern Africa*. PhD in Zoology (Ph.D Dissertation), Rhodes University, Grahamstown, South Africa.
- Polovina, J. J., Balazs, G. H., Howell, E. A., Parker, D. M., Seki, M. P., & Dutton, P. H. (2004). Forage and Migration Habitat of Loggerhead (*Caretta caretta*) and Olive Ridley (*Lepidochelys olivacea*) Sea Turtles in the Central North Pacific Ocean. *Fisheries Oceanography*, 13(1), 36-51.
- Polovina, J. J., Howell, E., Parker, D. M., & Balazs, G. H. (2003). Dive-Depth Distribution of Loggerhead (*Caretta caretta*) and Olive Ridley (*Lepidochelys olivacea*) Sea Turtles in the Central North Pacific: Might Deep Longling Sets Catch Fewer Turtles? *Fishery Bulletin*, 101(1), 189-193.
- Ponganis, P. J., Gentry, R. L., Ponganis, E. P., & Ponganis, K. V. (1992). Analysis of Swim Velocities During Deep and Shallow Dives of Two Northern Fur Seals *Callorhinus ursinus*. *Marine Mammal Science*, 8(1), 69-75.
- Porras-Peters, H., Aurioles-Gamboa, D., Cruz-Escalona, V. H., & Koch, P. L. (2008). Trophic Level and Overlap of Sea Lions (*Zalophus californianus*) in the Gulf of California, Mexico. *Marine Mammal Science*, 24(3), 554-576.
- Pusineri, C., Magnin, V., Meynier, L., Spitz, J., Hassani, S., & Ridoux, V. (2007). Food and Feeding Ecology of the Common Dolphin (*Delphinus delphis*) in the Oceanic Northeast Atlantic and Comparison with its Diet in Neritic Areas. *Marine Mammal Science*, 23(1), 30-47.
- Rankin, S., Oswald, J., Barlow, J., & Lammers, M. (2007). Patterned Burst-Pulse Vocalizations of the Northern Right Whale Dolphin, *Lissodelphis borealis*. *Journal of the Acoustical Society of America*, 121(12), 1213-1218.
- Ream, R. R., Sterling, J. T., & Loughlin, T. R. (2005). Oceanographic Features Related to Northern Fur Seal Migratory Movements. *Deep Sea Research II*, 52, 823-843.
- Recchia, C. A., & Read, A. J. (1989). Stomach Contents of Harbour Porpoises, *Phocoena phocoena* (L.), from the Bay of Fundy. *Canadian Journal of Zoology*, 67, 2140-2146.
- Reeves, R. R., Mitchell, E., & Whitehead, H. (1993). Status of the Northern Bottlenose Whale, *Hyperoodon ampullatus*. *Canadian Field-Naturalist*, 107(4), 490-508.
- Reeves, R. R., Smeenk, C., Kinze, C. C., Brownell Jr., R. L., & Lien, J. (1999). White-beaked Dolphin -- *Lagenorhynchus albirostris* Gray, 1846. In: Ridgway, S. H. & Harrison, S. R. (Eds.), *Handbook of Marine Mammals* (Vol. 6: The Second Book of Dolphins and the Porpoises, pp. 1-30). San Diego, CA: Academic Press.
- Reeves, R. R., Stewart, B. S., Clapham, P. J., & Powell, J. A. (2002). *Guide to Marine Mammals of the World*. New York, NY: Chanticleer Press Inc.
- Reilly, S. B. (1990). Seasonal Changes in Distribution and Habitat Differences Among Dolphins in the Eastern Tropical Pacific. *Marine Ecology Progress Series*, 66, 1-11.

- Renaud, M. L. (1995). Movements and Submergence Patterns of Kemp's Ridley Turtles (*Lepidochelys kempii*). *Journal of Herpetology*, 29(3), 370-374.
- Renaud, M. L., & Carpenter, J. A. (1994). Movements and Submergence Patterns of Loggerhead Turtles (*Caretta caretta*) in the Gulf of Mexico Determined through Satellite Telemetry. *Bulletin of Marine Science*, 55(1), 1-15.
- Renaud, M. L., & Williams, J. A. (1997). *Movements of Kemp's Ridley (Lepidochelys kempii) and Green (Chelonia mydas) Sea Turtles Using Lavaca Bay and Matagorda Bay, 1996 - 1997*. Dallas, TX: Environmental Protection Agency Office of Planning and Coordination. p. 62.
- Renaud, M. L., & Williams, J. A. (2005). Kemp's Ridley Sea Turtle Movements and Migrations. *Chelonian Conservation and Biology*, 4(4), 808-816.
- Reynolds III, J. E., & Powell, J. A. (2002). Manatees. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 709-720). San Diego, CA: Academic Press.
- Rice, M. R., & Balazs, G. H. (2008). Diving Behavior of the Hawaiian Green Turtle (*Chelonia mydas*) during Oceanic Migrations. *Journal of Experimental Marine Biology and Ecology*, 356(1-2), 121-127.
- Riedman, M. (1990). *The Pinnipeds: Seals, Sea Lions, and Walruses*. Berkley, CA: University of California Press.
- Riedman, M. L., & Estes, J. A. (1990). *The Sea Otter (Enhydra lutris): Behavior, Ecology, and Natural History*. (Biological Report 90(14)). Washington, DC: United States Department of the Interior. p. 33.
- Ringelstein, J., Pusineri, C., Hassani, S., Meynier, L., Nicolas, R., & Ridoux, V. (2006). Food and Feeding Ecology of the Striped Dolphin, *Stenella coeruleoalba*, in the Oceanic Waters of the North-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom*, 86, 909-918.
- Ritter, F. (2001, 6-10 May 2001). *21 Cetacean Species off La Gomera (Canary Islands): Possible Reasons for an Extraordinary Species Diversity*. Paper presented at the Proceedings of the 15th Annual Conference of the European Cetacean Society, Rome, Italy.
- Ritter, F., & Brederlau, B. (1999). Behavioural Observations of Dense Beaked Whales (*Mesoplodon densirostris*) off La Gomera, Canary Islands (1995-1997). *Aquatic Mammals*, 25(2), 55-61.
- Robertson, K. M., & Chivers, S. J. (1997). Prey Occurrence in Pantropical Spotted Dolphins, *Stenella attenuata*, from the Eastern Tropical Pacific. *Fishery Bulletin*, 95, 334-348.
- Robinson, P. W., Costa, D. P., Crocker, D. E., Gallo-Reynoso, J. P., Champagne, C. D., Fowler, M. A., . . . Yoda, K. (2012). Foraging Behavior and Success of a Mesopelagic Predator in the Northeast Pacific Ocean: Insights from a Data-Rich Species, the Northern Elephant Seal. *PloS one*, 7(5). doi: 10.1371/journal.pone.0036728.
- Robison, B. H., & Craddock, J. E. (1983). Mesopelagic Fishes Eaten by Fraser's Dolphin, *Lagenodelphis hosei*. *Fishery Bulletin*, 81(2), 283-289.

- Rosel, P. E., Dizon, A. E., & Heyning, J. E. (1994). Genetic Analysis of Sympatric Morphotypes of Common Dolphins (genus *Delphinus*). *Marine Biology*(119), 159-167.
- Ross, G. J. B. (1979). Records of Pygmy and Dwarf Sperm Whales, genus *Kogia*, from Southern Africa, with Biological Notes and Some Comparisons. *Annals of the Cape Provincial Museums (Natural History)*, 11(14), 259-327.
- Roszbach, K. A., & Herzing, D. L. (1997). Underwater Observations of Benthic-Feeding Bottlenose Dolphins (*Tursiops truncatus*) Near Grand Bahama Island, Bahamas. *Marine Mammal Science*, 13(3), 498-504.
- Ruchonnet, D., Boutoute, M., Guinet, C., & Mayzaud, P. (2006). Fatty Acid Composition of Mediterranean Fin Whale *Balaenoptera physalus* Blubber with Respect to Body Heterogeneity and Trophic Interaction. *Marine Ecology Progress Series*, 311, 165-174.
- Sakamoto, W., Uchida, I., Naito, Y., Kureha, K., Tujimura, M., & Sato, K. (1990). Deep Diving Behavior of the Loggerhead Turtle Near the Frontal Zone. *Nippon Suisan Gakkaishi*, 56(9), 1435-1443.
- Sale, A., Luschi, P., Mencacci, R., Lambardi, P., Hughes, G. R., Hays, G. C., . . . Papi, F. (2006). Long-Term Monitoring of Leatherback Turtle Diving Behaviour During Oceanic Movements. *Journal of Experimental Marine Biology and Ecology*, 328, 197-210.
- Santos, M. B., Pierce, G. J., Herman, J., Lopez, A., Guerra, A., Mente, E., & Clarke, M. R. (2001a). Feeding Ecology of Cuvier's Beaked Whale (*Ziphius cavirostris*): A Review with New Information on the Diet of this Species. *Journal of the Marine Biological Association of the United Kingdom*, 81, 687-694.
- Santos, M. B., Pierce, G. J., Lopez, A., Reid, R. J., Ridoux, V., & Mente, E. (2006). Pygmy Sperm Whales *Kogia breviceps* in the Northeast Atlantic: New Information on Stomach Contents and Strandings. *Marine Mammal Science*, 22(3), 600-616.
- Santos, M. B., Pierce, G. J., Smeenk, C., Addink, M. J., Kinze, C. C., Tougaard, S., & Herman, J. (2001b). Stomach Contents of Northern Bottlenose Whales *Hyperoodon ampullatus* Stranded in the North Sea. *Journal of the Marine Biological Association of the United Kingdom*, 81(1), 143-150.
- Sasaki, T., Nikaido, M., Hamilton, H., Goto, M., Kato, H., Kanda, N., . . . Okada, N. (2005). Mitochondrial Phylogenetics and Evolution of Mysticete Whales. *Systematic Biology*, 54(1), 77-90.
- Sasaki, T., Nikaido, M., Wada, S., Yamada, T. K., Cao, Y., Hasegawa, M., & Okada, N. (2006). *Balaenoptera omurai* is a Newly Discovered Baleen Whale that Represents an Ancient Evolutionary Lineage. *Molecular Phylogenetics and Evolution*, 41, 40-52.
- Sasso, C., & Witzell, W. N. (2006). Diving Behavior of an Immature Kemp's Ridley Turtle (*Lepidochelys kempii*) from Gullivan Bay, Ten Thousands Islands, South-west Florida. *Journal of the Marine Biological Association of the United Kingdom*, 86, 915-925.
- Saulitis, E., Matkin, C., Barrett-Lennard, L., Heise, K., & Ellis, G. (2000). Foraging Strategies of Sympatric Killer Whale (*Orcinus orca*) Populations in Prince William Sound, Alaska. *Marine Mammal Science*, 16(1), 94-109.

- Schlexer, F. V. (1984). *Diving Patterns of the Hawaiian Monk Seal, Lisianski Island, 1982*. (NOAA-TM-NMFS-SWFC-41). Honolulu, HI: National Marine Fisheries Service, NOAA. p. 7.
- Schoenherr, J. R. (1991). Blue whales Feeding on High Concentrations of Euphausiids Around Monterey Submarine Canyon. *Canadian Journal of Zoology*, 69, 583-594.
- Schorr, G. S., Falcone, E. A., Moretti, D. J., & Andrews, R. D. (2014). First Long-Term Behavioral Records from Cuvier's Beaked Whales (*Ziphius cavirostris*) Reveal Record-Breaking Dives. *PLOS ONE*, 9(3), e92633.
- Schreer, J. F., & Kovacs, K. M. (1997). Allometry of Diving Capacity in Air-Breathing Vertebrates. *Canadian Journal of Zoology*, 75, 339-358.
- Schroeder, B. A., Foley, A. M., & Bagley, D. A. (2003). Nesting Patterns, Reproductive Migrations, and Adult Foraging Areas of Loggerhead Turtles. In. Bolten, A. B. & Witherington, B. E. (Eds.), *Loggerhead Sea Turtles* (pp. 114-124). Washington, D.C.: Smithsonian Institution Press.
- Scott, M. D., & Chivers, S. J. (2009). Movements and Diving Behavior of Pelagic Spotted Dolphins. *Marine Mammal Science*, 25(1), 137-160.
- Scott, M. D., Hohn, A. A., Westgate, A. J., Nicolas, J. R., Whitaker, B. R., & Campbell, W. J. (2001). A note on the release and tracking of a rehabilitated pygmy sperm whale (*Kogia breviceps*). *Journal of Cetacean Research and Management*, 3(1), 87-94.
- Sears, R. (2002). Blue Whale. In. Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 112-116). San Diego, CA: Academic Press.
- Sears, R., & Perrin, W. F. (2008). Blue Whale, 2nd. In *Encyclopedia of Marine Mammals* (pp. 120-124). San Diego, CA: Academic Press.
- Sekiguchi, K., Klages, N. T. W., & Best, P. B. (1992). Comparative Analysis of the Diets of Smaller Odontocete Cetaceans Along the Coast of Southern Africa. *South African Journal of Marine Science*, 12, 843-861.
- Selzer, L. A., & Payne, P. M. (1988). The Distribution of White-Sided (*Lagenorhynchus acutus*) and Common Dolphins (*Delphinus delphis*) vs. Environmental Features of the Continental Shelf of the Northeastern United States. *Marine Mammal Science*, 4(2), 141-153.
- Seney, E. E., & Landry, A. M. (2008). Movements of Kemp's Ridley Sea Turtles Nesting on the Upper Texas Coast: Implications for Management. *Endangered Species Research*, 4, 73-84.
- Seney, E. E., & Musick, J. A. (2005). Diet Analysis of Kemp's Ridley Sea Turtles (*Lepidochelys kempii*) in Virginia. *Chelonian Conservation and Biology*, 4(4), 864-871.
- Shane, S. H. (1990). Behavior and Ecology of the Bottlenose Dolphin at Sanibel Island, Florida. In. Leatherwood, S. & Reeves, R. R. (Eds.), *The Bottlenose Dolphin* (pp. 245-265). New York, NY: Academic Press, Inc.
- Shapiro, A. D. (2008). *Orchestration: The Movement and Vocal Behavior of Free-Ranging Norwegian Killer Whales (Orcinus orca)*. Doctor of Philosophy (Doctrine), Massachusetts

- institute of Technology and the Woods Hole Oceanographic Institution, Woods Hole, MA.
- Shaver, D. J., & Rubio, C. (2007). Post-Nesting Movement of Wild and Head-Started Kemp's Ridley Sea Turtles *Lepidochelys kempii* in the Gulf of Mexico. *Endangered Species Research*, 3, 1-13.
- Shaver, D. J., Schroeder, B. A., Byles, R. A., Burchfield, P. M., Pena, J., Marquez, R., & Martinez, H. J. (2005). Movements and Home Ranges of Adult Male Kemp's Ridley Sea Turtles (*Lepidochelys kempii*) in the Gulf of Mexico Investigated by Satellite Telemetry. *Chelonian Conservation and Biology*, 4(4), 817-827.
- Siciliano, S., Santos, M. C. d. O., Vicente, A. F. C., Alvarenga, F. S., Zampirolli, E., Brito Jr., J. L., . . . Pizzorno, J. L. A. (2004). Strandings and Feeding Records of Bryde's Whales (*Balaenoptera edeni*) in South-Eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom*, 84, 857-859.
- Simard, P., & Gowans, S. (2008). Group Movements of White-Beaked Dolphins (*Lagenorhynchus albirostris*) near Halifax, Canada. *Aquatic Mammals*, 34(3), 331-337.
- Simard, P., Lawlor, J. L., & Gowans, S. (2006). Temporal Variability of Cetaceans near Halifax, Nova Scotia. *Canadian Field-Naturalist*, 120(1), 93-99.
- Sinclair, E. H. (1994). Prey of Juvenile Northern Elephant Seals (*Mirounga angustirostris*) in the Southern California Bight. *Marine Mammal Science*, 10(2), 230-239.
- Sivle, L. D., Kvadsheim, P. H., Fahlman, A., Lam, F. P. A., Tyack, P. L., & Miller, P. J. O. (2012). Changes in dive behavior during naval sonar exposure in killer whales, long-finned pilot whales, and sperm whales. *Frontiers in Physiology*, 3(Article 400), 1-11. doi: 10.3389/fphys.2012.00400.
- Smith, R. J., & Read, A. J. (1992). Consumption of Euphausiids by Harbour Porpoise (*Phocoena phocoena*) Calves in the Bay of Fundy. *Canadian Journal of Zoology*, 70, 1629-1632.
- Soto, N. A., Johnson, M. P., Madsen, P. T., Díaz, F., Domínguez, I., Brito, A., & Tyack, P. (2008). Cheetahs of the Deep Sea: Deep Foraging Sprints in Short-Finned Pilot Whales off Tenerife (Canary Islands). *Journal of Animal Ecology*, 77(5), 936-947.
- Southall, B., Calambokidis, J., Barlow, J., Moretti, D., Friedlaender, A., Stimpert, A., . . . Schorr, G. (2014). *Biological and behavioral response studies of marine mammals in southern California, 2013 ("SOCAL-13"), final project report*. Southall Environmental Associates, Inc. p. 54.
- Spotila, J. R. (2004). *Sea Turtles: A Complete Guide to Their Biology, Behavior, and Conservation*. Baltimore, MD: The Johns Hopkins University Press.
- Stacey, P. J., Leatherwood, S., & Baird, R. W. (1994). *Pseudorca crassidens*. *Mammalian Species*, 456, 1-6.
- Starbird, C. H., Hillis-Starr, Z.-M., Harvey, J. T., & Eckert, S. A. (1999). Internesting Movements and Behavior of Hawksbill Turtles (*Eretmochelys imbricata*) Around Buck Island Reef National Monument, St. Croix, U.S. Virgin Islands. *Chelonian Conservation and Biology*, 3(2), 237-243.

- Stelle, L. L., Megill, W. M., & Kinzel, M. R. (2008). Activity Budget and Diving Behavior of Gray Whales (*Eschrichtius robustus*) in Feeding Grounds Off Coastal British Columbia. *Marine Mammal Science*, 24(3), 462-478.
- Sterling, J. T., & Ream, R. R. (2004). At-Sea Behavior of Juvenile Male Northern Fur Seals (*Callorhinus ursinus*). *Canadian Journal of Zoology*, 82(10), 1621-1637.
- Stewart, B. S., Antonelis, G. A., Baker, J. D., & Yochem, P. K. (2006). Foraging Biogeography of Hawaiian Monk Seals in the Northwestern Hawaiian Islands. *Atoll Research Bulletin*, 543, 131-145.
- Stewart, B. S., Harvey, J., & Yochem, P. K. (2001). Post-Release Monitoring and Tracking of a Rehabilitated California Gray Whale. *Aquatic Mammals*, 27(3), 294-300.
- Stewart, B. S., & Huber, H. R. (1993). *Mirounga angustirostris*. *Mammalian Species*(1-10).
- Stewart, B. S., & Yochem, P. K. (2004a). *Use of Marine Habitats by Hawaiian Monk Seals (Monachus schauinslandi) from Kure Atoll: Satellite-Linked Monitoring in 2001-2002*. (Administrative Report H-04-01C). San Diego, CA: Pacific Islands Fisheries Science Center. p. 113.
- Stewart, B. S., & Yochem, P. K. (2004b). *Use of Marine Habitats by Hawaiian Monk Seals (Monachus schauinslandi) from Laysan Island: Satellite-Linked Monitoring in 2001-2002*. (Administrative Report H-04-02C). San Diego, CA: Hubbs-SeaWorld Research Institute. p. 131.
- Stewart, B. S., Yochem, P. K., DeLong, R. L., & Antonelis, G. A. (1987). *Interactions between Guadalupe fur seals and California sea lions at San Nicolas and San Miguel Islands, California*. Paper presented at the Status, Biology and Ecology of Fur Seals, Cambridge, England.
- Stimpert, A. K., DeRuiter, S., Southall, B., Moretti, D., Falcone, E., Goldbogen, J., . . . Calambokidis, J. (2014). Acoustic and foraging behavior of a Baird's beaked whale, *Berardius bairdii*, exposed to simulated sonar. *Scientific Reports*, 4. doi: 10.1038/srep07031.
- Stockin, K. A., Fairbairns, R. S., Parsons, E. C. M., & Sims, D. W. (2001). Effects of Diel and Seasonal Cycles on the Dive Duration of the Minke Whale (*Balaenoptera acutorostrata*). *Journal of the Marine Biological Association of the United Kingdom*, 81(1), 189-190.
- Storch, S., Wilson, R. P., Hillis-Starr, Z.-M., & Adelung, D. (2005). Cold-Blooded Divers: Temperature-Dependent Dive Performance in the Wild Hawksbill Turtle *Eretmochelys imbricata*. *Marine Ecology Progress Series*, 293, 263-271.
- Stroud, R. K., Fiscus, C. H., & Kajimura, H. (1981). Food of the Pacific White-Sided Dolphin, *Lagenorhynchus obliquidens*, Dall's Porpoise, *Phocoenoides dalli*, and Northern Fur Seal, *Callorhinus ursinus*, off California and Washington. *Fishery Bulletin*, 78(4), 951-959.
- Swartz, S. L. (1986). Gray Whale Migratory, Social and Breeding Behavior. In. Donovan, G. P. (Ed.), *Behaviour of Whales in Relation to Management: Incorporating the Proceedings of a Workshop held in Seattle, WA, 19-23 April 1982* (Vol. Special Issue 8, pp. 207-229). Cambridge: International Whaling Commission.

- Swimmer, Y., Arauz, R., McCracken, M., McNaughton, L., Ballesteros, J., Musyl, M., . . . Brill, R. (2006). Diving Behavior and Delayed Mortality of Olive Ridley Sea Turtles *Lepidochelys olivacea* After Their Release from Longline Fishing Gear. *Marine Ecology Progress Series*, 323, 253-261.
- Teilmann, J., Christiansen, C. T., Kjellerup, S., Dietz, R., & Nachman, G. (2013). Geographic, seasonal, and diurnal surface behavior of harbor porpoises *Marine Mammal Science*, 29(2), E60-E76. doi: 10.1111/j.1748-7692.2012.00597.x.
- Teloni, V., Mark, J. P., Patrick, M. J. O., & Peter, M. T. (2008). Shallow Food for Deep Divers: Dynamic Foraging Behavior of Male Sperm Whales in a High Latitude Habitat. *Journal of Experimental Marine Biology and Ecology*, 354, 119-131.
- Thomas, K., Harvey, J. T., Goldstein, T., Barakos, J., & Gulland, F. (2010). Movement, dive behavior, and survival of California sea lions (*Zalophus californianus*) posttreatment for domoic acid toxicosis. *Marine Mammal Science*, 26(1), 36-52.
- Thompson, D., Hammond, P. S., Nicholas, K. S., & Fedak, M. A. (1991). Movements, Diving and Foraging Behaviour of Grey Seals (*Halichoerus grypus*). *Journal of Zoology, London*, 224, 223-232.
- Threlfall, W. (1978). First Record of the Atlantic Leatherback Turtle (*Dermochelys coriacea*) from Labrador. *Canadian Field-Naturalist*, 92(3), 287.
- Tove, M. (1995). Live Sighting of *Mesoplodon cf. M. mirus*, True's Beaked Whale. *Marine Mammal Science*, 11(1), 80-85.
- Tyack, P. (2010). *Mesoplodon denirostris* Andros Island, Bahamas 10/23/2006 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 23 October 2006 from <http://hdl.handle.net/1912/3650>.
- Tyack, P. L., Johnson, M., Soto, N. A., Sturlese, A., & Madsen, P. T. (2006). Extreme Diving of Beaked Whales. *The Journal of Experimental Biology*, 209, 4238-4253.
- Tyne, J. A., Pollock, K. H., Johnston, D. W., & Bejder, L. (2014). Abundance and survival rates of the Hawai'i island associated spinner dolphin (*Stenella longirostris*) stock. *PLoS One*, 9(1), e86132.
- U.S. Department of the Navy. (In Prep.-a). *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing*.
- U.S. Department of the Navy. (In Prep.-b). *U.S. Navy Marine Species Density Database Phase III for the Atlantic Fleet Training and Testing Study Area*. Norfolk, VA: Naval Facilities Engineering Command Atlantic. p. 273.
- U.S. Department of the Navy. (In Prep.-c). *U.S. Navy Marine Species Density Database Phase III for the Hawaii-Southern California Training and Testing (HSTT) Study Area*. Pearl Harbor, HI: Naval Facilities Engineering Command Pacific.
- van Dam, R. P., & Diez, C. E. (1996). Diving Behavior of Immature Hawksbills (*Eretmochelys imbricata*) in Caribbean Cliff-Wall Habitat. *Marine Biology*, 127, 171-178.

- Vikingsson, G. A. (1997). Feeding of Fin Whales (*Balaenoptera physalus*) off Iceland - Diurnal and Seasonal Variation and Possible Rates. *Journal of Northwest Atlantic Fishery Science*, 22, 77-89.
- Waerebeek, K. V., & Würsig, B. (2002). Pacific White-Sided Dolphin and Dusky Dolphin. In Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 859-861). San Diego, CA: Academic Press.
- Walker, W. A. (1996). Summer Feeding Habits of Dall's Porpoise, *Phocoenoides dalli*, in the Southern Sea of Okhotsk. *Marine Mammal Science*, 12(2), 167-182.
- Walker, W. A., Hanson, M. B., Baird, R. W., & Guenther, T. J. (1998). Food Habits of the Harbor Porpoise, *Phocoena phocoena*, and Dall's Porpoise, *Phocoenoides dalli*, in the Inland Waters of British Columbia and Washington. In Alaska Fisheries Science Center, A. (Ed.), *Marine Mammal Protection Act and Endangered Species Act Implementation Program 1997, AFSC Processed Report 98-10* (pp. 14): National Marine Fisheries Service.
- Walker, W. A., & Jones, L. L. (1994). Food Habits of Northern Right Whale Dolphin, Pacific White-Sided Dolphin, and Northern Fur Seal Caught in the High Seas Driftnet Fisheries of the North Pacific Ocean, 1990. *International North Pacific Fisheries Bulletin*, 53(2), 285-295.
- Walker, W. A., Mead, J. G., & Brownell Jr., R. L. (2002). Diets of Baird's Beaked Whales, *Berardius bairdii*, in the Southern Sea of Okhotsk and off the Pacific Coast of Honshu, Japan. *Marine Mammal Science*, 18(4), 902-919.
- Wallace, B. P., Tiwari, M., & Girondot, M. (2013). *Dermochelys coriacea*. . *The IUCN Red List of Threatened Species*. Retrieved from www.iucnredlist.org
- Wang, M.-C., Walker, W. A., Shao, K.-T., & Chou, L.-S. (2002). Comparative Analysis of the Diets of Pygmy Sperm Whales and Dwarf Sperm Whales in Taiwanese Waters. *Acta Zoologica Taiwanica*, 13(2), 53-62.
- Wang, M.-C., Walker, W. A., Shao, K.-T., & Chou, L.-S. (2003). Feeding Habits of the Pantropical Spotted Dolphin, *Stenella attenuata*, off the Eastern Coast of Taiwan. *Zoological Studies*, 42(2), 368-378.
- Waring, G. T., Josephson, E., Fairfield, C. P., & Maze-Foley, K. (2006). *U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments 2005*. Woods Hole, MA: U.S. Department of Commerce National Oceanic and Atmospheric Administration National Marine Fisheries Service. p. 358.
- Waring, G. T., Josephson, E., Maze-Foley, K., & Patricia, R. E. (2016). *U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments 2015*. Woods Hole, MA: U.S. Department of Commerce National Oceanic and Atmospheric Administration National Marine Fisheries Service. p. 512.
- Wathne, J. A., Haug, T., & Lydersen, C. (2000). Prey Preference and Niche Overlap of Ringed Seals *Phoca hispida* and Harp Seals *P. groenlandica* in the Barents Sea. *Marine Ecology Progress Series*, 194, 233-239.
- Watkins, W. A., Daher, M. A., Frstrup, K., & Sciara, G. N. d. (1994). Fishing and Acoustic Behavior of Fraser's Dolphin (*Lagenodelphis hosei*) near Dominica, Southeast Caribbean. *Caribbean Journal of Science*, 30(1-2), 76-82.

- Watkins, W. A., Daher, M. A., Samuels, A., & Gannon, D. P. (1997). Observations of *Peponocephala electra*, the Melon-Headed Whale, in the Southeastern Caribbean. *Caribbean Journal of Science*, 33(1-2), 34-40.
- Watkins, W. A., Tyack, P., Moore, K. E., & Notarbartolo-di-Sciara, G. (1987). *Steno bredanensis* in the Mediterranean Sea. *Marine Mammal Science*, 3(1), 78-82.
- Watwood, S. L., Miller, P. J. O., Johnson, M., Madsen, P. T., & Tyack, P. L. (2006). Deep-Diving Foraging Behaviour of Sperm Whales (*Physeter macrocephalus*). *Journal of Animal Ecology*, 75, 814-825.
- Weinrich, M. T., Belt, C. R., & Morin, D. (2001). Behavior and Ecology of the Atlantic White-Sided Dolphin (*Lagenorhynchus acutus*) in Coastal New England Waters. *Marine Mammal Science*, 17(2), 231-248.
- Weir, C. R. (2007). Occurrence and Distribution of Cetaceans off Northern Angola, 2004/05. *Journal of Cetacean Research and Management*, 9(3), 225-239.
- Weir, C. R., Debrah, J., Ofori-Danson, P. K., Pierpoint, C., & Waerebeek, K. V. (2008). Records of Fraser's Dolphin *Lagenodelphis hosei* Fraser 1956 from the Gulf of Guinea and Angola. *African Journal of Marine Science*, 30(2), 241-246.
- Weise, M. J. (2006). *Foraging Ecology of Male California Sea Lion (Zalophus californianus): Movement, Diving and Foraging Behavior, and Diving Capacity*. Doctor of Philosophy (Doctrine), University of California, Santa Cruz, CA.
- Wells, R. S., Boness, D. J., & Rathbun, G. B. (1999). Behavior. In: Reynolds III, J. E. & Rommel, S. A. (Eds.), *Biology of Marine Mammals* (pp. 324-422). Washington, D.C.: Smithsonian Institution Press.
- Wells, R. S., Early, G. A., Gannon, J. G., Lingenfelter, R. G., & Sweeney, P. (2008). *Tagging and Tracking of Rough-Toothed Dolphins (Steno bredanensis) from the March 2005 Mass Stranding in the Florida Keys*. (NOAA Technical Memorandum NMFS-SEFSC-574). Miami, FL: United States Department of Commerce, National Oceanic and Atmospheric Administration, and National Marine Fisheries Service. p. 50.
- Wells, R. S., Fougères, E. M., Cooper, A. G., Stevens, R. O., Brodsky, M., Lingenfelter, R. G., . . . Douglas, D. C. (2013). Movements and dive patterns of short-finned pilot whales (*Globicephala macrorhynchus*) released from a mass stranding in the Florida Keys. *Aquatic Mammals*, 39(1), 61-72.
- Wells, R. S., Manire, C. A., Byrd, L., Smith, D. R., Gannon, J. G., Fauquier, D., & Mullin, K. D. (2009). Movements and Dive Patterns of a Rehabilitated Risso's Dolphin, *Grampus griseus*, in the Gulf of Mexico and the Atlantic Ocean. *Marine Mammal Science*, 25(2), 420-429.
- Wells, R. S., & Scott, M. D. (1999). Bottlenose Dolphin - *Tursiops truncatus* (Montagu, 1821). In: Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals* (Vol. 6: The Second Book of Dolphins and the Porpoises, pp. 137-182). San Diego, CA: Academic Press.
- Wells, R. S., & Scott, M. D. (2002). Bottlenose Dolphins, *Tursiops truncatus* and *T. aduncus*. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 122-128). San Diego, CA: Academic Press.

- Westgate, A. J., & Read, A. J. (1998). Applications of New Technology to the Conservation of Porpoises. *Marine Technology Science Journal*, 32(1), 70-81.
- Westgate, A. J., Read, A. J., Berggren, P., Koopman, H. N., & Gaskin, D. E. (1995). Diving Behaviour of Harbour Porpoises, *Phocoena phocoena*. *Canadian Journal of Fisheries and Aquatic Sciences*, 52, 1064-1073.
- Whitehead, H. (2002). Sperm Whale. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1165-1172). San Diego, CA: Academic Press.
- Whitehead, H., & Wimmer, T. (2005). Heterogeneity and the Mark-Recapture Assessment of the Scotian Shelf Population of Northern Bottlenose Whales (*Hyperoodon ampullatus*). *Canadian Journal of Fisheries and Aquatic Science*, 62, 2573-2585.
- Willis, P. M., & Baird, R. W. (1998). Status of the Dwarf Sperm Whale, *Kogia simus*, with Special Reference to Canada. *The Canadian Field-Naturalist*, 112(1), 114-125.
- Witzell, W. N. (1983). *Synopsis of Biological Data on the Hawksbill Turtle Eretmochelys imbricata (Linnaeus, 1766), Final*. (FAO Fisheries Synopsis No. 137). Miami, FL: National Oceanic and Atmospheric Administration, National Marine Fisheries Service. p. 81.
- Womble, J. N., Blundell, G. M., Gende, S. M., Horning, M., Sigler, M. F., & Csepp, D. J. (2014). Linking marine predator diving behavior to local prey fields in contrasting habitats in a subarctic glacial fjord. *Marine Biology*, 161, 1361-1374.
- Woodward, B. (2006). *Locomotory Strategies, Dive Dynamics, and Functional Morphology of the Mysticetes: Using Morphometrics, Osteology, and DTAG Data to Compare Swim Performance in Four Species*. (Doctor of Philosophy), The University of Maine, Orono, ME.
- Woodward, B. L., & Winn, J. P. (2006). Apparent Lateralized Behavior in Gray Whales Feeding off the Central British Columbia Coast. *Marine Mammal Science*, 22(1), 64-73.
- Würsig, B., Wells, R. S., & Croll, D. A. (1986). Behavior of Gray Whales Summering near St. Lawrence Island, Bering Sea. *Canadian Journal of Zoology*, 64, 611-621.
- Yamada, T. K. (2002). *On an Unidentified Beaked Whale Found Stranded in Kagoshima*. Tokyo, Japan: National Science Museum, Tokyo. p. 5.
- Young, R. E. (1978). Vertical distribution and photosensitive vesicles of pelagic cephalopods from Hawaiian waters. *Fisheries Bulletin*, 76, 583-615.
- Zerbini, A. N., & de Oliveira Santos, M. C. (1997). First Record of the Pygmy Killer Whale *Feresa attenuata* (Gray, 1874) for the Brazilian Coast. *Aquatic Mammals*, 23(2), 105-109.